

Biowell Fasmol 250mg/5ml Suspension

Paracetamol (250mg/5ml)

What is in this leaflet

1. What Biowell Fasmol 250mg/5ml Suspension is used for
2. How Biowell Fasmol 250mg/5ml Suspension works
3. Before you use Biowell Fasmol 250mg/5ml Suspension
4. How to use Biowell Fasmol 250mg/5ml Suspension
5. While you are using it
6. Side effects
7. Storage and disposal of Biowell Fasmol 250mg/5ml Suspension
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Biowell Fasmol 250mg/5ml Suspension is used for

Indicated for relief of fever and pain (such as headache and toothache) in children.

How Biowell Fasmol 250mg/5ml Suspension works

Paracetamol acts by inhibiting the production of hormone-like substances that participate in a wide range of body functions such as the contraction and relaxation of muscle, the dilation and constriction of blood vessels, control of blood pressure and modulation of inflammation. It is considered to affect the brain that controls body temperature.

Before you use Biowell Fasmol 250mg/5ml Suspension

When you must not use it

Tell your doctor if you are hypersensitive to Paracetamol.

- Before you start to use it

Pregnancy and lactation

Pregnancy: Not known to be harmful
Lactation: Amount too small to be harmful.

Ask your doctor or pharmacists for advice before taking Biowell Fasmol 250mg/5ml Suspension

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Paracetamol should not be taken concomitantly with

- Alcohol
- Cholestyramine (reduce cholesterol)
- Warfarin (prevent blood clot)

- Metoclopramide or Domperidone (treat nausea and vomiting)

Paracetamol should not be taken concomitantly with a medicine that acts on the liver.

How to use Biowell Fasmol 250mg/5ml Suspension

How much to use

Children aged 1 - 5 years: ½ to 1 teaspoonful (2.5 ml to 5 ml).

6 - 12 years: 1 to 2 teaspoonfuls (5 ml to 10 ml).

Doses may be repeated every 4 – 6 hours when necessary (maximum of 4 doses in 24 hours).

The directions given may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking Biowell Fasmol 250mg/5ml Suspension for as long as your doctor recommends.

Do not exceed the recommended dosage or use for more than 10 days unless prescribed.

If you forget to use it

Take the missed dose as soon as you remember. If it almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Features of Paracetamol poisoning are nausea and vomiting, followed by pain and tenderness in the right side of lower ribs area of the abdomen. Overdose can cause liver damage that is signified by yellowing of the skin and eyes, gastrointestinal bleeding and low blood glucose level.

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell the doctor, dentist and pharmacist treating you that you are taking Biowell Fasmol 250mg/5ml Suspension.

- Things you must not do

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Biowell Fasmol 250mg/5ml Suspension to anyone else, even if they have the same symptoms or condition as you.

This preparation contains PARACETAMOL. Do not take any other PARACETAMOL-containing medicines at the same time.

- Things to be careful of

This medicine should be used with care in liver and kidney impairment.

Side effects

Side effects of the Biowell Fasmol 250mg/5ml Suspension usually mild. Nausea, gastro-intestinal upsets, rashes and allergic conditions have occurred.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my (Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal of Biowell Fasmol 250mg/5ml Suspension

Keep out of reach of children.

Shake well before use.

Store below 30°C. Protect from light.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of

Biowell Fasmol 250mg/5ml Suspension

Paracetamol (250mg/5ml)

medicines no longer required. These measures will help to protect the environment.

Product Description

- *What it looks like*

Pink colour suspension, opaque and with raspberry flavour.

- *Ingredients*

- Active ingredients

Paracetamol

- Inactive ingredients

Raspberry Flavour, Red (Carmoisine), Propyl Paraben, Sorbitol 70% - Non Crystal, Xanthan Gum, Gum Tragacanth, Methyl Paraben, Sodium Saccharin, Sucrose (Refined Sugar) & Purified Water.

- *MAL number*

MAL

Product Registration Holder

SSJ PHARMA SDN BHD
658-D, Jalan Bukit Melaka 1/1
Taman Bukit Melaka,
Bukit Beruang, Bukit Baru,
75450 Melaka Tengah,
Melaka, Malaysia.

Manufacturer

Dynapharm (M) Sdn. Bhd.
2497, Mk 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3, 13600
Perai, Pulau Pinang, Malaysia.

Date of revision:

27/03/2026, rev 01

Serial Number:

NPRA (R2) 26/039

Biowell Fasmol 250mg/5ml Suspension

Paracetamol (250mg/5ml)
