

# REDISTRA (IMATINIB TABLETS 100MG)

# REDISTRA (IMATINIB TABLETS 400MG)

## *Imatinib (100mg and 400mg)*

### What is in this leaflet

1. What Redistra is used for
2. How Redistra works
3. Before you use Redistra
4. How to use Redistra
5. While you are using it
6. Side effects
7. Storage and Disposal of Redistra
8. Product Description
9. Manufacturer
10. Product Registration Holder
11. Date of revision
12. Serial Number

### What Redistra is used for

Redistra is a medicine containing an active substance called imatinib. Redistra is used in the treatment for adult and children for:

- Chronic myeloid leukaemia (CML). Leukaemia is a cancer of white blood cells. These white cells usually help the body to fight infection. Chronic myeloid leukaemia is a form of leukaemia in which certain abnormal white cells (named myeloid cells) start growing out of control.
- Philadelphia chromosome positive acute lymphoblastic leukaemia (Ph-positive ALL). Leukaemia is a cancer of white blood cells. These white cells usually help the body to fight infection. Acute lymphoblastic leukaemia is a form of leukaemia in which certain abnormal white cells (named lymphoblasts) start growing out of control. Redistra inhibits the growth of these cells

Redistra is also used in the treatment for adult for:

- Myelodysplastic/myeloproliferative diseases (MDS/MPD). These are a group of blood diseases in which some blood cells start growing out of control. Redistra inhibits the growth of these cells in a certain subtype of these diseases
- Hypereosinophilic syndrome (HES) and/or chronic eosinophilic leukaemia (CEL). These are blood diseases in which some blood cells (named eosinophils) start growing out of control. Redistra inhibits the growth of these cells in a certain

subtype of these diseases.

- Gastrointestinal stromal tumours (GIST). GIST is a cancer of the stomach and bowels. It arises from uncontrolled cell growth of the supporting tissues of these organs.
- Dermatofibrosarcoma protuberans (DFSP). DFSP is a cancer of the tissue beneath the skin in which some cells start growing out of control. Redistra inhibits the growth of these cells
- Aggressive systemic mastocytosis (ASM). ASM is a rare condition where certain white blood cells (mast cells) build up in certain tissue and organs in the body.

In the rest of this leaflet, we will use the abbreviations when talking about these diseases. If you have any questions about how Redistra works or why this medicine has been prescribed for you, ask your doctor.

### How Redistra works

This medicine works by inhibiting the growth of abnormal cells in the diseases listed above. These include some types of cancer.

### Before you use Redistra

#### - When you must not use it

If you are allergic to *imatinib* or any of the other ingredients of this medicine listed in the 'inactive ingredient' below.

#### - Before you start to use it

Talk to your doctor before taking Redistra:

- if you have or have ever had a liver, kidney or heart problem.
- if you are taking the medicine levothyroxine because your thyroid has been removed.
- if you have ever had or might now have a hepatitis B infection. This is because Redistra could cause hepatitis B to become active again, which can be fatal in some cases. Patients will be carefully checked by their doctor for signs of this infection before treatment is started.
- if you experience bruising, bleeding, fever, fatigue and confusion when taking Redistra, contact your doctor.

This may be a sign of damage to blood vessels known as thrombotic microangiopathy (TMA).

If any of these apply to you, tell your doctor before taking Redistra.

#### *Pregnancy, breast-feeding and fertility*

- If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.
- Redistra is not recommended during pregnancy unless clearly necessary as it may harm your baby. Your doctor will discuss with you the possible risks of taking Redistra during pregnancy.
- Women who might become pregnant are advised to use effective contraception during treatment and for 15 days after ending treatment.
- Do not breast-feed during the treatment with Redistra and for 15 days after ending treatment, as it may harm your baby.
- Patients who are concerned about their fertility while taking Redistra are advised to consult with their doctor.

#### - Taking other medicines

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription and including herbal medicines. Some medicines can interfere with the effect of Redistra when taken together. They may increase or decrease the effect of Redistra, either leading to increased side effects or making Redistra less effective. Redistra may do the same to some other medicines.

Some medication will cause complications when taken together with Redistra, such as:

- CYP3A4 inhibitors (e.g. protease inhibitors such as indinavir, lopinavir/ritonavir, ritonavir, saquinavir, telaprevir, nelfinavir, boceprevir; azole antifungals including ketoconazole, itraconazole, posaconazole, voriconazole; certain macrolides such as erythromycin, clarithromycin and telithromycin) might cause decreased metabolism and increased imatinib concentration in the

# REDISTRA (IMATINIB TABLETS 100MG)

# REDISTRA (IMATINIB TABLETS 400MG)

## Imatinib (100mg and 400mg)

body.

- Inducers of CYP3A4 activity (e.g. dexamethasone, phenytoin, carbamazepine, rifampicin, phenobarbital, fosphenytoin, primidone or St. John's Wort) may significantly reduce exposure to Redistra, potentially increasing the risk of Imatinib to lose its efficacy.
- Redistra may increase concentration of other CYP3A4 metabolised drugs (e.g. triazolobenzodiazepines, dihydropyridine calcium channel blockers, certain HMG-CoA reductase inhibitors, i.e. statins, etc.)
- Narrow therapeutic medications (e.g. cyclosporine, pimozone), where Imatinib might increase the plasma level of these medications.
- Warfarin, as it may cause the increase in risk of bleeding.
- Paracetamol
- Levothyroxine, as Imatinib might cause this medication to reduce its effectiveness.

### **How to use Redistra**

#### - How much to use

##### *Use in adults*

Your doctor will tell you exactly how many tablets of Redistra to take.

- 400 mg to be taken as one tablet once a day
- 600 mg to be taken as one tablet of 400 mg plus 2 tablets of 100 mg once a day
- 800 mg to be taken as one tablets of 400mg twice a day

#### If you are being treated for CML:

Depending on your condition the usual dose is either 400 mg or 600 mg and your doctor may decide to increase the dose increased up to 800 mg

#### If you are being treated for GIST:

The starting dose is 400 mg. Your doctor may decide to increase the dose to 600mg or 800mg

#### If you are being treated for Ph-positive ALL:

The starting dose is 600 mg.

#### If you are being treated for MDS/MPD:

The starting dose is 400 mg.

#### If you are being treated for HES/CEL:

The starting dose is 100 mg. Your doctor may decide to increase the dose to 400 mg.

#### If you are being treated for ASM:

The starting dose is 100 mg. Your doctor may decide to increase the dose to 400 mg.

#### If you are being treated for DFSP:

The dose is 800 mg per day

#### *Use in children and adolescents*

The doctor will tell you how many tablets of Redistra to give to your child. The amount of Redistra given will depend on your child's condition, body weight and height. The total daily dose in children must not exceed 600 mg with Ph+ALL and CML. The treatment can either be given to your child as a once-daily dose or alternatively the daily dose can be split into two administrations (half in the morning and half in the evening).

#### - When to use it

- Take Redistra with a meal. This will help protect you from stomach problems when taking Redistra.

- Swallow the tablets whole with a large glass of water.

If you are unable to swallow the tablets, you can dissolve them in a glass of still water or apple juice:

- Use about 50ml for each 100mg tablet and 200 ml for each 400 mg tablet.
- Stir with a spoon until the tablets have completely dissolved.
- Once the tablet has dissolved, drink everything in the glass straight away.

#### - How long to use it

Keep taking Redistra every day for as long as your doctor tells you.

#### - If you forget to use it

- If you forget a dose, take it as soon as you remember. However, if it is nearly time for the next dose, skip the

missed dose.

- Then continue with your normal schedule.

- Do not take a double dose to make up a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

#### - If you use too much (overdose)

If you have accidentally taken too many tablets, talk to your doctor straight away. You may require medical attention. Take the medicine pack with you

### **While you are using it**

#### - Things you must do

- You may become more sensitive to the sun while taking Redistra. It is important to cover sun-exposed areas of skin and use sunscreen with high sun protection factor (SPF). These precautions are also applicable to children.

- During treatment with Redistra, tell your doctor straight away if you put on weight very quickly. Redistra may cause your body to retain water (severe fluid retention).

- While you are taking Redistra, your doctor will regularly check whether the medicine is working. You will also have blood tests and be weighed regularly.

#### - Children and adolescents

Redistra is also a treatment for children with CML. There is no experience in children with CML below 2 years of age. There is limited experience in children with Ph-positive ALL and very limited experience in children with MDS/MPD, DFSP, GIST and HES/CEL. Some children and adolescents taking Redistra may have slower than normal growth. The doctor will monitor the growth at regular visits.

#### - Things you must not do

- Do not stop taking your medicines without consulting your doctor.

- Do not use Redistra after the expiry date which is stated on the carton and bottle.

- Do not give this medication to anyone else, even if they have the same

**REDISTRA (IMATINIB TABLETS 100MG)****REDISTRA (IMATINIB TABLETS 400MG)*****Imatinib (100mg and 400mg)***

condition as you.

***- Things to be careful of******Driving and using Machines***

You may feel dizzy or drowsy or get blurred vision while taking this medicine. If this happens, do not drive or use any tools or Machines until you are feeling well again.

**Side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them. They are usually mild to moderate.

Some side effects may be serious. Tell your doctor straight away if you get any of the following:

***Very common or common:***

- Rapid weight gain. Redistra may cause your body to retain water (severe fluid retention).
- Signs of infection such as fever, severe chills, sore throat or mouth ulcers. Redistra can reduce the number of white blood cells, so you might get infections more easily.
- Unexpected bleeding or bruising (when you have not hurt yourself).

If you get any of the above, tell your doctor straight away.

Other side effects may include:

Very common:

- Neutropenia (low level of white blood cells), thrombocytopenia (low level of platelet), anaemia (low level of red blood cells)
  - abdominal pain
  - Headache or feeling tired.
  - Feeling sick (nausea), being sick (vomiting), diarrhoea or indigestion.
  - Rash.
  - Muscle cramps or joint, muscle or bone pain, during Redistra treatment or after you have stopped taking Redistra.
  - Swelling such as round your ankles or puffy eyes.
  - Weight gain.
- If any of these affects you severely, tell your doctor.

Common:

- Pancytopenia (low level of red blood cell, white blood cell & platelet)
  - febrile neutropenia (fever in patients with low level of white blood cells)
  - Paresthesia (abnormal sensation of the skin)
  - hypoaesthesia (partial or total loss of sensation in a part of your body)
  - Swelling of the eyelid
  - Hemorrhage (blood loss)
  - Cough
  - Increase in liver enzymes level
  - Swelling of the face
  - Anasarca (general swelling of the body)
  - Raised body temperature
  - chills, Rigor
  - Anorexia, weight loss or a disturbed sense of taste.
  - Feeling dizzy or weak.
  - Difficulty in sleeping (insomnia).
  - Discharge from the eye with itching, redness and swelling (conjunctivitis), watery eyes or having blurred vision.
  - Nose bleeds.
  - Pain or swelling in your abdomen, flatulence, heartburn or constipation.
  - Itching.
  - Unusual hair loss or thinning.
  - Numbness of the hands or feet.
  - Mouth ulcers.
  - Joint pain with swelling.
  - Dry mouth, dry skin or dry eye.
  - Decreased or increased skin sensitivity.
  - Hot flushes, chills or night sweats.
- If any of these affects you severely, tell your doctor.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) (Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

**Storage and Disposal of Redistra*****- Storage***

Keep out of the reach and sight of children. Do not store above 30 °C.

***- Disposal***

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

**Product Description*****- What it looks like***

REDISTRA (Imatinib Tablets 100 mg): Yellow to brownish orange, round film coated tablets debossed with “1” on one side and score line on the other side

REDISTRA (Imatinib Tablets 400 mg): Yellow to brownish orange, ovaloid film coated tablets debossed with “4” on one side and score line on the other side.

***- Ingredients******- Active ingredient***

REDISTRA (Imatinib Tablets 400 mg): *Imatinib (as mesilate)* 100mg

REDISTRA (Imatinib Tablets 400 mg): *Imatinib (as mesilate)* 400mg

***- Inactive ingredients***

Sodium Stearyl Fumarate, Colloidal Silica Anhydrous, Crospovidone, Microcrystalline cellulose, Lactose anhydrous, magnesium stearate & Opadry Brown

***- MAL number:*****Manufacturer**

Dr. Reddy's Laboratories Ltd.  
Formulation Unit -VII,  
Plot Nos, P1 to P9, Phase III,  
VSEZ, Duvvada, Visakhapatnam District  
-530046, Andhra Pradesh, India.

**Product Registration Holder**

Dr. Reddy's Laboratories Malaysia Sdn. Bhd. (1238154-X)  
Unit 10-06, Menara MBMR  
No. 1, Jalan Syed Putra,  
58000 Kuala Lumpur, Malaysia

**Date of revision**

10/02/2021

**Serial Number**

NPRA (R2) 21/004