

WELLMEX ATOMIC ENEMA 原子浣腸

ACTIVE INGREDIENT

Each 100ml contains :

Sodium chloride.....	15g
Preservative : Methylparaben	0.05g/100ml
Sodium Benzoate	0.05g/100ml

PHARMACODYNAMICS

Sodium salt act by retaining fluid in the bowel by osmosis or by changing the pattern of water distribution in faeces.

PHARMACOKINETICS

Sodium chloride is readily absorbed from the gastrointestinal tract. It is present in all the body fluids and is mainly found in the extra cellular fluid. The amount of sodium chloride lost in sweat is small and the osmotic equilibrium is maintained by the excretion of surplus amounts in the urine.

INDICATION

It is used for relieving constipation.

For rectally use only.

ADVERSE EFFECTS

Enema can cause rectal irritations, which are characterized by rectal bleeding, blistering, burning, itching or pain. There can also be skin irritation surrounding the rectal area.

WARNING/PRECAUTION

Enemas should be used with caution in the elderly and should not be used by people with acute gastrointestinal conditions or inflammatory bowel disease. Enemas should not be used in patients with congenital megacolon, bowel obstruction, imperforate anus or congestive heart failure. Use with caution in patients with impaired renal function, pre-existing electrolyte disturbances or in patients on diuretics or other medications which may affect electrolyte levels or where colostomy exists. Since enema contains sodium chloride, there is a risk of elevated serum levels of sodium and decreased levels of potassium and consequent hypernatremia. This is of particular concern in children with megacolon or any other condition where there is retention of enema solution.

CONTRAINDICATIONS

It is contraindicated in patients with ileus, gastric retention, gastrointestinal obstruction, bowel perforation, toxic colitis and toxic megacolon.

DRUG INTERACTION

Not known.

DOSAGE & ADMINISTRATION

Rectally used. Adults : 20-40ml, once daily.

Children : 10-20ml, once daily.

SYMPTOMS OF OVERDOSAGE & TREATMENT

The sign and symptoms would be the exaggeration of the side effects, especially rectal irritation. The frequency of bowel movement would also be increased. Treatment would be the cessation of use of the medication. If symptoms continue, professional medical advice should be obtained.

STORAGE

Store below 30°C in a cool dry place and protected from sunlight. Keep out of reach of children.

SHELF-LIFE

See the instructions on the package.

PACK SIZE

10ml/bottle & 20ml/bottle.

PRODUCT DESCRIPTION

A colorless, clear and transparent solution.

Product Registration Holder:

NGL TRADING SDN. BHD., (1020904-A),
4-1-5, Kompleks Kantonmen Prima,
Batu 4 ½, Jalan Ipoh, 51200 Kuala Lumpur,
Wilayah Persekutuan Kuala Lumpur,

Manufactured by:

HEALTH CHEMICAL PHARMACEUTICAL
CO., LTD.
No.12 Yeou-4th Rd., Ta-Chia Yowshy Ind. Dist,
Taichung City, Taiwan, R.O.C

Date of revision: 6th November 2025

13

品名：原子浣腸說明書(馬來西亞)

尺寸：130x150mm

材質：模造紙60g

印刷：單面印刷 黑刷

青水印刷113. 04. 19

版次：13