

DIFFLAM HEXTRA™ SORE THROAT LOZENGES 2.4MG

Hexylresorcinol (2.4mg)

What is in this leaflet

1. What is Difflam HEXTRA Sore Throat Lozenges 2.4mg is used for
2. How Difflam HEXTRA Sore Throat Lozenges 2.4mg works
3. Before you use Difflam HEXTRA Sore Throat Lozenges 2.4mg
4. How to use Difflam HEXTRA Sore Throat Lozenges 2.4mg
5. While you are using it
6. Side effects
7. Storage and Disposal of Difflam HEXTRA Sore Throat Lozenges 2.4mg
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What is Difflam HEXTRA Sore Throat Lozenges 2.4mg is used for

Difflam HEXTRA Sore Throat Lozenges 2.4mg is used as an antiseptic and local anaesthetic for the relief of sore throat and its associated pain.

How Difflam HEXTRA Sore Throat Lozenges 2.4mg works

Difflam HEXTRA Sore Throat Lozenges 2.4mg contain Hexylresorcinol which is an anaesthetic to bring soothing relief of pain and discomfort.

Before you use Difflam HEXTRA Sore Throat Lozenges 2.4mg

-When you must not use it

Do not use this medicine if you are hypersensitive to any of the ingredients in this product.

Please do not take this medicine if you have hereditary fructose intolerance, glucose-galactose malabsorption syndrome or sucrose-isomaltase deficiency.

If you are unsure about this, kindly ask your doctor or pharmacist for advice.

-Before you start to use it

Please read all instructions carefully before using this product. Full details are available in the package insert of this product. If you are unsure about any information, kindly seek advice from your doctor or pharmacist before use.

-Taking other medicines

It is not known if this medicine interacts significantly with other medications. However, if you are currently on any medications, whether it is bought over the counter, without a prescription from the pharmacy or from a health food store, please consult your doctor or pharmacist before use.

How to use Difflam HEXTRA Sore Throat Lozenge 2.4mg

-How much to use

Please follow instructions carefully as stated in the product's package insert. If you are unsure, kindly seek advice from your doctor or pharmacist.

Adults & Children 6 years & over:
1 lozenge to be dissolved slowly in the mouth every 3 hours as required up to a maximum of 12 lozenges per day.

Children under 6 years:
Not recommended.

-When to use it

Use as directed by your doctor or pharmacist.

-How long to use it

Use this medicine as directed. If your symptoms persist, have not improved or have worsened, please consult your doctor or pharmacist for advice.

-If you forgot to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Gastrointestinal discomfort is a symptom of overdosage of this medicine.

While you are using it

-Things you must do

Take this medicine exactly as instructed.

Tell all the doctors and pharmacists treating you that you are taking Difflam HEXTRA Sore Throat Lozenges 2.4mg.

-Things you must not do

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Difflam HEXTRA Sore Throat Lozenges 2.4mg to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of**Driving and using machines:**

This medicine has no or negligible influence on the ability to drive or use machines.

Pregnancy and lactation:

There is lack of evidence of safety of the product in human pregnancy and in animals, but hexylresorcinol has been used widely in lozenges for many years without apparent ill consequence. However, as with all medicines, caution should be exercised during pregnancy and breastfeeding.

Side effects

Like all medicines, Difflam HEXTRA Sore Throat Lozenges 2.4mg can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npdra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and disposal of Difflam Forte Anti-inflammatory Throat Spray**-Storage**

Store below 30°C.

Keep out of reach of children.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.

Product description**-What it looks like**

Purple coloured, round, flat, bevelled shaped lozenges.

-Ingredients**-Active ingredient(s)**

Hexylresorcinol

-Inactive ingredients

- Levomenthol
- Liquid Glucose
- Sucrose
- Sucralose
- Anhydrous Citric Acid
- Purple Grape Flavour,
- FD & C Red No.40
- FD & C Blue No. 1
- Purified Water

-MAL Number

MAL20076056X

Manufacturer

Unique Pharmaceutical Laboratories
(A Div of J. B. Chemicals & Pharmaceuticals Ltd.)
Survey No. 101/2 & 102/1,
Daman Industrial Estate,
Airport Road, Kadaiya, Daman,
396 210, India.

Product Registration Holder

iNova Pharmaceuticals (Singapore) Pte
Ltd (Incorporated in Singapore)
Malaysia Branch
5F-2, 5th Floor,
Tower 2, @PFCC,
Jalan Puteri 1/2, Bandar Puteri,
47100 Puchong,
Selangor, Malaysia

Date of revision

05/ 03/ 2025

Serial Number

NPRA (R2) 20/019