

TOKUSHIN MEDICAL PLASTER

Menthol (6.50%w/w), Methyl Salicylate (5.00%w/w), Camphor (1.25%w/w)

What is in this leaflet?

1. What TOKUSHIN MEDICAL PLASTER is used for
2. How TOKUSHIN MEDICAL PLASTER works
3. Before you use TOKUSHIN MEDICAL PLASTER
4. How to use TOKUSHIN MEDICAL PLASTER
5. While you are using it
6. Side effects
7. Storage and Disposal of TOKUSHIN MEDICAL PLASTER
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What TOKUSHIN MEDICAL PLASTER is used for

For symptomatic treatment of rheumatism, neuralgia, stiff shoulder, bruise, sprain, and all other muscular aches.

How TOKUSHIN MEDICAL PLASTER works

TOKUSHIN MEDICAL PLASTER contains active ingredients Menthol, Methyl Salicylate, and Camphor. Menthol, when applied to skin, dilates the blood vessels causing a sensation of coldness followed by an analgesic effect. Applied externally. Methyl Salicylate is used to relieve pain in musculoskeletal, joint, and soft-tissue disorders. Camphor acts as a rubefacient and mild analgesic.

Before you use TOKUSHIN MEDICAL PLASTER

When you must not use it

Do not use TOKUSHIN MEDICAL PLASTER if

1. You are taking warfarin, excessive application onto the skin for muscles or joint pains may increase the chances of bleeding
2. You have skin sensitization or contact dermatitis.
3. The application area is directly into the nostrils

Before you start to use it

Check with your doctor or pharmacist
If you are allergic to Methyl Salicylate, Menthol, Camphor, or to any other ingredients in this product.

Taking other medicines

Tell your doctor if you are currently taking warfarin.

How to use TOKUSHIN MEDICAL PLASTER

How much to use

Adult: 1 to 2 times a day
Remove after at most 8 hours.

When to use it

Use to provides for relief of aches and pains associated with: rheumatism, neuralgia, stiff shoulder, bruise, sprain, and all other muscular aches.

How long to use it

Usage should continue when there are signs of rheumatism, neuralgia, stiff shoulder, bruise, sprain, and all other muscular aches.

Consult a doctor or pharmacist if the pain does not improve after 5 to 6 days of use.

If you forgot to use it

Use the product as soon as possible and continue to use as normal usage.

If you use too much (overdose)

For user taking warfarin, excessive application on to the skin for muscle and joint pains may increase chances of bleeding.

If you use too much of this product, consult your doctor or pharmacist.

While you are using it

Things you must do

Keep the applied area from getting wet.

Things you must not do

This product should not be used on infants below 2 years of age and is contraindicated in those with sensitivity to any of the ingredients of the product.

Things to be careful of

For external use only.
Keep out of reach of children.

Side effects

Do not use if you are allergic to aspirin or salicylate. Stop use and ask a doctor if condition worsen, rash, itching, or excessive skin irritation occur.

You may report any side effects or adverse drug reactions directly to the National Center for Adverse Drug Reaction Monitoring by visiting the website www.npra.gov.my [Consumers -> Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of TOKUSHIN MEDICAL PLASTER

Storage

Store at below 30°C.
Protect from light and moisture.

Disposal

Dispose the product after the expired date.

Product Description

What it looks like TOKUSHIN MEDICAL PLASTER

Rectangular fabric plaster coated with a thin layer of medicated colloid.

Ingredients

The ingredients are Menthol, Methyl Salicylate, Camphor, Liquid Paraffin, Rubber RSS1, WingStay L, Polyisobutene, Zinc Oxide, Hydrocarbon Resin, Polybutene and Lanolin Anhydrous.

MAL number

Manufacturer & Product Registration Holder

THP Medical Sdn. Bhd (1135528-K)

Lot PT 1281 (Plot 308), Jalan Perindustrian Bukit Minyak 18, Kawasan Perindustrian Bukit Minyak, 14100 Simpang Ampat, Pulau Pinang, Malaysia.

Date of Revision

01/10/2024

Serial Number

NPRA(R2)24/110