

# THINHET 60 FILM-COATED TABLETS

Ticagrelor 60 mg

## What is in this leaflet

1. What THINHET 60 is used for
2. How THINHET 60 works
3. Before you use THINHET 60
4. How to use THINHET 60
5. While you are using it
6. Side effects
7. Storage and Disposal of THINHET 60
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision
11. Serial Number

## What THINHET 60 is used for

THINHET 60 contains an active substance called ticagrelor. This belongs to a group of medicines called anti-platelet medicines.

THINHET 60 in combination with acetylsalicylic acid (another antiplatelet agent) is to be used in adults only. You have been given THINHET 60 because you have:

- Had a heart attack, over a year ago.
- coronary artery disease and type 2 diabetes, with no previous heart attacks, and have previously had a procedure to open a blocked or narrowed heart blood vessel (percutaneous coronary intervention or PCI). This may have included placement of a small tube (stent) to keep the blocked or narrowed blood vessel open.

It reduces the chances of you having another heart attack, stroke or dying from a disease related to your heart or blood vessels.

## How THINHET 60 works

THINHET 60 affects cells called 'platelets' (also called thrombocytes). These very small blood cells help stop bleeding by clumping together to plug tiny holes in blood vessels that are cut or damaged.

However, platelets can also form clots inside diseased blood vessels in the

heart and brain. This can be very dangerous because:

- the clot can cut off the blood supply completely; this can cause a heart attack (myocardial infarction) or stroke, or
- the clot can partly block the blood vessels to the heart; this reduces the blood flow to the heart and can cause chest pain which comes and goes (called 'unstable angina').

THINHET 60 helps stop the clumping of platelets. This reduces the chance of a blood clot forming that can reduce blood flow.

## Before you use THINHET 60

### -When you must not use it

- You are allergic to ticagrelor or any of the other ingredients of this medicine (listed in section "Ingredients").
- You are bleeding now.
- You have had a stroke caused by bleeding in the brain.
- You have severe liver disease.
- You are taking any of the following medicines:
  - ketoconazole (used to treat fungal infections)
  - clarithromycin (used to treat bacterial infections)
  - nefazodone (an antidepressant)
  - ritonavir and atazanavir (used to treat HIV infection and AIDS).

Do not take THINHET 60 if any of the above applies to you. If you are not sure, talk to your doctor or pharmacist before taking this medicine.

### -Before you start to use it

Talk to your doctor or pharmacist before taking THINHET 60 if:

- You have an increased risk of bleeding because of:
  - a recent serious injury
  - recent surgery (including dental work, ask your dentist

about this)

- you have a condition that affects blood clotting
- recent bleeding from your stomach or gut (such as a stomach ulcer or colon 'polyps')
- You are due to have surgery (including dental work) at any time while taking THINHET 60. This is because of the increased risk of bleeding. Your doctor may want you to stop taking this medicine 5 days prior to surgery.
- Your heart rate is abnormally low (usually lower than 60 beats per minute) and you do not already have in place a device that paces your heart (pacemaker).
- You have asthma or other lung problems or breathing difficulties.
- You develop irregular breathing patterns such as speeding up, slowing down or short pauses in breathing. Your doctor will decide if you need further evaluation.
- You have had any problems with your liver or have previously had any disease which may have affected your liver.
- You have had a blood test that showed more than the usual amount of uric acid.

If any of the above apply to you (or you are not sure), talk to your doctor or pharmacist before taking this medicine. If you have coronary artery disease and type 2 diabetes, your doctor should regularly monitor if your treatment is appropriate for you. Your doctor will discuss any risks and ways to prevent possible side effects while you are taking THINHET 60.

If you are taking both THINHET 60 and heparin:

- Your doctor may require a sample of your blood for diagnostic tests if they suspect a rare platelet disorder caused by heparin. It is important

# THINHET 60 FILM-COATED TABLETS

Ticagrelor 60 mg

that you inform your doctor that you are taking both THINHET 60 and heparin, as THINHET 60 may affect the diagnostic test.

## Use in children

THINHET 60 is not recommended for children and adolescents under 18 years.

## -Taking other medicines

Please tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This is because THINHET 60 can affect the way some medicines work, and some medicines can have an effect on THINHET 60.

Tell your doctor or pharmacist if you are taking any of the following medicines:

- more than 40 mg daily of either simvastatin or lovastatin (medicines used to treat high cholesterol)
- rifampicin (an antibiotic)
- phenytoin, carbamazepine and phenobarbital (used to control seizures)
- digoxin (used to treat heart failure)
- cyclosporine (used to lessen your body's defenses)
- quinidine and diltiazem (used to treat abnormal heart rhythms)
- beta blockers and verapamil (used to treat high blood pressure).
- morphine and other opioids (used to treat severe pain)

In particular, tell your doctor or pharmacist if you are taking any of the following medicines that increase your risk of bleeding:

- 'oral anticoagulants' often referred to as 'blood thinners' which include warfarin.
- Non-Steroidal Anti-Inflammatory Drugs (abbreviated as NSAIDs) often taken as painkillers such as ibuprofen and naproxen.
- Selective Serotonin Reuptake

Inhibitors (abbreviated as SSRIs) taken as antidepressants such as paroxetine, sertraline and citalopram.

- other medicines such as ketoconazole (used to treat fungal infections), clarithromycin (used to treat bacterial infections), nefazodone (an antidepressant), ritonavir and atazanavir (used to treat HIV infection and AIDS), cisapride (used to treat heartburn), ergot alkaloids (used to treat migraines and headaches).

Also tell your doctor that because you are taking THINHET 60, you may have an increased risk of bleeding if your doctor gives you fibrinolytics, often called 'clot solvers', such as streptokinase or alteplase.

## **How to use THINHET 60**

### -How much to use

The usual dose is one tablet of 60 mg twice a day. Your doctor will usually also tell you to take acetylsalicylic acid. This is a substance present in many medicines used to prevent blood clotting. Your doctor will tell you how much to take (usually between 75-150 mg daily).

### -When to use it

Take this medicine around the same time every day (for example, one tablet in the morning and one in the evening). You can take the tablet with or without food.

If you have trouble swallowing the tablet you can crush it and mix with water as follows:

- Crush the tablet to a fine powder.
- Pour the powder into half a glass of water.
- Stir and drink immediately.
- To make sure there is no medicine left, rinse the empty glass with another half a glass of water and drink it.

If you are in the hospital you may be given this tablet mixed with some water and given through a tube via the nose (nasogastric tube).

### -How long to use it

Continue taking THINHET 60 as long as your doctor tells you.

### -If you forget to use it.

- If you forget to take a dose, just take your next dose as normal.
- Do not take a double dose (two doses at the same time) to make up for the forgotten dose.

### -If you use too much (overdose)

If you take more THINHET 60 than you should, talk to a doctor or go to hospital straight away. Take the medicine pack with you. You may be at increased risk of bleeding.

## **While you are using THINHET 60**

### -Things you must do

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

### -Things you must not do

Do not stop taking THINHET 60 without talking to your doctor. Take this medicine on a regular basis and for as long as your doctor keeps prescribing it. If you stop taking THINHET 60, it may increase your chances of having another heart attack or stroke or dying from a disease related to your heart or blood vessels.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### -Things to be careful of

#### Pregnancy

It is not recommended to use THINHET 60 if you are pregnant or may become pregnant. Women should use appropriate contraceptive measures to avoid pregnancy while taking this

# THINHET 60 FILM-COATED TABLETS

Ticagrelor 60 mg

medicine.

Talk to your doctor before taking this medicine if you are breast-feeding. Your doctor will discuss with you the benefits and risks of taking THINHET 60 during this time. If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

## Driving and using machines

THINHET 60 is not likely to affect your ability to drive or use machines. If you feel dizzy or confusion while taking this medicine, be careful while driving or using machines.

## Side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. The following side effects may happen with this medicine:

THINHET 60 affects blood clotting, so most side effects are related to bleeding. Bleeding may occur in any part of the body. Some bleeding is common (like bruising and nose-bleeds). Severe bleeding is uncommon but can be life threatening.

**See a doctor straight away if you notice any of the following – you may need urgent medical treatment:**

- **Bleeding into the brain or inside the skull is an uncommon side effect, and may cause signs of a stroke such as:**
  - sudden numbness or weakness of your arm, leg or face, especially if only on one side of the body
  - sudden confusion, difficulty speaking or understanding others
  - sudden difficulty in walking or loss of balance or co-ordination
  - suddenly feeling dizzy or sudden severe headache with no known cause

- **Signs of bleeding such as:**
  - bleeding that is severe or that you cannot control
  - unexpected bleeding or bleeding that lasts a long time
  - pink, red, or brown urine
  - vomiting red blood or your vomit looks like ‘coffee grounds’
  - red or black stools (look like tar)
  - coughing up or vomiting blood clots
- **Fainting (syncope)**
  - a temporary loss of consciousness due to sudden drop in blood flow to the brain (common)
- **Signs of a blood clotting problem called Thrombotic Thrombocytopenic Purpura (TTP) such as:**
  - Fever and purplish spots (called purpura) on the skin or in the mouth, with or without yellowing of the skin or eyes (jaundice), unexplained extreme tiredness or confusion

**Discuss with your doctor if you notice any of the following:**

- **Feeling short of breath** - this is very common. It might be due to your heart disease or another cause, or it might be a side effect of THINHET 60. THINHET 60-related breathlessness is generally mild and characterised as a sudden, unexpected hunger for air usually occurring at rest and may appear in the first weeks of therapy and for many may disappear. If your feeling of shortness of breath gets worse or lasts a long time, tell your doctor. Your doctor will decide if it needs treatment or further investigations.

**Other possible side effects: Very common (may affect more than 1 in 10 people)**

- High level of uric acid in your blood (as seen in tests)
- Bleeding caused by blood disorders

**Common (may affect up to 1 in 10 people)**

- Bruising
- Headache
- Feeling dizzy or like the room is spinning
- Diarrhoea or indigestion
- Feeling sick (nausea)
- Constipation
- Rash
- Itching
- Severe pain and swelling in your joints – these are signs of gout
- Feeling dizzy or light-headed, or having blurred vision – these are signs of low blood pressure
- Nosebleed
- Bleeding after surgery or from cuts (for example while shaving) and wounds more than is normal
- Bleeding from your stomach lining (ulcer)
- Bleeding gums

**Uncommon (may affect up to 1 in 100 people)**

- Allergic reaction – a rash, itching, or a swollen face or swollen lips/tongue may be signs of an allergic reaction.
- Confusion
- Visual problems caused by blood in your eye
- Vaginal bleeding that is heavier, or happens at different times, than your normal period (menstrual) bleeding
- Bleeding into your joints and muscles causing painful swelling
- Blood in your ear
- Internal bleeding, this may cause dizziness or light-headedness

# THINHET 60 FILM-COATED TABLETS

Ticagrelor 60 mg

---

## Not known (cannot be estimated from the available data)

- Abnormally low heart rate (usually lower than 60 beats per minute)

If you get any side effects or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist, but do not stop taking THINHET 60 until you have spoken to them.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

## Storage and Disposal of THINHET 60

### -Storage

- Do not store above 30°C.
- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the month and year shown by the numbers following EX (or EXP) on the container. The first two numbers indicate the month; the last two numbers indicate the year.

### -Disposal

Medicines should not be disposed of via wastewater or household water. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## Product Description

### -What it looks like

Peach to light brown, round shaped, biconvex film-coated tablets with approx. diameter of 8.05mm debossed with “68” on one side and “V1” on other side.

Pack size: 3 x 10's Clear  
PVC/PE/PVdC-Aluminium Blister  
Pack

### -Ingredients:

-Active ingredients:

- The active substance is ticagrelor. Each film-coated tablet contains 60 mg of ticagrelor.

-Inactive ingredients:

- The other ingredients are:  
Tablet core: Mannitol (Pearlitol 160 C), Sodium starch Glycolate Type A (Primojel), Povidone (Kollidon 30 LP), Purified water, Mannitol (Pearlitol SD 200), Cellulose Microcrystalline (Emcocel 102), Magnesium stearate (Ligamed MF-2-V)

Film-coating: Instacoat universal Brown (A05E04420), Purified water

### -MAL number:

MAL26036020AZ

### Manufacturer

Annora Pharma Private Limited  
Sy. No. 261, Annaram Village,  
Gummadidala Mandal,  
Sangareddy District, Telangana State,  
502313, India.

### Product Registration Holder

Camber Laboratories Sdn. Bhd  
Unit E-13A-02, Menara SUEZCAP 2,  
No. 2, KL Gateway, Jalan Kerinchi,  
Gerbang Kerinchi Lestari,  
59200 Kuala Lumpur, Malaysia.

### Date of revision

08/01/2026

### Serial Number:

NPRA (R2) 24/128