

BACTOLID (LINEZOLID FILM-COATED TABLETS 600MG)

Linezolid 600 mg

What is in this leaflet

1. What BACTOLID is used for
2. How BACTOLID works
3. Before you use BACTOLID
4. How to use BACTOLID
5. While you are using it
6. Side effects
7. Storage and Disposal of BACTOLID
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What BACTOLID is used for

BACTOLID is an antibiotic. It is used to treat pneumonia, skin infections, and some bacterial infections that are resistant to other antibiotics.

How BACTOLID works

BACTOLID stops or slows the growth of bacterial cells.

Before you use BACTOLID

-When you must not use it

Do not use BACTOLID if you:

- are allergic to *linezolid* or any other ingredients in this product
- are taking or have taken within the last 2 weeks any medicines known as monoamine oxidase inhibitors (e.g. *phenelzine*, *isocarboxazid*)
- have uncontrolled hypertension (high blood pressure)
- have pheochromocytoma (adrenal gland tumor)
- have thyrotoxicosis (excessive thyroid hormone)
- have carcinoid syndrome (signs and symptoms that occur secondary to carcinoid tumours)
- If you are taking any of the following medicines:
 - *Pseudoephedrine*
 - *phenylpropanolamine*
 - *epinephrine*

- *norepinephrine*
- *dopamine*
- *dobutamine*
- serotonin re-uptake inhibitors
- tricyclic antidepressants
- serotonin 5-HT1 receptor agonists
- *meperidine*
- *bupirone*

Pregnancy and lactation

The effect of BACTOLID in pregnant women is not known. Therefore, it should not be taken in pregnancy unless advised by your doctor.

It is not known if linezolid is excreted in human milk. Do not take BACTOLID if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

-Before you start to use it

Talk to your doctor before taking BACTOLID if you:

- are at increased risk for bleeding
- have a pre-existing condition called myelosuppression in which your bone marrow activity is decreased
- are on medications that may decrease hemoglobin levels or platelet count
- have a history of seizures or have risk factors for seizures
- are taking any serotonergic agents, including antidepressants such as selective serotonin reuptake inhibitors (SSRIs) and opioids
- have Hyponatremia and/or Syndrome of Inappropriate Antidiuretic Hormone Secretion (SIADH)
- are at high risk for life threatening systemic infections, such as those with infections related to central venous catheters in intensive care units

-Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

BACTOLID should not be taken with the following medicines

- Monoamine oxidase inhibitors (e.g. *phenelzine*, *isocarboxazid*) in the last two weeks
- Sympathomimetic agents (e.g. *pseudoephedrine*, *phenylpropanolamine*)
- Vasopressive agents (e.g. *epinephrine*, *norepinephrine*)
- Dopaminergic agents (e.g. *dopamine*, *dobutamine*)
- *Rifampin*
- Serotonin re-uptake inhibitors
- Tricyclic antidepressants
- Serotonin 5-HT1 receptor agonists
- *Meperidine* or *bupirone*

How to use BACTOLID

-How much to use

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help. Recommended dosage for adults and adolescents (12 years and above):

BACTOLID (LINEZOLID FILM-COATED TABLETS 600MG)

Linezolid 600 mg

Infection	Dosage	Recommended duration of treatment (consecutive days)
Complicated skin and skin structure infections	600 mg every 12 hours	10-14
Community-acquired pneumonia, including concurrent bacteremia		
Nosocomial pneumonia		
Vancomycin-resistant <i>Enterococcus faecium</i> infections, including concurrent bacteremia	600 mg every 12 hours	14-28
Uncomplicated skin and skin structure infections	Adults: 400 mg every 12 hours Adolescents: 600 mg every 12 hours	10-14

Recommended dosage for pediatric patients (birth through 11 years):

Infection	Dosage	Recommended duration of treatment

Infection	Dosage	(consecutive days)
Complicated skin and skin structure infections	10 mg/kg every 8 hours	10-14
Community-acquired pneumonia, including concurrent bacteremia		
Nosocomial pneumonia		
Vancomycin-resistant <i>Enterococcus faecium</i> infections, including concurrent bacteremia	10 mg/kg every 8 hours	14-28
Uncomplicated skin and skin structure infections	Below 5 years: 10 mg/kg every 8 hours 5-11 years: 10 mg/kg every 12 hours	10-14

-When to use it

Use as directed by your doctor or pharmacist.

-How long to use it

Continue taking for as long as your doctor recommends.

-If you forget to use it.

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

-Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking BACTOLID. Tell your doctor immediately if you become pregnant while taking this medication.

-Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give BACTOLID to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

Driving and using machines

This medicine may affect your ability to drive or use machines. If it makes

BACTOLID (LINEZOLID FILM-COATED TABLETS 600MG)

Linezolid 600 mg

you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Side effects

Like all medicines, BACTOLID can cause side effects, although not everybody gets them.

You may develop diarrhoea while taking or after taking BACTOLID. If this becomes severe or persistent or you notice that your stool contains blood or mucus, you should stop taking BACTOLID immediately and consult your doctor.

If you take BACTOLID for longer periods you may get problems with your vision such as blurred vision, changes in colour vision, difficulty in seeing detail or if your field of vision becomes restricted.

Other side effects some patients have experienced while taking BACTOLID are:

- Abdominal pain/cramps/distension
- Nausea
- Vomiting
- Moniliasis (yeast infection)
- Abnormal blood tests
- Abnormal liver function tests
- Headache
- Taste alteration
- Reversible anemia
- Sideroblastic anaemia (a form of anemia in which the bone marrow produces ringed sideroblasts)
- Leucopenia (decrease number of white blood cells)
- Thrombocytopenia (decrease number of platelets)
- Pancytopenia (decrease number of red blood cells, white blood cells and platelets)

- Optic neuropathy (loss of vision due to damage to the optic nerve)
- Anaphylaxis (serious allergic reaction)
- Lactic acidosis
- Peripheral neuropathy
- Convulsions
- Rash
- Angioedema (swelling of the skin and mucous membranes)
- Tongue discoloration
- Bullous skin disorders (e.g. Stevens Johnson syndrome, toxic epidermal necrolysis)
- Superficial tooth discoloration.
- Rhabdomyolysis (muscle pain, tenderness or weakness and dark urine)

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of BACTOLID

-Storage

- Keep out of the reach and sight of children.
- Store below 30°C. Protect from light and moisture.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

-What it looks like

White to off white, oval shaped, bevel edged, biconvex film coated tablets debossed with 'H' on one side and 'L8' on the other side.

Pack size: 3 x 10's Alu-Alu Blister pack

- Ingredients:

-Active ingredients: Linezolid

-Inactive ingredients:

Tablet core: Lactose monohydrate, Maize starch, Hydroxypropyl cellulose, Sodium starch glycolate, Magnesium stearate, Purified Water

Tablet film-coated: Opadry white (03B58895) Hypromellose (E464), Titanium dioxide (E171), Macrogol (E1521), Carnauba wax.

Sodium content

This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially 'sodium-free'.

-MAL number:

MALXXXXXXXXXX

Manufacturer

Hetero Labs Limited
Unit V Block-V and Block VA,
TSIC Formulation SEZ S. Nos. 439
440 441 and 458,
Polepally Village, Jadcherla Mandal,
Mahabubnagar, Telangana 509301
India.

BACTOLID (LINEZOLID FILM-COATED TABLETS 600MG)

Linezolid 600 mg

Product Registration Holder

Camber Laboratories Sdn. Bhd
Unit E13A-02, Menara SUEZCAP 2,
No. 2, KL Gateway, Jalan Kerinchi,
Gerbang Kerinchi Lestari, 59200 Kuala
Lumpur, Malaysia.

Date of revision

30 March 2026

Serial Number:

NPRA (R2) 24/148