

AKLIEF® CREAM

Trifarotene 50mcg/g

What is in this leaflet

1. What Aklief Cream is used for
2. How Aklief Cream works
3. Before you use Aklief Cream
4. How to use Aklief Cream
5. While you are using it
6. Side effects
7. Storage and Disposal of Aklief Cream
8. Product Description
9. Manufacturer
10. Product Registration Holder
11. Date of revision
12. Serial Number

1. What Aklief Cream is used for
Aklief is used for the cutaneous treatment of *Acne Vulgaris* of the face and/or the trunk in patients from 12 years of age and older, when many comedones (whiteheads and blackheads), papules and pustules (inflammatory pimples) are present.

2. How Aklief Cream works

Aklief contains the active substance trifarotene that belongs to a group of medicines called retinoids. Trifarotene modulates retinoid target genes in skin.

3. Before you use Aklief Cream

- When you must not use it

Do not use Aklief:

- If you are a woman planning pregnancy or if you are pregnant (see section "Pregnancy and breast-feeding")
- If you are allergic to trifarotene or any of the other ingredients of this medicine (listed in section 8)

Use in children

Aklief should not be used by children below 12 years of age.

- Before you start to use it

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Important: Aklief is intended for patients from 12 years of age and older only for use on the skin of the face and/or the trunk. Do not use this medicine on any other parts of your body. Do not swallow.

Keep Aklief away from children.

- Taking other medicines

Tell your doctor if you are using, have recently used or might use any other medicines.

Caution should be exercised if cosmetics or acne medications with peeling, irritant or drying effects are used, as they may produce additive irritant effects with the medicinal product. If your skin becomes irritated, contact your doctor.

4. How to use Aklief Cream

- How much to use

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

- When to use it

- Before using the pump for the first time, prime it by pressing down several times until a small amount of medicine is dispensed (up to 10 times maximum). The pump is now ready to use. Apply a thin layer of Aklief cream to the affected areas of the face (forehead, nose, chin and right and left cheeks) and all affected areas of the trunk **once a day, in the evening**, on a clean and dry skin:

- One (1) pump actuation should be enough to cover

the face (i.e. forehead, cheeks, nose and chin).

- Two (2) pump actuations should be enough to cover the upper trunk (i.e. reachable upper back, shoulders and chest). One (1) additional pump actuation may be used for middle and lower back if acne is present.
- More than four (4) pump actuations in one day is not recommended.
- Avoid contact with the eyes, eyelids, lips and mucous membranes such as inside the nose or the mouth. If you accidentally get cream in any of these areas wash it immediately with plenty of lukewarm water.
- Wash your hands immediately after applying the cream.

You are recommended to use a moisturizer as frequently as needed from the initiation of the Aklief treatment. The moisturizer can either be applied before or after Aklief, allowing sufficient time to let the skin to dry between the moisturizer and Aklief application.

If you stop using Aklief, the spots (whiteheads, blackheads and inflammatory pimples) will be reduced only after several application of this medicine. It is important that you continue using Aklief as long as prescribed by your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

- How long to use it

Your doctor will tell you how long you will need to use Aklief. After three months of treatment your doctor may need to assess the

AKLIEF[®] CREAM

Trifarotene 50mcg/g

continued improvement of your acne.

- If you forget to use it

If you forget to use Akliief in the evening, use it the next evening. Do not apply a double dose to make up for a forgotten dose.

- If you use too much (overdose)

If you use more Akliief than you should on your skin, you will not get rid of your acne any quicker, but your skin may become irritated, scaly and red. Talk to the doctor if you have used more Akliief than you should.

Contact a doctor or the national poison centre immediately if:

- a child has accidentally used this medicine
- you or someone else accidentally swallow this medicine.

Your doctor will advise you on what action needs to be taken.

5. While you are using it

- Things you must do

Redness, peeling, dryness, and stinging/burning may be experienced with the use of Akliief cream (see section 6 “Side effects”). Talk to a doctor if you experience these symptoms. You are recommended to apply a moisturizer from the initiation of treatment, which may help prevent such reactions. If symptoms do occur the doctor may instruct you to start using a moisturizer (if you have not already), to use the cream less often or to stop for a short time. If the symptoms persist, despite these measures, you may be asked to stop the cream altogether.

- Things you must not do

- Akliief should not be used on cuts, scrapes, abraded or eczematous skin.

- Akliief should not come into contact with the eyes, eyelids, lips, or mucous membranes. If the product accidentally enters the eye, wash immediately and abundantly with luke warm water. Be careful when applying to sensitive areas of the skin such as the neck or armpits.
- Caution should be exercised when Akliief cream is applied at the same time as other preparation used on the skin including cosmetics (see also section “Taking other medicines”).
- You should not use “waxing” as a depilatory method on skin treated with Akliief.
- If a reaction suggesting sensitivity to any component of the formula occurs, the use of Akliief should be discontinued.
- Akliief should not be used on sunburned skin. Excessive exposure to sunlight, including sunlamps or phototherapy should be avoided during the treatment. Use of sunscreen with Sun Protection Factor (SPF) of at least 30 and protective clothing (such as a hat and a shirt) over treated areas is recommended when exposure cannot be avoided. If nevertheless your face, chest, shoulders or back become sunburned, stop medication on the affected area until your skin is healed.

- Things to be careful of

Pregnancy and breast-feeding

Pregnancy

DO NOT use Akliief if you are pregnant or thinking of becoming pregnant. Your doctor can give you more information.

If you discover you are pregnant during treatment, stop application of this medicine and consult a doctor immediately.

Breast-feeding

When using Akliief there is a risk that the active substance in cream passes into your breast milk and a risk to the newborn/infant cannot be excluded. You and your doctor must make a decision whether to discontinue breast-feeding or to abstain from Akliief therapy, taking into account the benefit of breast-feeding for the child and the benefit of therapy for the mother.

To avoid the risk of ingestion by, and/or contact exposure of, an infant, nursing women should not apply Akliief to the chest or breast area.

Driving and using machines

Akliief has no or negligible influence on the ability to drive and use machines.

Akliief contains propylene glycol (E1520) which may cause skin irritation.

This medicine also contains 50 mg alcohol (ethanol) in each gram which is equivalent to 5% w/w. It may cause burning sensation on damaged skin.

6. Side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Application site reactions such as redness, peeling, dryness, and stinging/burning of the skin may often be experienced with the use of Akliief cream. See section 5 “While you are using it” for information on what to do if you experience such symptoms.

AKLIEF® CREAM

Trifarotene 50mcg/g

Aklief may cause the following side effects:

Common side effect (may affect up to 1 in 10 people):

- Application site irritation, itch, sunburn.

Uncommon side effect (may affect up to 1 in 100 people):

- Pain of the skin
- Dry skin
- Discolouration (loss of skin pigmentation)
- Erosion (skin loss)
- Rash
- Swelling
- Skin irritation
- Acne
- Dermatitis allergic (skin allergy)
- Erythema (redness)

Rare side effect (may affect up to 1 in 1000 people):

- Urticaria (hives)
- Vesicles
- Eczema “asteatotic” (dry skin with scales and fissures)
- Seborrheic dermatitis (red, scaly and itchy skin)
- Skin burning sensation
- Skin fissures
- Skin hyperpigmentation (darkening of skin pigmentation)
- Eyelid exfoliation (peeling of the eyelid skin) or oedema (swelling of the eyelid skin)
- Chapped lips
- Flushing (red face)

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

By reporting side effects, you can help provide more information on the safety of this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

7. Storage and Disposal of Aklief Cream

- Storage

Do not freeze. Do not store above 30°C.

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and tube/pump after EXP. The expiry date refers to the last day of that month.

Discard the tube or pump 6 months after first opening.

- Disposal

Do not throw away unused Aklief cream via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

8. Product Description

- What it looks like

Aklief is a white and homogenous cream.

Aklief is available in tube containing 5 grams of cream or pump of 15, 30 or 75 grams of cream.

Pack sizes of 1 tube or 1 pump.

Not all pack sizes may be marketed.

- Ingredients

- Active ingredient(s)

The active substance is trifarotene, one gram of cream contains 50 micrograms of trifarotene.

- Inactive ingredients

The other excipients are allantoin, Simulgel 600 PHA (copolymer of acrylamide and sodium acryloyldimethyltaurate, isohexadecane, polysorbate 80, sorbitan oleate), cyclomethicone, ethanol, phenoxyethanol, propylene glycol (E1520), triglycerides medium-chain and purified water.

- MAL number

MAL24076001ARZ

9. Manufacturer

Laboratoires Galderma
Z.I. Montdesir
74540 Alby-sur-Chéran
France

10. Product Registration Holder

Galderma Malaysia Sdn. Bhd.
Unit 904, Level 9, Uptown One
Jalan SS 21/58, Damansara Utama
47400 Petaling Jaya,
Selangor Darul Ehsan, Malaysia

11. Date of revision

December 2022

12. Serial Number

NPRA (R1/1)13082024/156