

FRONT PAGE

148 mm

Advocate™

Spot-on Solution for Cats

Product Description

Advocate Spot-on Solution for Small Cats and Ferrets is a clear yellow to brownish yellow colour solution. Each 0.4 ml pipette contains 40 mg imidacloprid and 4 mg moxidectin.

Advocate Spot-on Solution for Large Cats is a clear yellow to brownish yellow colour solution. Each 0.8 ml pipette contains 80 mg imidacloprid and 8 mg moxidectin.

Pharmacodynamics

Imidacloprid, 1-(6-Chloro-3-pyridylmethyl)-N-nitro-imidazolidin-2-ylideneamine is an ectoparasiticide belonging to the chloronicotinyl group of compounds. Chemically, it is more accurately described as a chloronicotinyl nitroguanidine. Imidacloprid is effective against larval flea stages and adult fleas. Flea larvae in the pet's surroundings are killed after contact with a pet treated with the veterinary medicinal product. Imidacloprid has a high affinity for the nicotinic acetylcholine receptors in the post-synaptic region of the central nervous system (CNS) of the flea. The ensuing inhibition of cholinergic transmission in insects results in paralysis and death. Due to the weak nature of the interaction with mammalian nicotinic receptors and the postulated poor penetration through the blood-brain barrier in mammals, it has virtually no effect on the mammalian CNS. Imidacloprid has minimal pharmacological activity in mammals.

Moxidectin, 23-(O-methylxime)-F28249 alpha is a second-generation macrocyclic lactone of the milbemycin family. It is a parasiticide which is active against many internal and external parasites. Moxidectin is active against larval stages (L3, L4) of *Dirofilaria immitis*. It is also active against gastrointestinal nematodes. Moxidectin interacts with GABA and glutamate-gated chloride channels. This leads to opening of the chloride channels on the postsynaptic junction, the inflow of chloride ions and induction of an irreversible resting state. The result is flaccid paralysis of affected parasites, followed by their death and/or expulsion. The veterinary medicinal product has a persistent action and protects cats for 4 weeks after a single application against reinfection with *Dirofilaria immitis*.

Pharmacokinetics

After topical administration of the veterinary medicinal product, imidacloprid is rapidly distributed over the animal's skin within one day of application. It can be found on the body surface throughout the treatment interval. Moxidectin is absorbed through the skin, reaching maximum plasma concentrations approximately 1 to 2 days after treatment in cats. Following absorption from the skin, moxidectin is distributed systemically throughout the body tissues but due to its lipophilicity it is concentrated mainly in the fat. It is slowly eliminated from the plasma as manifested by detectable moxidectin concentrations in plasma throughout the treatment interval of one month. The mean $t_{1/2}$ in cats ranges between 18.7 and 25.7 days. Studies evaluating the pharmacokinetic behaviour of moxidectin after multiple applications have indicated that steady state serum levels are achieved following approximately 4 consecutive monthly treatments in cats.

Target Species: Cats and ferrets (for Advocate Spot-on Solution for Small Cats and Ferrets only).

Indication

For cats suffering from, or at risk from, mixed parasitic infections. The veterinary medicinal product is only indicated when use against fleas and one or more of the other target parasites is indicated at the same time:

- the treatment and prevention of flea infestation (*Ctenocephalides felis*),
 - the treatment of ear mite infestation (*Otodectes cynotis*),
 - the treatment of notoedric mange (*Notoedres cati*),
 - the treatment of the lungworm *Eucoleus aerophilus* (syn. *Capillaria aerophila*) (adults),
 - the prevention of lungworm disease (L3/L4 larvae of *Aelurostrongylus abstrusus*),
 - the treatment of the lungworm *Aelurostrongylus abstrusus* (adults),
 - the treatment of the eye worm *Thelazia callipaeda* (adults),
 - the prevention of heartworm disease (L3 and L4 larvae of *Dirofilaria immitis*),
 - the treatment of infections with gastrointestinal nematodes (L4 larvae, immature adults and adults of *Toxocara cati* and *Ancylostoma tubaeforme*).
- The veterinary medicinal product can be used as part of a treatment strategy for flea allergy dermatitis (FAD).

For ferrets suffering from, or at risk from, mixed parasitic infections. The veterinary medicinal product is only indicated when use against fleas and the prevention of heartworm disease is indicated at the same time:

- the treatment and prevention of flea infestation (*Ctenocephalides felis*),
- the prevention of heartworm disease (L3 and L4 larvae of *Dirofilaria immitis*).

Recommended Dosage

To ensure a correct dosage, body weight should be determined as accurately as possible.

Dosage schedule for cats:

The recommended minimum doses are 10 mg/kg bodyweight imidacloprid and 1.0 mg/kg bodyweight moxidectin, equivalent to 0.1 ml/kg bodyweight of the veterinary medicinal product.

For treatment or prevention of infestations with the parasites indicated for use of this veterinary medicinal product, the need for and frequency of re-treatment(s) should be based on professional advice and should take into account the local epidemiological situation and the animal's lifestyle.

Weight of cat (kg)	Pipette size to be used	Volume (ml)	Imidacloprid (mg/kg bw)	Moxidectin (mg/kg bw)
≤ 4 kg	Advocate Spot-on Solution for Small Cats and Ferrets	0.4	minimum of 10	minimum of 1
> 4-8 kg	Advocate Spot-on Solution for Large Cats	0.8	10 – 20	1 – 2
> 8 kg	the appropriate combination of pipettes			

Flea treatment and prevention (*Ctenocephalides felis*)

One treatment prevents future flea infestation for 4 weeks. Existing pupae in the environment may emerge for 6 weeks or longer after treatment is initiated, depending upon climatic conditions. Therefore, it may be necessary to combine veterinary medicinal product treatment with environmental treatments aimed at breaking the flea life cycle in the surroundings. This can result in a more rapid reduction in the household flea population. The

PA550274X

veterinary medicinal product should be administered at monthly intervals when used as part of a treatment strategy for flea allergy dermatitis.

Treatment of ear mite infestation (*Otodectes cynotis*)

A single dose of the veterinary medicinal product should be administered. A further veterinary examination 30 days after treatment is recommended as some animals may require a second treatment. Do not apply directly to the ear canal.

Treatment of notoedric mange (*Notoedres cati*)

A single dose of the veterinary medicinal product should be administered.

Treatment of the lungworm *Eucoleus aerophilus* (syn. *Capillaria aerophila*) (adults)

A single dose of the veterinary medicinal product should be administered.

Prevention of *Aelurostrongylus abstrusus*

The veterinary medicinal product should be administered monthly.

Treatment of *Aelurostrongylus abstrusus*

The veterinary medicinal product should be administered monthly for three consecutive months.

Treatment of the eye worm *Thelazia callipaeda* (adults)

A single dose of the veterinary medicinal product should be administered.

Heartworm prevention (*Dirofilaria immitis*)

Cats in areas endemic for heartworm, or those which have travelled to endemic areas, may be infected with adult heartworms. Therefore prior to treatment with the veterinary medicinal product, the advice provided in section Warnings and Precautions should be considered.

For prevention of heartworm disease, the veterinary medicinal product must be applied at regular monthly intervals during the time of the year when mosquitoes (the intermediate hosts which carry and transmit heartworm larvae) are present. The veterinary medicinal product may be administered throughout the year. The first dose may be given after first possible exposure to mosquitoes, but not more than one month after this exposure. Treatment should continue at regular monthly intervals until 1 month after the last exposure to mosquitoes. To establish a treatment routine, it is recommended that the same day or date be used each month. When replacing another heartworm preventative product in a heartworm prevention programme, the first treatment with the veterinary medicinal product must be given within 1 month of the last dose of the former veterinary medicinal product. In non-endemic areas there should be no risk of cats having heartworm. Therefore, they can be treated without special precautions.

Roundworm and hookworm treatment (*Toxocara cati* and *Ancylostoma tubaeforme*)

In areas endemic for heartworm, monthly treatment may significantly reduce the risk of re-infection caused by the respective roundworms and hookworms. In areas non-endemic for heartworm, the veterinary medicinal product can be used as part of a seasonal prevention programme against fleas and gastrointestinal nematodes.

Dosage schedule for ferrets:

One pipette of the veterinary medicinal product (0.4 ml) should be administered per animal. Do not exceed the recommended dose. For treatment or prevention of infestations with the parasites indicated for use of this veterinary medicinal product, the need for and frequency of re-treatment(s) should be based on professional advice and should take into account the local epidemiological situation and the animal's lifestyle.

Flea treatment and prevention (*Ctenocephalides felis*)

One treatment prevents future flea infestation for 3 weeks. Under heavy flea pressure it may be necessary to repeat administration after 2 weeks.

Heartworm prevention (*Dirofilaria immitis*)


Ferrets in areas endemic for heartworm, or those which have travelled to endemic areas, may be infected with adult heartworms. Therefore prior to treatment with the veterinary medicinal product, the advice provided in section Warnings and Precautions should be considered.

For prevention of heartworm disease, the veterinary medicinal product must be applied at regular monthly intervals during the time of the year when mosquitoes (the intermediate hosts which carry and transmit heartworm larvae) are present. The veterinary medicinal product may be administered throughout the year. The first dose may be given after first possible exposure to mosquitoes, but not more than one month after this exposure. Treatment should continue at regular monthly intervals until 1 month after the last exposure to mosquitoes. In non-endemic areas there should be no risk of ferrets having heartworm. Therefore, they can be treated without special precautions.

Method of Administration

External use.

Remove one pipette from the package. Then hold the pipette in an upright position, and twist and pull off the cap. Reverse the cap and use it to twist and remove the seal from the pipette, as shown.



Part the fur on the animal's neck at the base of the skull until the skin is visible. Place the tip of the pipette on the skin and squeeze the pipette firmly several times to empty its contents directly onto the skin. Application at the base of the skull will minimise the opportunity for the animal to lick the veterinary medicinal product. Apply only to undamaged skin.



Contraindications

Do not use in kittens under 9 weeks of age. Do not use in cases of hypersensitivity to the active substances or to any of the excipients. For ferrets: Do not use Advocate Spot-on Solution for Large Cats (0.8 ml) or Advocate Spot-on Solution for Dogs (any size). Do not use in dogs. Instead, the corresponding "Advocate Spot-on Solution for Dogs" product, which contains 100 mg/ml imidacloprid and 25 mg/ml moxidectin, must be used for dogs. Do not use on canaries.

Warnings and Precautions

The veterinary medicinal product's efficacy has not been tested in ferrets weighing over 2 kg and therefore the duration of effect might be shorter in these animals. Brief contact of the animal with water on one or two occasions between

• Item code placement is flexible and can be moved to suit artwork

PRODUCT INFO

BLUE #:	550274M	Product Name:	Advocate SO Solution for Small Cats and Ferrets/Large Cat
Item Code:	PA550274X	Component:	Package Inserts and Leaflets
Product Code:	N/A	Pack Size:	3 x 0.4, 0.8 ml
Previous Item Code	PA502162X		

ARTWORK INFO

Template:	PI_148x297mm_Kiel_v2a	Packaging Spec(s):	TPM_016191
Barcodes/Type:	Pharmacode: 249	Add. Info:	N/A
		Minimum Core Data Point Size:	7pt
GTIN:	GTIN Not Required	Proof #	P2a
		By/Date	TCK 27-NOV-2025



ELANCO ARTWORK LEGEND v16

75 mm

Pharmacode
Read Direction

297 mm

148 mm

monthly treatments is unlikely to significantly reduce the efficacy of the veterinary medicinal product. However, frequent shampooing or immersion of the animal in water after treatment may reduce the efficacy of the veterinary medicinal product.

The possibility that other animals in the same household can be a source of re-infection with fleas, mites, gastrointestinal nematodes, heartworm and/or lungworm should be considered, and these should be treated as necessary with an appropriate product.

Unnecessary use of antiparasitics or use deviating from the instructions given in the SPC may increase the resistance selection pressure and lead to reduced efficacy. The decision to use the product should be based on confirmation of the parasitic species and burden, or of the risk of infestation based on its epidemiological features, for each individual animal.

Special precautions for safe use in the target species:

The treatment of cats weighing less than 1 kg and ferrets weighing less than 0.8 kg should be based on a benefit-risk assessment.

There is limited experience on the use of the veterinary medicinal product in sick and debilitated animals, thus the veterinary medicinal product should only be used based on a benefit-risk assessment for these animals.

Do not apply in the mouth, in the eyes or the ears of the animal.

Care should be taken that the veterinary medicinal product is not ingested by animals and does not come into contact with the eyes or mouth of the recipient and/or other animals.

Consider carefully the correct application method described in section Method of Administration, especially that the veterinary medicinal product should be applied to the site specified in order to minimise the risk for the animal to lick the veterinary medicinal product.

Do not allow recently treated animals to groom each other. Do not allow treated animals to come into contact with untreated animals until the application site is dry.

It is recommended that cats and ferrets living in, or travelling to areas endemic for heartworm are treated monthly with the veterinary medicinal product to protect them from heartworm disease.

Whilst the accuracy of diagnosis of heartworm infection is limited, it is recommended that attempts be made to check the heartworm status of any cat and ferret aged over 6 months, before beginning prophylactic treatment, as use of the veterinary medicinal product on cats or ferrets which have adult heartworms may cause serious adverse effects, including death. If adult heartworm infection is diagnosed, the infection should be treated in accordance with current scientific knowledge.

In certain individual cats *Notodres cati* infestation may be severe. In these severe cases concomitant supportive treatment is necessary as treatment with the veterinary medicinal product alone may not be sufficient to prevent death of the animal.

The safety of the veterinary medicinal product has not been established in cats with severe clinical signs of *T. brevior*. Use of the veterinary medicinal product in such cases should be based on the benefit-risk assessment of the veterinarian.

Imidacloprid is toxic to birds, especially canaries.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This veterinary medicinal product can cause skin, eye, or mouth irritation.

In very rare cases the veterinary medicinal product may cause skin sensitisation or transient skin reactions (for example numbness, irritation or burning/tingling sensation).

In very rare cases the veterinary medicinal product may cause respiratory irritation in sensitive individuals.

People with known hypersensitivity to benzyl alcohol, imidacloprid or moxidectin should administer the veterinary medicinal product with caution. Avoid contact with skin, eyes or mouth.

Do not eat, drink or smoke during application.

Wash hands thoroughly after use.

After application do not stroke or groom animals until the application site is dry.

In case of accidental spillage onto skin, wash off immediately with soap and water.

If the veterinary medicinal product accidentally gets into eyes, they should be thoroughly flushed with water.

If skin or eye symptoms persist, or the veterinary medicinal product is accidentally swallowed, seek medical advice immediately and show the package leaflet or the label to the physician.

Special precautions for the protection of the environment:

The veterinary medicinal product should not enter water courses as imidacloprid and moxidectin may be dangerous for fish and other aquatic organisms.

Other precautions:

The solvent in the veterinary medicinal product may stain or damage certain materials including leather, fabrics, plastics and finished surfaces. Allow the application site to dry before permitting contact with such materials.

Interaction with Other Medicinal Products and Other Forms of Interaction

During treatment with the veterinary medicinal product no other antiparasitic macrocyclic lactone should be administered.

No interactions between the veterinary medicinal product and routinely used veterinary medicinal products or medical or surgical procedures have been observed.

Use During Pregnancy, Lactation or Lay

The safety of the veterinary medicinal product has not been established during pregnancy and lactation in the target species.

Pregnancy and lactation:

The use is not recommended during pregnancy and lactation.

Fertility:

Do not use in breeding animals.

Adverse Events

Cats and ferrets:

Rare (1 to 10 animals / 10,000 animals treated):	Application site greasy fur ¹ Vomiting ¹ Hypersensitivity reaction (local) Erythema ¹
Very rare (< 1 animal / 10,000 animals treated, including isolated reports):	Behavioural disorder (e.g. agitation) ² Hypersalivation ^{3,4} Neurological signs ³ Pruritus ⁵ Inappetence ² , Lethargy ²

¹ These signs disappear without further treatment.

² Transiently noted and related to sensation at application site.

³ If the animal licks the application site, in most cases transient.

⁴ This is not a sign of intoxication and disappears within minutes without treatment. Correct application will minimise licking of the application site.

⁵ In cats, transient.

Overdose

Up to 10 times the recommended dose was tolerated in cats with no evidence of adverse effects or undesirable clinical signs.

The veterinary medicinal product was administered to kittens at up to 5 times the recommended dose, every 2 weeks for 6 treatments, and there were no serious safety concerns. Transient mydriasis, salivation, vomiting and transient rapid respiration were observed.

After accidental oral ingestion or overdose, neurological signs (most of which are transient) such as ataxia, generalised tremors, ocular signs (dilated pupils, little pupillary reflex, nystagmus), abnormal respiration, salivation and vomiting may occur in very rare cases.

The veterinary medicinal product was administered to ferrets at 5 times the recommended dose, every 2 weeks for 4 treatments, and there was no evidence of adverse effects or undesirable clinical signs.

In case of accidental oral uptake, symptomatic treatment should be administered. There is no known specific antidote. The use of activated charcoal may be beneficial.

Withdrawal Period

Not applicable.

Storage Condition

Do not store above 30°C.

Dosage forms and packaging available

Advocate Spot-on Solution for Cats is available in triple pipettes per box for the following:

Advocate Spot-on Solution for Small Cats and Ferrets

Advocate Spot-on Solution for Large Cats

Marketing Authorisation Holder:

Elanco Malaysia Sdn Bhd
Unit 5.04, Level 5 & 6, Tower Block, The Bousteador,
No. 10, Jalan PJU 7/6, Mutiara Damansara,
47800 Petaling Jaya, Selangor, Malaysia

Manufactured by:

KVP Pharma + Veterinär Produkte GmbH
Projensdorfer Straße 324, D-24106 Kiel, Germany.

Date of revision: Oct 2025

Advocate, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2025 Elanco or its affiliates



Pharmacode
Read Direction

75 mm

297 mm

Elanco