

VIAGRA[®] 50mg Orodispersible Films

Sildenafil Citrate (50mg)

PLD Title : Sildenafil Citrate (VIAGRA 50mg Orodispersible Films)
PLD Date : 10 March 2026
Country : Malaysia
Reference Document : EU SmPC, Commission Decision on 9 Feb 2024 (EMA/H/C/000202/X/0115)
Reason for change : TW3429559 - Update of new registration number.

VIAGRA[®] 50mg Orodispersible Films

Sildenafil Citrate (50mg)

What is in this leaflet

1. What VIAGRA is used for
2. How VIAGRA works
3. Before you use VIAGRA
4. How to use VIAGRA
5. While you are using it
6. Side effects
7. Storage and Disposal of VIAGRA
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision

What VIAGRA is used for

VIAGRA is a treatment for adult men with erectile dysfunction, sometimes known as impotence. This is when a man cannot get, or keep a hard, erect penis suitable for sexual activity.

How VIAGRA works

VIAGRA contains the active substance sildenafil which belongs to a group of medicines called phosphodiesterase type 5 (PDE5) inhibitors. It works by helping to relax the blood vessels in your penis, allowing blood to flow into your penis when you get sexually excited. VIAGRA will only help you to get an erection if you are sexually stimulated.

Before you use VIAGRA

When you must not use it

- If you are allergic to sildenafil or any of the other ingredients of this medicine.
- If you are taking medicines called nitrates, as the combination may lead to a dangerous fall in your blood pressure. Tell your doctor if you are taking any of these medicines which are often

given for relief of angina pectoris (or "chest pain"). If you are not certain, ask your doctor or pharmacist.

- If you are using any of the medicines known as nitric oxide donors such as amyl nitrite ("poppers"), as the combination may also lead to a dangerous fall in your blood pressure.

- If you are taking riociguat. This medicine is used to treat pulmonary arterial hypertension (i.e., high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (i.e., high blood pressure in the lungs secondary to blood clots). PDE5 inhibitors, such as VIAGRA have been shown to increase the hypotensive effects of this medicine. If you are taking riociguat or are unsure tell your doctor.

- If you have a severe heart or liver problem.

- If you have recently had a stroke or a heart attack, or if you have low blood pressure.

- If you have certain rare inherited eye diseases (such as retinitis pigmentosa).

- If you have ever had loss of vision due to non-arteritic anterior ischaemic optic neuropathy (NAION).

Before you start to use it

Talk to your doctor, pharmacist or nurse before taking VIAGRA

- If you have sickle cell anaemia (an abnormality of red blood cells), leukaemia (cancer of blood cells), multiple myeloma (cancer of bone marrow).

- If you have a deformity of your penis or Peyronie's disease.

- If you have problems with your heart. Your doctor should carefully check whether your heart can take the additional strain of having sex.

- If you currently have a stomach ulcer, or a bleeding problems (such as haemophilia).

- If you experience sudden decrease or loss of vision, stop taking VIAGRA and contact your doctor immediately.

You should not use VIAGRA with any other oral or local treatments for erectile dysfunction.

You should not use VIAGRA with treatments for pulmonary arterial hypertension (PAH) containing sildenafil or any other PDE5 inhibitors.

You should not take VIAGRA if you do not have erectile dysfunction.

You should not take VIAGRA if you are a woman.

Special considerations for patients with kidney or liver problems

You should tell your doctor if you have kidney or liver problems. Your doctor may decide on a lower dose for you.

Children and adolescents

VIAGRA should not be given to individuals under the age of 18.

Taking other medicines

Tell your doctor or pharmacist if you are taking, have recently

VIAGRA[®] 50mg Orodispersible Films

Sildenafil Citrate (50mg)

taken or might take any other medicines.

VIAGRA may interfere with some medicines, especially those used to treat chest pain. In the event of a medical emergency, you should tell your doctor, pharmacist or nurse that you have taken VIAGRA and when you did. Do not take VIAGRA with other medicines unless your doctor tells you that you can.

You should not take VIAGRA if you are taking medicines called nitrates, as the combination of these medicines may lead to a dangerous fall in your blood pressure. Always tell your doctor, pharmacist or nurse if you are taking any of these medicines that are often used for the relief of angina pectoris (or “chest pain”).

You should not take VIAGRA if you are using any of the medicines known as nitric oxide donors such as amyl nitrite (“poppers”) as the combination may also lead to a dangerous fall in your blood pressure.

Tell your doctor or pharmacist if you are already taking riociguat.

If you are taking medicines known as protease inhibitors, such as for the treatment of HIV, your doctor may start you on the lowest dose (25 mg film-coated tablets) of VIAGRA.

Some patients who take alpha-blocker therapy for the treatment of high blood pressure or prostate enlargement may experience dizziness or light-headedness, which may be caused by low blood pressure upon sitting or

standing up quickly. Certain patients have experienced these symptoms when taking VIAGRA with alpha-blockers. This is most likely to happen within 4 hours after taking VIAGRA. To reduce the chance that these symptoms might happen, you should be on a regular daily dose of your alpha-blocker before you start VIAGRA. Your doctor may start you on a lower dose (25 mg film-coated tablets) of VIAGRA.

Tell your doctor or pharmacist if you are taking medicines containing sacubitril/valsartan, used to treat heart failure.

Pregnancy, breast-feeding and fertility

VIAGRA is not indicated for use by women.

How to use VIAGRA

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much to use

The recommended starting dose is 50 mg.

You should not take VIAGRA more than once a day.

Do not take VIAGRA orodispersible films in combination with other sildenafil containing products including VIAGRA film-coated tablets or VIAGRA orodispersible tablets.

When to use it

You should take VIAGRA about one hour before you plan to have sex. The amount of time VIAGRA takes to work varies from person to person,

but it normally takes between half an hour and one hour.

Carefully peel open the aluminium pouch with dry hands. Do not cut it open. Take out the orodispersible film with a dry finger and immediately place the orodispersible film on the tongue, where it will disintegrate in seconds with or without water. During disintegration, saliva may be swallowed; however, the film should not be intentionally swallowed or kept in the mouth longer than necessary.

The orodispersible film should be taken on an empty stomach as you may find that it takes longer to start working if you take it with a heavy meal.

If you require a second 50 mg orodispersible film to make a 100 mg dose, you should wait until the first one has completely disintegrated before taking the second orodispersible film.

If you feel that the effect of VIAGRA is too strong or too weak, talk to your doctor or pharmacist.

VIAGRA will only help you to get an erection if you are sexually stimulated.

If VIAGRA does not help you to get an erection, or if your erection does not last long enough for you to complete sexual intercourse you should tell your doctor.

If you use too much (overdose)

You may experience an increase in side effects and their severity. Doses above 100 mg do not increase the efficacy.

VIAGRA[®] 50mg Orodispersible Films

Sildenafil Citrate (50mg)

You should not take more films than your doctor tells you to.

Contact your doctor if you take more films than you should.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

While you are using VIAGRA

Things you must do

Use your medicine exactly as your doctor has told you.

Things you must not do

Drinking alcohol can temporarily impair your ability to get an erection. To get the maximum benefit from your medicine, you are advised not to drink excessive amounts of alcohol before taking VIAGRA.

Things to be careful of

VIAGRA can cause dizziness and can affect vision. You should be aware of how you react to VIAGRA before you drive or use machinery.

Side effects

Like all medicines, this medicine can cause side effects although not everybody gets them. The side effects reported in association with the use of VIAGRA are usually mild to moderate and of a short duration.

If you experience any of the following serious side effects stop taking VIAGRA and seek medical help immediately:

- An allergic reaction – this occurs **uncommonly** (may affect up to 1 in 100 people)
Symptoms include sudden wheeziness, difficulty in breathing or dizziness, swelling of the eyelids, face, lips or throat.
- Chest pains - this occurs **uncommonly**
If this occurs during or after intercourse:
- Get in a semi-sitting position and try to relax.
- **Do not use nitrates** to treat your chest pain.
- Prolonged and sometimes painful erections – this occurs **rarely** (may affect up to 1 in 1 000 people)
If you have an erection which lasts for more than 4 hours, you should contact a doctor immediately.
- A sudden decrease or loss of vision - this occurs **rarely**
- Serious skin reactions - this occurs **rarely**
Symptoms may include severe peeling and swelling of the skin, blistering of the mouth, genitals and around the eyes, fever.
- Seizures or fits - this occurs **rarely**

Other side effects:

Very common (may affect more than 1 in 10 people): headache.

Common (may affect up to 1 in 10 people): nausea, facial flushing, hot flush (symptoms include a sudden feeling of heat in your upper body), indigestion, colour tinge to vision, blurred vision, visual

disturbance, stuffy nose and dizziness.

Uncommon (may affect up to 1 in 100 people): vomiting, skin rash, eye irritation, bloodshot eyes /red eyes, eye pain, seeing flashes of light, visual brightness, light sensitivity, watery eyes, pounding heartbeat, rapid heartbeat, high blood pressure, low blood pressure, muscle pain, feeling sleepy, reduced sense of touch, vertigo, ringing in the ears, dry mouth, blocked or stuffy sinuses, inflammation of the lining of the nose (symptoms include runny nose, sneezing and stuffy nose), upper abdominal pain, gastro-oesophageal reflux disease (symptoms include heartburn), presence of blood in urine, pain in the arms or legs, nosebleed, feeling hot and feeling tired.

Rare (may affect up to 1 in 1 000 people): fainting, stroke, heart attack, irregular heartbeat, temporary decreased blood flow to parts of the brain, feeling of tightening of the throat, numb mouth, bleeding at the back of the eye, double vision, reduced sharpness of vision, abnormal sensation in the eye, swelling of the eye or eyelid, small particles or spots in your vision, seeing halos around lights, dilation of the pupil of the eye, discolouration of the white of the eye, penile bleeding, presence of blood in semen, dry nose, swelling of the inside of the nose, feeling irritable and sudden decrease or loss of hearing.

From post-marketing experience cases of unstable angina (a heart condition) and sudden death have been reported rarely. Of note, most, but not all, of the men who experienced these side effects had heart problems before

VIAGRA[®] 50mg Orodispersible Films

Sildenafil Citrate (50mg)

taking this medicine. It is not possible to determine whether these events were directly related to VIAGRA.

Reporting of side effects

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine. This includes any possible side effects not listed in this leaflet. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF)].

Storage and Disposal of VIAGRA

Storage

Keep this medicine out of the sight and reach of children.

Store below 30°C. Do not use this medicine after the expiry date which is stated on the carton and pouch after EXP. This medicine does not require any special storage conditions.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

What it looks like

Thin pale red coloured orodispersible film (approximately 24 mm x 32 mm).

Each orodispersible film is packed in an individual foil pouch.

They are supplied in cartons containing 4 pouches.

What VIAGRA contains

- Ingredients

The active substance is sildenafil. Each orodispersible film contains 50 mg of sildenafil (as the citrate salt).

- Inactive ingredients

The other ingredients are hydroxypropylcellulose (E463), macrogol 400, crospovidone (E1202), povidone (E1201), sucralose (E955), Polyvinyl alcohol/polyethylene glycol/graft copolymer, l-menthol, hypromellose (E464), titanium dioxide (E171), ferric oxide (red) (E172), anhydrous Ethanol, purified water.

MAL Numbers

MAL26036002ACRZ

Manufacturer

Kyukyu Pharmaceutical Co., Ltd.
Toyama Plant
32-7 Hibari, Imizu City,
Toyama,
Japan

Product Registration Holder

Viatrix Sdn Bhd
15-03 & 15-04, Level 15,
Imazium,
No. 8, Jalan SS 21/37,
Damansara Uptown,
47400 Petaling Jaya,
Selangor, Malaysia.

Date of Revision

10/03/2026

Serial Number

NPRA (R1) 25/031

PLD-VIAGRAODF-0326