

Trajenta Duo®



1. NAME OF THE MEDICINAL PRODUCT

Trajenta Duo 2.5 mg/500 mg film-coated tablets
Trajenta Duo 2.5 mg/850 mg film-coated tablets
Trajenta Duo 2.5 mg/1000 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 2.5 mg of linagliptin and 500 mg, 850 mg or 1000 mg of metformin hydrochloride.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet (tablet).

Oval, biconvex, light yellow, film-coated tablet of 16.3 mm x 8.6 mm debossed with "D2/500" on one side and the company logo on the other.

Oval, biconvex, light orange, film-coated tablet of 19.2 mm x 9.4 mm debossed with "D2/850" on one side and the company logo on the other.

Oval, biconvex, light pink, film-coated tablet of 21.1 mm x 9.7 mm debossed with "D2/1000" on one side and the company logo on the other.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Trajenta Duo is indicated as an adjunct to diet and exercise to improve glycaemic control in adults with type 2 diabetes mellitus when treatment with both linagliptin and metformin is appropriate (see *Posology and Method of Administration* and *Clinical Studies*)

4.2 Posology and method of administration

Posology

Adults with normal renal function ($GFR \geq 90$ ml/min)

The dose of antihyperglycaemic therapy with Trajenta Duo should be individualised on the basis of the patient's current regimen, effectiveness, and tolerability, while not exceeding the maximum recommended daily dose of 5 mg linagliptin plus 2000 mg of metformin hydrochloride.

For patients currently not treated with metformin

In patient currently not treated with metformin, initiate treatment with 2.5mg linagliptin/ 500mg metformin hydrochloride twice daily.

For patients inadequately controlled on maximal tolerated dose of metformin monotherapy

For patients not adequately controlled on metformin alone, the usual starting dose of Trajenta Duo should provide linagliptin dosed as 2.5 mg twice daily (5 mg total daily dose) plus the dose of metformin already being taken.

For patients switching from co-administration of linagliptin and metformin

For patients switching from co-administration of linagliptin and metformin, Trajenta Duo should be initiated at the dose of linagliptin and metformin already being taken.

For patients inadequately controlled on dual combination therapy with the maximal tolerated dose of metformin and a sulphonylurea

The dose of Trajenta Duo should provide linagliptin dosed as 2.5 mg twice daily (5 mg total daily dose) and a dose of metformin similar to the dose already being taken. When linagliptin plus metformin hydrochloride is used in combination with a sulphonylurea, a lower dose of the sulphonylurea may be required to reduce the risk of hypoglycaemia (see section 4.4).

For patients inadequately controlled on dual combination therapy with insulin and the maximal tolerated dose of metformin

The dose of Trajenta Duo should provide linagliptin dosed as 2.5mg twice daily (5mg total daily dose) and a dose of metformin similar to the dose already being taken. When linagliptin plus metformin hydrochloride is used in combination with insulin, a lower dose of insulin may be required to reduce the risk of hypoglycaemia (see section 4.4).

For the different doses of metformin, Trajenta Duo is available in strengths of 2.5 mg linagliptin plus 500 mg metformin hydrochloride, 2.5 mg linagliptin plus 850 mg metformin hydrochloride and 2.5 mg linagliptin plus 1000 mg metformin hydrochloride.

Special populations

Elderly

As metformin is excreted by the kidney, Trajenta Duo should be used with caution as age increases. Monitoring of renal function is necessary to aid in prevention of metformin-associated lactic acidosis, particularly in the elderly (see sections 4.3 and 4.4).

Renal impairment

An eGFR should be assessed before initiation of treatment with metformin containing products and at least annually thereafter. In patients at an increased risk of further progression of renal impairment and in the elderly, renal function should be assessed more frequently, e.g. every 3-6 months. Factors that may increase the risk of lactic acidosis (see 4.4) should be reviewed before considering initiation of metformin in patients with eGFR<60 ml/min.

If no adequate strength of Trajenta Duo is available, individual monocomponents should be used instead of the fixed dose combination.

Table 1: Posology for renally impaired patients

GFR ml/min	Metformin	Linagliptin
60 - 89	Maximum daily dose is 3000 mg. Dose reduction may be considered in relation to declining renal function.	No dose adjustment is required.
45 - 59	Maximum daily dose is 2000 mg. The starting dose is at most half of the	No dose adjustment is required.

	maximum dose.	
30 - 44	Maximum daily dose is 1000 mg. The starting dose is at most half of the maximum dose.	No dose adjustment is required.
<30	Metformin is contraindicated.	No dose adjustment is required.

Hepatic impairment

Trajenta Duo is not recommended in patients with hepatic impairment due to the active substance metformin (see sections 4.3 and 5.2). Clinical experience with Trajenta Duo in patients with hepatic impairment is lacking.

Paediatric population

A clinical trial did not establish efficacy of linagliptin in paediatric patients 10 to 17 years of age (see section Clinical Studies). Therefore, treatment of children and adolescents with Trajenta Duo is not recommended. Linagliptin has not been studied in paediatric patients under 10 years of age.

Method of administration

Trajenta Duo should be taken twice daily with meals to reduce the gastrointestinal adverse reactions associated with metformin.

All patients should continue their diet with an adequate distribution of carbohydrate intake during the day. Overweight patients should continue their energy-restricted diet.

If a dose is missed, it should be taken as soon as the patient remembers. However, a double dose should not be taken at the same time. In that case, the missed dose should be skipped.

4.3 Contraindications

- Hypersensitivity to the active substances or to any of the excipients listed in section 6.1.
- Any type of acute metabolic acidosis (such as lactic acidosis, diabetic ketoacidosis)
- Diabetic pre-coma
- Severe renal failure ($\text{CrCl} < 30 \text{ ml/min}$ or $\text{eGFR} < 30 \text{ ml/min/1.73m}^2$).
- Acute conditions with the potential to alter renal function such as: dehydration, severe infection, shock, intravascular administration of iodinated contrast agents (see section 4.4).
- Disease which may cause tissue hypoxia (especially acute disease, or worsening of chronic disease) such as: decompensated heart failure, respiratory failure, recent myocardial infarction, shock.
- Hepatic impairment, acute alcohol intoxication, alcoholism (see section 4.5).

4.4 Special warnings and precautions for use

General

Trajenta Duo should not be used in patients with type 1 diabetes or for the treatment of diabetic ketoacidosis.

Hypoglycaemia

When linagliptin was added to a sulphonylurea on a background of metformin, the incidence of hypoglycaemia was increased over that of placebo.

Sulphonylureas and insulin are known to cause hypoglycaemia. Therefore, caution is advised when Trajenta Duo is used in combination with a sulphonylurea and/or insulin. A dose reduction of the sulphonylurea or insulin may be considered (see section 4.2).

Hypoglycaemia is not identified as adverse reaction for linagliptin, metformin, or linagliptin plus metformin. In clinical trials, the incidence rates of hypoglycaemia were comparably low in patients taking linagliptin in combination with metformin or metformin alone.

Lactic acidosis

Lactic acidosis, a very rare but serious metabolic complication, most often occurs at acute worsening of renal function or cardiorespiratory illness or sepsis. Metformin accumulation occurs at acute worsening of renal function and increases the risk of lactic acidosis.

In case of dehydration (severe diarrhoea or vomiting, fever or reduced fluid intake), Trajenta Duo should be temporarily discontinued and contact with a health care professional is recommended.

Medicinal products that can acutely impair renal function (such as antihypertensives, diuretics and NSAIDs) should be initiated with caution in metformin-treated patients. Other risk factors for lactic acidosis are excessive alcohol intake, hepatic impairment, inadequately controlled diabetes, ketosis, prolonged fasting and any conditions associated with hypoxia, as well as concomitant use of medicinal products that may cause lactic acidosis (see sections 4.3 and 4.5).

Patients and/or care-givers should be informed of the risk of lactic acidosis. Lactic acidosis is characterised by acidotic dyspnea, abdominal pain, muscle cramps, asthenia and hypothermia followed by coma. In case of suspected symptoms, the patient should stop taking Trajenta Duo and seek immediate medical attention. Diagnostic laboratory findings are decreased blood pH (<7.35), increased plasma lactate levels (>5 mmol/l), and an increased anion gap and lactate/pyruvate ratio.

Patients with known or suspected mitochondrial diseases

In patients with known mitochondrial diseases such as Mitochondrial Encephalopathy with Lactic Acidosis, and Stroke-like episodes (MELAS) syndrome and Maternal inherited diabetes and deafness (MIDD), metformin is not recommended due to the risk of lactic acidosis exacerbation and neurologic complications which may lead to worsening of the disease.

In case of signs and symptoms suggestive of MELAS syndrome or MIDD after the intake of metformin, treatment with Trajenta Duo should be withdrawn immediately and prompt diagnostic evaluation should be performed.

Administration of iodinated contrast agent

Intravascular administration of iodinated contrast agents may lead to contrast induced nephropathy, resulting in metformin accumulation and an increased risk of lactic acidosis. Trajenta Duo should be discontinued prior to or at the time of the imaging procedure and not restarted until at least 48 hours after, provided that renal function has been re-evaluated and found to be stable, see sections 4.2 and 4.5.

Renal function

GFR should be assessed before treatment initiation and regularly thereafter, see section 4.2. Trajenta Duo is contraindicated in patients with GFR<30 ml/min and should be temporarily discontinued in the presence of conditions that alter renal function (see section 4.3).

Cardiac function

Patients with heart failure are more at risk of hypoxia and renal impairment. In patients with stable chronic heart failure, Trajenta Duo may be used with a regular monitoring of cardiac and renal function.

For patients with acute and unstable heart failure, Trajenta Duo is contraindicated due to the metformin component (see section 4.3).

Surgery

Metformin must be discontinued at the time of surgery under general, spinal or epidural anaesthesia. Therapy may be restarted no earlier than 48 hours following surgery or resumption of oral nutrition and provided that renal function has been re-evaluated and found to be stable.

Elderly

Caution should be exercised when treating patients 80 years and older (see section 4.2).

Change in clinical status of patients with previously controlled type 2 diabetes

As Trajenta Duo contains metformin, a patient with previously well controlled type 2 diabetes on Trajenta Duo who develops laboratory abnormalities or clinical illness (especially vague and poorly defined illness) should be evaluated promptly for evidence of ketoacidosis or lactic acidosis. Evaluation should include serum electrolytes and ketones, blood glucose and, if indicated, blood pH, lactate, pyruvate, and metformin levels. If acidosis of either form occurs, Trajenta Duo must be stopped immediately and other appropriate corrective measures initiated.

Acute Pancreatitis

Use of DPP-4 inhibitors has been associated with a risk of developing acute pancreatitis. Acute pancreatitis has been observed in patients taking linagliptin. In a cardiovascular and renal safety study (CARMELINA) with median observation period of 2.2 years, adjudicated acute pancreatitis was reported in 0.3% of patients treated with linagliptin and in 0.1% of patients treated with placebo. Patients should be informed of the characteristic symptom of acute pancreatitis. If pancreatitis is suspected, Trajenta Duo should be discontinued; if acute pancreatitis is confirmed, Trajenta Duo should not be restarted. Caution should be exercised in patients with a history of pancreatitis.

Bullous pemphigoid

Bullous pemphigoid has been observed in patients taking linagliptin. If bullous pemphigoid is suspected, Trajenta Duo should be discontinued.

Vitamin B12

The risk of low vitamin B12 levels increases with increasing metformin dose, treatment duration, and/or in patients with risk factors known to cause vitamin B12 deficiency. In case of suspicion of vitamin B12 deficiency (such as anaemia or neuropathy), vitamin B12 serum levels should be monitored. Periodic vitamin B12 monitoring could be necessary in patients with risk factors for vitamin B12 deficiency. Metformin therapy should be continued for as long as it is tolerated and not contra-indicated and appropriate corrective treatment for vitamin B12 deficiency provided in line with current clinical guidelines.

4.5 Interaction with other medicinal products and other forms of interaction

Pharmacokinetic drug interaction studies with Trajenta Duo have not been performed; however, such studies have been conducted with the individual active substances, i.e. linagliptin and metformin. Co-administration of multiple doses of linagliptin and metformin did not meaningfully alter the pharmacokinetics of either linagliptin or metformin in healthy volunteers and patients.

Linagliptin

In vitro assessment of interactions

Linagliptin is a weak competitive and a weak to moderate mechanism-based inhibitor of CYP isozyme CYP3A4, but does not inhibit other CYP isozymes. It is not an inducer of CYP isozymes.

Linagliptin is a P-glycoprotein substrate, and inhibits P-glycoprotein mediated transport of digoxin with low potency. Based on these results and *in vivo* drug interaction studies, linagliptin is considered unlikely to cause interactions with other P-gp substrates.

In vivo assessment of interactions

Effects of other medicinal products on linagliptin

Clinical data described below suggest that the risk for clinically meaningful interactions by coadministered medicinal products is low.

Metformin:

Co-administration of multiple three-times-daily doses of 850 mg metformin hydrochloride with 10 mg linagliptin once daily did not clinically meaningfully alter the pharmacokinetics of linagliptin in healthy subjects.

Sulphonylureas:

The steady-state pharmacokinetics of 5 mg linagliptin was not changed by concomitant administration of a single 1.75 mg dose glibenclamide (glyburide).

Ritonavir:

Co-administration of a single 5 mg oral dose of linagliptin and multiple 200 mg oral doses of ritonavir, a potent inhibitor of P-glycoprotein and CYP3A4, increased the AUC and C_{max} of linagliptin approximately twofold and threefold, respectively. The unbound concentrations, which are usually less than 1% at the therapeutic dose of linagliptin, were increased 4-5-fold after co-administration with ritonavir. Simulations of steady-state plasma concentrations of linagliptin with and without ritonavir indicated that the increase in exposure will not be associated with an increased accumulation. These changes in linagliptin pharmacokinetics were not considered to be clinically relevant. Therefore, clinically relevant interactions would not be expected with other P-glycoprotein/CYP3A4 inhibitors.

Rifampicin:

Multiple co-administration of 5 mg linagliptin with rifampicin, a potent inducer of P-glycoprotein and CYP3A4, resulted in a 39.6% and 43.8% decreased linagliptin steady-state AUC and C_{max} respectively, and about 30% decreased DPP-4 inhibition at trough. Thus full efficacy of linagliptin in combination with strong P-gp inducers might not be achieved, particularly if these are administered long-term. Co-administration with other potent inducers of P-glycoprotein and CYP3A4, such as carbamazepine, phenobarbital and phenytoin has not been studied.

Effects of linagliptin on other medicinal products

In clinical studies, as described below, linagliptin had no clinically relevant effect on the pharmacokinetics of metformin, glyburide, simvastatin, warfarin, digoxin or oral contraceptives providing *in vivo* evidence of a low propensity for causing interactions with substrates of CYP3A4, CYP2C9, CYP2C8, P-glycoprotein, and organic cationic transporter (OCT).

Metformin:

Co-administration of multiple daily doses of 10 mg linagliptin with 850 mg metformin hydrochloride, an OCT substrate, had no relevant effect on the pharmacokinetics of metformin in healthy subjects. Therefore, linagliptin is not an inhibitor of OCT-mediated transport.

Sulphonylureas:

Co-administration of multiple oral doses of 5 mg linagliptin and a single oral dose of 1.75 mg glibenclamide (glyburide) resulted in clinically not relevant reduction of 14% of both AUC and C_{max} of glibenclamide. Because glibenclamide is primarily metabolised by CYP2C9, these data also support the conclusion that linagliptin is not a CYP2C9 inhibitor. Clinically meaningful interactions would not be expected with other sulphonylureas (e.g., glipizide, tolbutamide, and glimepiride) which, like glibenclamide, are primarily eliminated by CYP2C9.

Digoxin:

Co-administration of multiple daily doses of 5 mg linagliptin with multiple doses of 0.25 mg digoxin had no effect on the pharmacokinetics of digoxin in healthy subjects. Therefore, linagliptin is not an inhibitor of P-glycoprotein-mediated transport *in vivo*.

Warfarin:

Multiple daily doses of 5 mg linagliptin did not alter the pharmacokinetics of S(-) or R(+) warfarin, a CYP2C9 substrate, administered in a single dose.

Simvastatin:

Multiple daily doses of linagliptin had a minimal effect on the steady state pharmacokinetics of simvastatin, a sensitive CYP3A4 substrate, in healthy subjects. Following administration of a supratherapeutic dose of 10 mg linagliptin concomitantly with 40 mg of simvastatin daily for 6 days, the plasma AUC of simvastatin was increased by 34%, and the plasma C_{max} by 10%.

Oral contraceptives:

Co-administration with 5 mg linagliptin did not alter the steady-state pharmacokinetics of levonorgestrel or ethinylestradiol.

Metformin

Combination requiring precautions for use

Glucocorticoids (given by systemic and local routes), beta-2-agonists, and diuretics have intrinsic hyperglycaemic activity. The patient should be informed and more frequent blood glucose monitoring performed, especially at the beginning of treatment with such medicinal products. If necessary, the dose of the anti-hyperglycaemic medicinal product should be adjusted during therapy with the other medicinal product and on its discontinuation.

Some medicinal products can adversely affect renal function which may increase the risk of lactic acidosis, e.g. NSAIDs, including selective cyclo-oxygenase (COX) II inhibitors, ACE inhibitors, angiotensin II receptor antagonists and diuretics, especially loop diuretics. When starting or using such products in combination with metformin, close monitoring of renal function is necessary.

Combinations not recommended

Alcohol

Alcohol intoxication is associated with an increased risk of lactic acidosis, particularly in cases of fasting, malnutrition or hepatic impairment.

Iodinated contrast agents

Trajenta Duo must be discontinued prior to, or at the time of the imaging procedure and not to be restarted until at least 48 hours after, provided that renal function has been re-evaluated and found to be stable (see section 4.2 and 4.4).

Organic cation transporters (OCT)

Metformin is a substrate of both transporters OCT1 and OCT2. Co-administration of metformin with:

- Inhibitors of OCT1 (such as verapamil) may reduce efficacy of metformin.
- Inducers of OCT1 (such as rifampicin) may increase gastrointestinal absorption and efficacy of metformin.
- Inhibitors of OCT2 (such as cimetidine, dolutegravir, ranolazine, trimethoprim, vandetanib, isavuconazole) may decrease the renal elimination of metformin and thus lead to an increase in metformin plasma concentration.
- Inhibitors of both OCT1 and OCT2 (such as crizotinib, olaparib) may alter efficacy and renal elimination of metformin.

Caution is therefore advised, especially in patients with renal impairment, when these drugs are co-administered with metformin, as metformin plasma concentration may increase. If needed, dose adjustment of metformin may be considered as OCT inhibitors/inducers may alter the efficacy of metformin.

4.6 Fertility, pregnancy and lactation

Pregnancy

The use of linagliptin has not been studied in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3).

A limited amount of data suggests that the use of metformin in pregnant women is not associated with an increased risk of congenital malformations. Animal studies with metformin do not indicate harmful effects with respect to pregnancy, embryonic or foetal development, parturition or postnatal development (see section 5.3).

Non-clinical reproduction studies did not indicate an additive teratogenic effect attributed to the co-administration of linagliptin and metformin.

Trajenta Duo should not be used during pregnancy. If the patient plans to become pregnant, or if pregnancy occurs, treatment with Trajenta Duo should be discontinued and switched to insulin treatment as soon as possible in order to lower the risk of foetal malformations associated with abnormal blood glucose levels.

Breast-feeding

Studies in animals have shown excretion of both metformin and linagliptin into milk in lactating rats. Metformin is excreted in human milk in small amounts. It is not known whether linagliptin is excreted into human milk. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from Trajenta Duo therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

Fertility

No studies on the effect on human fertility have been conducted for Trajenta Duo. No adverse effects of linagliptin on fertility were observed in male or female rats (see section 5.3).

4.7 Effects on ability to drive and use machines

Trajenta Duo has no or negligible influence on the ability to drive and use machines. However, patients should be alerted to the risk of hypoglycaemia when Trajenta Duo is used in combination with other antidiabetic medicinal products known to cause hypoglycaemia (e.g. sulphonylureas).

4.8 Undesirable effects

Summary of the safety profile

The safety of linagliptin 2.5 mg twice daily (or its bioequivalent of 5 mg once daily) in combination with metformin has been evaluated in over 6800 patients with type 2 diabetes mellitus (T2DM).

In placebo-controlled studies, more than 1800 patients were treated with the therapeutic dose of either 2.5 mg linagliptin twice daily (or its bioequivalent of 5 mg linagliptin once daily) in combination with metformin for $\geq 12/24$ weeks.

In the pooled analysis of the seven placebo-controlled trials, the overall incidence of adverse events (AEs) in patients treated with placebo and metformin was comparable to linagliptin 2.5 mg and metformin (54.3% and 49.0%). Discontinuation of therapy due to AEs was comparable in patients who received placebo and metformin to patients treated with linagliptin and metformin (3.8% and 2.9%).

Due to the impact of the background therapy on AEs (e.g. on hypoglycaemias), adverse events were analysed and displayed based on the respective treatment regimens, add-on to metformin, and add-on to metformin plus sulphonylurea.

The placebo-controlled studies included 7 studies where linagliptin was given as add-on to metformin and 1 study where linagliptin was given as add-on to metformin + sulphonylurea.

Tabulated summary of adverse reactions

The following adverse reactions derived from the use of the linagliptin/metformin combination or the use of the monocomponents (linagliptin or metformin) in clinical trials or from post-marketing experience are shown in the table below. Undesirable effects previously reported with one of the individual components may be potential undesirable effects with TRAJENTA DUO even if not observed in clinical trials with this product.

The adverse reactions are listed by system organ class and absolute frequency. Frequencies are defined as very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), or very rare ($< 1/10,000$) and not known (cannot be estimated from the available data).

MedDRA System Organ Class terminology	Linagliptin and Metformin Adverse reactions	Frequencies
Infections and infestations	Nasopharyngitis ^{1; 3}	Uncommon

Immune system disorders	Hypersensitivity ^{1; 3} Angioedema ⁴ Urticaria ^{2; 4}	Uncommon Rare [§] Rare [§]
Metabolism and nutrition disorders	Lactic acidosis ² Vitamin B ₁₂ decrease/deficiency ² Hypoglycaemia (when linagliptin and metformin were combined with a sulphonylurea)	Very Rare Common Very common
Nervous system disorders	Taste disturbance ²	Common
Respiratory, thoracic and mediastinal disorders	Cough ^{1; 3}	Uncommon
Gastrointestinal disorders***	Decreased appetite ^{3; 5} Diarrhoea ^{3; 5} Nausea ^{3; 5} Pancreatitis ³ Vomiting ^{3; 5} Constipation (when linagliptin and metformin were combined with insulin) Abdominal pain ² Mouth ulceration ⁴	Uncommon Common Uncommon Rare ^Ω Uncommon Uncommon Very common Rare
Hepato-biliary disorders	Liver function test abnormal ² Hepatitis ²	Very Rare Very Rare
Skin and subcutaneous tissue disorders	Pruritus ^{3; 5} Erythema ² Rash ⁴ Bullous pemphigoid ^{4; §}	Uncommon Very Rare Uncommon Rare ^Ω
Investigations	Lipase increased ^{3;†} Amylase increased ^ψ	Common Uncommon

¹ Adverse reactions reported also in patients treated with linagliptin monotherapy

² Adverse reactions of metformin as monotherapy; refer to the Summary of Product Characteristics for metformin for additional information

³ Adverse reactions of FDC linagliptin + metformin (pooled analysis of placebo-controlled studies)

⁴ Adverse reactions identified from post-marketing experience with linagliptin

⁵ Adverse reactions reported in patients who received FDC linagliptin + metformin and also in patients who received metformin monotherapy

*** Gastrointestinal disorders such as abdominal pain and nausea, vomiting, diarrhoea, decreased appetite occur most frequently during initiation of therapy with metformin hydrochloride and resolve spontaneously in most cases. To prevent them, it is recommended that metformin hydrochloride be taken in 2 daily doses during or after meals if administered as monotherapy.

§ See also *Linagliptin cardiovascular and renal safety study (CARMELINA)* below

† Based on lipase increase >3 ULN

ψ In the CAROLINA study comparing linagliptin with active comparator glimepiride (see section clinical trials) laboratory analysis of amylase showed increases to >3xULN in 0.99% of patients treated with linagliptin and in 0.54% of patients treated with glimepiride

§ ADR based on post-marketing experience, for which no drug-related AEs were reported in the reference dataset (the pool of trials with Linagliptin and metformin alone or with various backgrounds was used to represent post marketing conditions). As suggested in European SmPC Guideline, in case of no reported events, the frequency category was derived from estimated worst case frequency of 3/X, with X representing the total number of patients in the reference dataset.

Ω Frequency based on CARMELINA study (1218.22). Frequency for pancreatitis was calculated based on adjudication-confirmed, drug-related (by investigator assessment) events n=3/3494 (0.086%) resulting in frequency 'rare' (source: CTR

listing 16.2.7.8). Frequency for bullous pemphigoid was calculated based on drug-related (by investigator assessment) events n=3/3494 (0.086%) resulting in frequency 'rare' (source: CTR table 15.3.1.3.20).

In placebo-controlled studies the most frequently reported adverse reaction for linagliptin plus metformin was diarrhoea (1.6%) with a comparable rate on metformin plus placebo (2.4%).

Adverse reactions reported when linagliptin and metformin were combined with sulphonylurea

When linagliptin and metformin were administered in combination with a sulphonylurea, hypoglycaemia was the most commonly reported adverse reaction (linagliptin plus metformin plus sulphonylurea 23.9% vs 16.0% in the placebo group) and identified as an additional adverse reaction under these conditions. None of the hypoglycaemias episodes were classified as severe (requiring assistance).

Adverse reactions reported when linagliptin and metformin were combined with insulin

When linagliptin and metformin were administered in combination with insulin, hypoglycaemia was the most frequently reported adverse reactions, but occurred at comparable rate when placebo and metformin were combined with insulin (linagliptin plus metformin plus insulin 29.5% versus 30.9% in the placebo plus metformin plus insulin group) with a low rate of severe (requiring assistance) episodes (1.5% vs 0.9%).

Linagliptin cardiovascular outcome and renal safety study (CARMELINA)

The CARMELINA study evaluated the cardiovascular and renal safety of linagliptin versus placebo in patients with type 2 diabetes and with increased CV risk evidenced by a history of established macrovascular or renal disease (see section Clinical Trials). The study included 3494 patients treated with linagliptin (5 mg) and 3485 patients treated with placebo. Both treatments were added to standard of care targeting regional standards for HbA1c and CV risk factors; with 54% on metformin. The overall incidence of adverse events and serious adverse events in patients receiving linagliptin was similar to that in patients receiving placebo. Safety data from this study was in line with previous known safety profile of linagliptin.

In the treated population, severe hypoglycaemic events (requiring assistance) were reported in 3.0% of patients on linagliptin and in 3.1% on placebo. Among patients who were using sulphonylurea at baseline, the incidence of severe hypoglycaemia was 2.0% in linagliptin-treated patients and 1.7% in placebo treated patients. Among patients who were using insulin at baseline, the incidence of severe hypoglycaemia was 4.4% in linagliptin-treated patients and 4.9% in placebo treated patients.

In the overall study observation period adjudicated acute pancreatitis was reported in 0.3% of patients treated with linagliptin and in 0.1% of patients treated with placebo.

In the CARMELINA study, bullous pemphigoid was reported in 0.2% of patients treated with linagliptin and in no patient treated with placebo.

4.9 Overdose

Linagliptin

During controlled clinical trials in healthy subjects, single doses of up to 600 mg linagliptin (equivalent to 120 times the recommended dose) were not associated with a dose dependent increase in adverse events. There is no experience with doses above 600 mg in humans.

Metformin

Hypoglycaemia has not been seen with metformin hydrochloride doses of up to 85 g, although lactic acidosis has occurred in such circumstances. High overdose of metformin hydrochloride or concomitant

risks may lead to lactic acidosis. Lactic acidosis is a medical emergency and must be treated in hospital. The most effective method to remove lactate and metformin hydrochloride is haemodialysis.

Management

In the event of an overdose, it is reasonable to employ the usual supportive measures, e.g. remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring, and institute clinical measures if required.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Drugs used in diabetes, combinations of oral blood glucose lowering drugs, ATC code: A10BD11.

Mechanism of action and pharmacodynamic effects

Trajenta Duo combines two antihyperglycaemic medicinal products with complementary mechanisms of action to improve glycaemic control in patients with type 2 diabetes: linagliptin, a dipeptidyl peptidase 4 (DPP-4) inhibitor, and metformin hydrochloride, a member of the biguanide class.

Linagliptin

Linagliptin is an inhibitor of the enzyme DPP-4 (Dipeptidyl peptidase 4) an enzyme which is involved in the inactivation of the incretin hormones GLP-1 and GIP (glucagon-like peptide-1, glucose-dependent insulinotropic polypeptide). These hormones are rapidly degraded by the enzyme DPP-4. Both incretin hormones are involved in the physiological regulation of glucose homeostasis. Incretins are secreted at a low basal level throughout the day and levels rise immediately after meal intake. GLP-1 and GIP increase insulin biosynthesis and secretion from pancreatic beta cells in the presence of normal and elevated blood glucose levels. Furthermore GLP-1 also reduces glucagon secretion from pancreatic alpha cells, resulting in a reduction in hepatic glucose output. Linagliptin binds very effectively to DPP-4 in a reversible manner and thus leads to a sustained increase and a prolongation of active incretin levels. Linagliptin glucose-dependently increases insulin secretion and lowers glucagon secretion thus resulting in an overall improvement in the glucose homeostasis. Linagliptin binds selectively to DPP-4 and exhibits a > 10,000 fold selectivity *versus* DPP-8 or DPP-9 activity *in vitro*.

Metformin

Metformin is a biguanide with antihyperglycaemic effects, on both basal and postprandial hyperglycaemia. It does not stimulate insulin secretion and therefore does not cause hypoglycaemia. Metformin reduces basal hyperinsulinemia, and in combination with insulin, reduces insulin requirement.

Metformin exerts its antihyperglycaemic effect via multiple mechanisms: Metformin reduces hepatic glucose production. Metformin facilitates peripheral glucose uptake and utilization, in part by increasing insulin action. Metformin alters glucose turnover in the gut: Uptake from circulation is increased and absorption from food is decreased.

Additional mechanisms attributed to the gut include an increase in release of glucagon-like peptide 1 (GLP-1) and a decrease of bile acid resorption. Metformin alters the gut microbiome.

Metformin can improve the lipid profile in hyperlipidaemic individuals.

In clinical studies, use of metformin was associated with either a stable body weight or modest weight loss. Metformin is an adenosine monophosphate-protein-kinase (AMPK) activator and increases the transport capacity of all types of membrane glucose transporters (GLUTs).

CLINICAL STUDIES

Linagliptin as add-on to metformin therapy

The efficacy and safety of linagliptin in combination with metformin in patients with insufficient glycaemic control on metformin monotherapy was evaluated in a double-blind placebo-controlled study of 24 weeks duration. Linagliptin added to metformin provided significant improvements in HbA1c, (-0.64% change compared to placebo), from a mean baseline HbA1c of 8%. Linagliptin also showed significant improvements in fasting plasma glucose (FPG) by -21.1 mg/dl and 2-hour post-prandial glucose (PPG) by -67.1 mg/dl compared to placebo, as well as a greater portion of patients achieving a target HbA1c of < 7.0% (28.3% on linagliptin *versus* 11.4% on placebo). The observed incidence of hypoglycaemia in patients treated with linagliptin was similar to placebo. Body weight did not differ significantly between the groups.

In a 24-week placebo-controlled factorial study of initial therapy, linagliptin 2.5 mg twice daily in combination with metformin (500 mg or 1,000 mg twice daily) provided significant improvements in glycaemic parameters compared with either monotherapy as summarised in Table 2 (mean baseline HbA1c 8.65%).

Table 2: Glycaemic Parameters at Final Visit (24-Week Study) for Linagliptin and Metformin, Alone and in Combination in Patients with Type 2 Diabetes Mellitus Inadequately Controlled on Diet and Exercise

	Placebo	Linagliptin 5 mg Once Daily¹	Metformin HCl 500 mg Twice Daily	Linagliptin 2.5 mg Twice Daily¹ + Metformin HCl 500 mg Twice Daily	Metformin HCl 1,000 mg Twice Daily	Linagliptin 2.5 mg Twice Daily¹ + Metformin HCl 1,000 mg Twice Daily
HbA1c (%)						
Number of patients	n = 65	n = 135	n = 141	n = 137	n = 138	n = 140
Baseline (mean)	8.7	8.7	8.7	8.7	8.5	8.7
Change from baseline (adjusted mean)	0.1	-0.5	-0.6	-1.2	-1.1	-1.6
Difference from placebo (adjusted mean) (95% CI)	--	-0.6 (-0.9, -0.3)	-0.8 (-1.0, -0.5)	-1.3 (-1.6, -1.1)	-1.2 (-1.5, -0.9)	-1.7 (-2.0, -1.4)
Patients (n, %) achieving HbA1c <7%	7 (10.8)	14 (10.4)	27 (19.1)	42 (30.7)	43 (31.2)	76 (54.3)
Patients (%) receiving rescue	29.2	11.1	13.5	7.3	8.0	4.3

treatment						
FPG (mg/dL)						
Number of patients	n = 61	n = 134	n = 136	n = 135	n = 132	n = 136
Baseline (mean)	203	195	191	199	191	196
Change from baseline (adjusted mean)	10	-9	-16	-33	-32	-49
Difference from placebo (adjusted mean) (95% CI)	--	-19 (-31, -6)	-26 (-38, -14)	-43 (-56, -31)	-42 (-55, -30)	-60 (-72, -47)

¹ Total daily dose of linagliptin is equal to 5 mg

Mean reductions from baseline in HbA_{1c} were generally greater for patients with higher baseline HbA_{1c} values. Effects on plasma lipids were generally neutral. The decrease in body weight with the combination of linagliptin and metformin was similar to that observed for metformin alone or placebo; there was no change in weight from baseline for patients on linagliptin alone. The incidence of hypoglycaemia was similar across treatment groups (placebo 1.4%, linagliptin 5 mg 0%, metformin 2.1%, and linagliptin 2.5 mg plus metformin twice daily 1.4%).

The efficacy and safety of linagliptin 2.5 mg twice daily *versus* 5 mg once daily in combination with metformin in patients with insufficient glycaemic control on metformin monotherapy was evaluated in a double-blind placebo-controlled study of 12 weeks duration. Linagliptin 5 mg once daily and 2.5 mg twice daily provided comparable (CI: -0.07; 0.19) significant HbA_{1c} reductions of -0.80% (from baseline 7.98%), and -0.74% (from baseline 7.96%) compared to placebo. The observed incidence of hypoglycaemia in patients treated with linagliptin was similar to placebo. Body weight did not differ significantly between the groups.

Linagliptin as add-on to a combination of metformin and sulphonylurea therapy

A placebo-controlled study of 24 weeks in duration was conducted to evaluate the efficacy and safety of linagliptin 5 mg to placebo, in patients not sufficiently treated with a combination with metformin and a sulphonylurea. Linagliptin provided significant improvements in HbA_{1c} (-0.62% change compared to placebo), from a mean baseline HbA_{1c} of 8.14%. Linagliptin also showed significant improvements in patients achieving a target HbA_{1c} of < 7.0% (31.2% on linagliptin *versus* 9.2% on placebo), and also for fasting plasma glucose (FPG) with -12.7 mg/dl reduction compared to placebo. Body weight did not differ significantly between the groups.

Linagliptin as add on to a combination of metformin and empagliflozin

In patients inadequately controlled with metformin and empagliflozin (10 mg (n=247) or 25 mg (n=217)), 24-weeks treatment with add-on therapy of linagliptin 5 mg provided adjusted mean HbA_{1c} reductions from baseline by -0.53% (significant difference to add-on placebo -0.32% (95% CI -0.25, -0.13) and -0.58% (significant difference to add-on placebo -0.47% (95% CI -0.66; -0.28), respectively. A statistically significant greater proportion of patients with a baseline HbA_{1c} ≥7.0% and treated with linagliptin 5 mg achieved a target HbA_{1c} of <7% compared to placebo.

In prespecified subgroups of patients with baseline HbA_{1c} greater or equal than 8.5% (n=66 and n=42 patients on metformin plus empagliflozin 10 mg or 25 mg, respectively), the adjusted mean HbA_{1c} reductions from baseline to 24 weeks on add-on therapy with linagliptin 5 mg were -0.97% (p=0.0875, for difference to add-on placebo) and -1.16% (p=0.0046 for difference to add-on placebo), respectively.

Linagliptin in combination with metformin and insulin

A 24-week placebo-controlled study was conducted to evaluate the efficacy and safety of linagliptin (5mg once daily) added to insulin with or without metformin. 83% of patients were taking metformin in combination with insulin in this trial. Linagliptin in combination of metformin plus insulin provided significant improvements in HbA_{1c} in this subgroup with -0.68% (CI: -0.78; -0.57) adjusted mean change from baseline (mean baseline HbA_{1c} 8.28%) compared to placebo in combination with metformin plus insulin. There was no meaningful change from baseline in body weight in either group.

Linagliptin 24 month data, as add-on to metformin in comparison with glimepiride

In a study comparing the efficacy and safety of the addition of linagliptin 5 mg or glimepiride (mean dose 3 mg) in patients with inadequate glycaemic control on metformin monotherapy, mean reductions in HbA_{1c} were -0.16% with linagliptin (mean baseline HbA_{1c} 7.69%) and -0.36% with glimepiride (mean baseline HbA_{1c} 7.69%) with a mean treatment difference of 0.20% (97.5% CI: 0.09, 0.299). The incidence of hypoglycaemia in the linagliptin group (7.5%) was significantly lower than that in the glimepiride group (36.1%). Patients treated with linagliptin exhibited a significant mean decrease from baseline in body weight compared to a significant weight gain in patients administered glimepiride (-1.39 versus +1.29 kg).

Linagliptin as add-on therapy in elderly patients (age ≥ 70 years) with type 2 diabetes

The efficacy and safety of linagliptin in elderly (age ≥ 70 years) type 2 diabetes patients was evaluated in a double-blind study of 24 weeks duration. Patients received metformin and/or sulphonylurea and/or insulin as background therapy. Doses of background antidiabetic medications were kept stable during the first 12 weeks, after which adjustments were permitted. Linagliptin provided significant improvements in HbA_{1c} (-0.64% change compared to placebo after 24 weeks), from a mean baseline HbA_{1c} of 7.8%. Linagliptin also showed significant improvements in fasting plasma glucose (FPG) compared to placebo. Body weight did not differ significantly between the groups.

In a pooled analysis of elderly (age ≥ 70 years) patients with type 2 diabetes (n=183) who were taking both metformin and basal insulin as background therapy, linagliptin in combination with metformin plus insulin provided significant improvements in HbA_{1c} parameters with -0.81% (CI: -1.01; -0.61) adjusted mean change from baseline (mean baseline HbA_{1c} 8.13%) compared to placebo in combination with metformin plus insulin.

Linagliptin and initial combination with Linagliptin and Metformin in recently diagnosed treatment naïve patients with marked hyperglycaemia:

The efficacy and safety of the initial combination of linagliptin 5 mg once daily and metformin twice daily (up-titrated in the first 6 weeks to 1500 mg or 2000 mg/d) compared to linagliptin 5 mg once has been studied in a 24 week trial in recently diagnosed treatment naïve patients with type 2 diabetes mellitus and marked hyperglycaemia (baseline HbA_{1c} 8.5–12.0%). After 24 weeks both linagliptin monotherapy as well as the initial combination of linagliptin and metformin significantly reduced HbA_{1c} levels by -2.0% and -2.8% respectively, from a baseline HbA_{1c} of 9.9% and 9.8% respectively. The treatment difference of -0.8% (95% CI -1.1 to -0.5) showed superiority for the initial combination over monotherapy (p<0.0001). Notably, 40% and 61% of patients in the monotherapy and combination arms achieved HbA_{1c} <7.0%.

Linagliptin cardiovascular and renal safety study (CARMELINA)

CARMELINA was a randomized study in 6979 patients with type 2 diabetes with increased CV risk evidenced by a history of established macrovascular or renal disease who were treated with linagliptin 5 mg (3494) or placebo (3485) added to standard of care targeting regional standards for HbA_{1c}, CV risk factors and renal disease. The study population included 1,211 (17.4%) patients ≥ 75 years of age and

4,348 (62.3%) patients with renal impairment. Approximately 19% of the population had eGFR ≥ 45 to < 60 mL/min/1.73 m², 28% of the population had eGFR ≥ 30 to < 45 mL/min/1.73 m² and 15% had eGFR < 30 mL/min/1.73m². The mean HbA_{1c} at baseline was 8.0%.

The study was designed to demonstrate non-inferiority for the primary cardiovascular endpoint which was a composite of the first occurrence of cardiovascular death or a non-fatal myocardial infarction (MI) or a non-fatal stroke (3P-MACE). The renal composite endpoint was defined as renal death or sustained end stage renal disease or sustained decrease of 40% or more in eGFR.

After a median follow up of 2.2 years, linagliptin, when added to standard of care, did not increase the risk of major adverse cardiovascular events or renal outcome events (Table 3 and Figure 1). There was no increased risk in hospitalization for heart failure which was an additional adjudicated endpoint observed compared to standard of care without linagliptin in patients with type 2 diabetes (Table 4).

Table 3: Major adverse cardiovascular events (MACE) and renal outcome events by treatment group in the CARMELINA study

	Linagliptin 5mg		Placebo		Hazard Ratio (95% CI)
	Number of Subjects (%)	Incidence Rate per 1000 PY*	Number of Subjects (%)	Incidence Rate per 1000 PY*	
Number of patients	3494		3485		
Primary CV composite (Cardiovascular death, non-fatal MI, non-fatal stroke)	434 (12.4)	57.7	420 (12.1)	56.3	1.02 (0.89, 1.17) **
Secondary renal composite (renal death, ESRD, 40% sustained decrease in eGFR)	327 (9.4)	48.9	306 (8.8)	46.6	1.04 (0.89, 1.22

*PY=patient years

** Test on non-inferiority to demonstrate that the upper bound of the 95% CI for the hazard ratio is less than 1.3

Figure 1: Time to first occurrence of 3P-MACE in CARMELINA




Table 4: Hospitalization for heart failure and mortality by treatment group in the CARMELINA study

	Linagliptin 5mg		Placebo		Hazard Ratio (95% CI)
	Number of Subjects (%)	Incidence Rate per 1000 PY*	Number of Subjects (%)	Incidence Rate per 1000 PY*	
Number of patients	3494		3485		
All-cause mortality	367 (10.5)	46.9	373 (10.7)	48.0	0.98 (0.84, 1.13)
CV death	255 (7.3)	32.6	264 (7.6)	34	0.96 (0.81, 1.14)
Hospitalization for heart failure	209 (6.0)	27.7	226 (6.5)	30.4	0.90 (0.74, 1.08)

*PY=patient years

In analyses for albuminuria progression (change from normoalbuminuria to micro- or macroalbuminuria, or from microalbuminuria to macroalbuminuria) the estimated hazard ratio was 0.86 (95% CI 0.78, 0.95) for linagliptin versus placebo. The microvascular endpoint was defined as the composite of renal death, sustained ESRD, sustained decrease of $\geq 50\%$ in eGFR, albuminuria progression, use of retinal photocoagulation or intravitreal injections of an anti-VEGF therapy for diabetic retinopathy or vitreous haemorrhage or diabetes-related-blindness. The estimated hazard ratio for time to first occurrence for the composite microvascular endpoint was 0.86 (95% CI 0.78, 0.95) for linagliptin versus placebo, mainly driven by albuminuria progression.

Linagliptin cardiovascular safety study (CAROLINA)

CAROLINA was a randomized study in 6033 patients with early type 2 diabetes and increased CV risk or established complications who were treated with linagliptin 5 mg (3023) or glimepiride 1-4 mg (3010)

added to standard of care (including background therapy with metformin in 83% of patients) targeting regional standards for HbA_{1c} and CV risk factors. The mean age for study population was 64 years and included 2030 (34%) patients ≥ 70 years of age. The study population included 2089 (35%) patients with cardiovascular disease and 1130 (19%) patients with renal impairment with an eGFR $< 60\text{ml/min/1.73m}^2$ at baseline. The mean HbA_{1c} at baseline was 7.15%.

The study was designed to demonstrate non-inferiority for the primary cardiovascular endpoint which was a composite of the first occurrence of cardiovascular death or a non-fatal myocardial infarction (MI) or a non-fatal stroke (3P-MACE).

After a median follow up of 6.25 years, linagliptin, when added to standard of care, did not increase the risk of major adverse cardiovascular events (Table 5) as compared to glimepiride. Results were consistent for patients treated with or without metformin.

Table 5: Major adverse cardiovascular events (MACE) and mortality by treatment group in the CAROLINA study

	Linagliptin 5mg		Glimepiride (1-4mg)		Hazard Ratio (95% CI)
	Number of Subjects (%)	Incidence Rate per 1000 PY*	Number of Subjects (%)	Incidence Rate per 1000 PY*	
Number of patients	3023		3010		
Primary CV composite (Cardiovascular death, non-fatal MI, non-fatal stroke)	356 (11.8)	20.7	362 (12.0)	21.2	0.98 (0.84, 1.14)**

* PY=patient years

** Test on non-inferiority to demonstrate that the upper bound of the 95% CI for the hazard ratio is less than 1.3

The composite of treatment sustainability, a key secondary endpoint, was defined as the proportion of patients on study treatment following initial titration period (16 weeks) that maintain glycaemic control (HbA_{1c} $\leq 7.0\%$) at final visit without need for additional antidiabetic drug therapy (rescue medication) without any moderate (symptomatic with glucose value $\leq 70\text{mg/dL}$) or severe (requiring assistance) hypoglycaemic episodes and without $> 2\%$ weight gain. A higher number of patients on linagliptin (481, 16.0%) achieved this key secondary endpoint compared to glimepiride (305, 10.2%).

For the entire treatment period (median time on treatment 5.9 years) the rate of patients with moderate or severe hypoglycaemia was 6.5% on linagliptin versus 30.9% on glimepiride, severe hypoglycaemia occurred in 0.3% of patients on linagliptin versus 2.2% on glimepiride.

Paediatric population

The clinical efficacy and safety of empagliflozin 10 mg with a possible dose-increase to 25 mg or linagliptin 5mg once daily has been studied in children and adolescents from 10 to 17 years of age with T2DM in a double-blind, randomised, placebo-controlled, parallel group study (DINAMO) over 26 weeks, with a double-blind active treatment safety extension period up to 52 weeks.

A total of 157 patients were treated with either empagliflozin (10 mg or 25 mg; N=52), linagliptin (N=52) or placebo (N=53). Background therapies as adjunct to diet and exercise included metformin (51%), a combination of metformin and insulin (40.1%), insulin (3.2%), or none (5.7%). The mean HbA1c was 8.03% at baseline.

The study population consisted of 38.2% male and 61.8% female patients with a mean age of 14.5 years (range: 10-17 years); 51.6% were 15 years of age or older. 49.7% of the study population were White, 5.7% Asian and 31.2% Black/African American. The mean BMI was 36.04 kg/m², the mean body weight was 99.92 kg. Only patients with an eGFR of ≥ 60 ml/min/1.73 m² were included in the DINAMO study.

The primary endpoint of the study was the change in HbA1c from baseline to the end of 26 weeks, regardless of rescue therapy or treatment discontinuation. Treatment with linagliptin 5 mg did not provide significant improvement in HbA1c. The treatment difference of adjusted mean change in HbA1c between linagliptin and placebo was -0.34% (95% CI -0.99, 0.30; p=0.2935). The adjusted mean change in HbA1c from baseline was 0.33% in patients treated with linagliptin and 0.68% in patients treated with placebo.

Overall, the safety profile of linagliptin was similar to that observed in the adult population.

Metformin

The prospective randomised (UKPDS) study has established the long-term benefit of intensive blood glucose control in type 2 diabetes. Analysis of the results for overweight patients treated with metformin after failure of diet alone showed:

- a significant reduction of the absolute risk of any diabetes-related complication in the metformin group (29.8 events/1,000 patient-years) *versus* diet alone (43.3 events/1,000 patient-years), p=0.0023, and *versus* the combined sulphonylurea and insulin monotherapy groups (40.1 events/1,000 patient-years), p=0.0034,
- a significant reduction of the absolute risk of any diabetes-related mortality: metformin 7.5 events/1,000 patient-years, diet alone 12.7 events/1,000 patient-years, p=0.017,
- a significant reduction of the absolute risk of overall mortality: metformin 13.5 events/1,000 patient-years *versus* diet alone 20.6 events/1,000 patient-years, (p=0.011), and *versus* the combined sulphonylurea and insulin monotherapy groups 18.9 events/1,000 patient-years (p=0.021),
- a significant reduction in the absolute risk of myocardial infarction: metformin 11 events/1,000 patient-years, diet alone 18 events/1,000 patient-years, (p=0.01).

5.2 Pharmacokinetic properties

Trajenta Duo

Bioequivalence studies in healthy subjects demonstrated that the Trajenta Duo (linagliptin/metformin hydrochloride) combination tablets are bioequivalent to co-administration of linagliptin and metformin hydrochloride as individual tablets.

Administration of Trajenta Duo 2.5/1000 mg with food resulted in no change in overall exposure of linagliptin. With metformin there was no change in AUC, however mean peak serum concentration of metformin was decreased by 18% when administered with food. A delayed time to peak serum concentrations by 2 hours was observed for metformin under fed conditions. These changes are not likely to be clinically meaningful.

The following statements reflect the pharmacokinetic properties of the individual active substances of Trajenta Duo.

Linagliptin:

The pharmacokinetics of linagliptin has been extensively characterised in healthy subjects and patients with type 2 diabetes. After oral administration of a 5 mg dose to healthy volunteers or patients, linagliptin was rapidly absorbed, with peak plasma concentrations (median T_{max}) occurring 1.5 hours post-dose.

Plasma concentrations of linagliptin decline in a triphasic manner with a long terminal half-life (terminal half-life for linagliptin more than 100 hours), that is mostly related to the saturable, tight binding of linagliptin to DPP-4 and does not contribute to the accumulation of the drug. The effective half-life for accumulation of linagliptin, as determined from oral administration of multiple doses of 5 mg linagliptin, is approximately 12 hours. After once daily dosing of 5 mg linagliptin, steady-state plasma concentrations are reached by the third dose. Plasma AUC of linagliptin increased approximately 33% following 5 mg doses at steady-state compared to the first dose. The intra-subject and inter-subject coefficients of variation for linagliptin AUC were small (12.6% and 28.5%, respectively). Due to the concentration dependent binding of linagliptin to DPP-4, the pharmacokinetics of linagliptin based on total exposure is not linear; indeed total plasma AUC of linagliptin increased in a less than dose-proportional manner, while unbound AUC increases in a roughly dose proportional manner. The pharmacokinetics of linagliptin was generally similar in healthy subjects and in patients with type 2 diabetes.

Absorption

The absolute bioavailability of linagliptin is approximately 30%. Co-administration of a high-fat meal with linagliptin prolonged the time to reach C_{max} by 2 hours and lowered C_{max} by 15%, but no influence on AUC_{0-72h} was observed. No clinically relevant effect of C_{max} and T_{max} changes is expected; therefore linagliptin may be administered with or without food.

Distribution

As a result of tissue binding, the mean apparent volume of distribution at steady-state following a single 5 mg intravenous dose of linagliptin to healthy subjects is approximately 1110 litres, indicating that linagliptin extensively distributes to the tissues. Plasma protein binding of linagliptin is concentration-dependent, decreasing from about 99% at 1 nmol/l to 75-89% at ≥ 30 nmol/l, reflecting saturation of binding to DPP-4 with increasing concentration of linagliptin. At high concentrations, where DPP-4 is fully saturated, 70-80% of linagliptin was bound to other plasma proteins than DPP-4, hence 30-20% were unbound in plasma.

Biotransformation

Following a [14 C] linagliptin oral 10 mg dose, approximately 5% of the radioactivity was excreted in urine. Metabolism plays a subordinate role in the elimination of linagliptin. One main metabolite with a relative exposure of 13.3% of linagliptin at steady state was detected which was found to be pharmacologically inactive, and thus does not contribute to the plasma DPP-4 inhibitory activity of linagliptin.

Elimination

Following administration of an oral [14 C] linagliptin dose to healthy subjects, approximately 85% of the administered radioactivity was eliminated in faeces (80%) or urine (5%) within 4 days of dosing. Renal clearance at steady state was approximately 70 ml/min.

Special populations

Renal impairment

Under steady-state conditions, linagliptin exposure in patients with mild renal impairment was comparable to healthy subjects. In moderate renal impairment, a moderate increase in exposure of about

1.7 fold was observed compared with control. Exposure in T2DM patients with severe RI was increased by about 1.4 fold compared to T2DM patients with normal renal function. Steady-state predictions for AUC of linagliptin in patients with ESRD indicated comparable exposure to that of patients with moderate or severe renal impairment. In addition, linagliptin is not expected to be eliminated to a therapeutically significant degree by hemodialysis or peritoneal dialysis. No dose adjustment of linagliptin is recommended in patients with renal impairment; therefore, linagliptin may be continued as a single entity tablet at the same total daily dose of 5 mg if Trajenta Duo is discontinued due to evidence of renal impairment.

Hepatic impairment

In patients with mild moderate and severe hepatic impairment (according to the Child-Pugh classification), mean AUC and C_{max} of linagliptin were similar to healthy matched controls following administration of multiple 5 mg doses of linagliptin.

Body Mass Index (BMI)

Body mass index had no clinically relevant effect on the pharmacokinetics of linagliptin based on a population pharmacokinetic analysis of Phase I and Phase II data. The clinical trials before marketing authorization have been performed up to a BMI equal to 40 kg/m².

Gender

Gender had no clinically relevant effect on the pharmacokinetics of linagliptin based on a population pharmacokinetic analysis of Phase I and Phase II data.

Elderly

Age did not have a clinically relevant impact on the pharmacokinetics of linagliptin based on a population pharmacokinetic analysis of Phase I and Phase II data. Elderly subjects (65 to 80 years, oldest patient was 78 years) had comparable plasma concentrations of linagliptin compared to younger subjects. Linagliptin trough concentrations were also measured in elderly (age ≥ 70 years) with type 2 diabetes patients in a phase III study of 24 weeks duration. Linagliptin concentrations in this study were within the range of values previously observed in younger type 2 diabetes patients.

Paediatric population

Pharmacokinetics and pharmacodynamics of 1 mg and 5 mg linagliptin were investigated in children and adolescents 10 to 17 years of age with type 2 diabetes mellitus. The observed pharmacokinetic and pharmacodynamic responses were consistent with those found in adult subjects. Linagliptin 5 mg showed superiority over 1 mg with regard to trough DPP-4 inhibition (72% vs 32%, $p=0.0050$) and a numerically larger reduction with regard to placebo-corrected adjusted mean change from baseline in HbA1c (-0.63% vs -0.48%, n.s.) after 12 weeks.

Pharmacokinetics and pharmacodynamics (HbA1c change from baseline) of 5 mg linagliptin were investigated in children and adolescents 10 to 17 years of age with type 2 diabetes mellitus. The observed exposure-response relationship was overall comparable in paediatric and adult patients. Oral administration of linagliptin resulted in exposure within the range observed in adult patients. The observed geometric mean trough concentrations and geometric mean concentrations at 1.5 hours post-administration at steady state were 4.30 nmol/L and 12.6 nmol/L, respectively.

Race

Race had no obvious effect on the plasma concentrations of linagliptin based on a composite analysis of available pharmacokinetic data, including patients of Caucasian, Hispanic, African, and Asian origin. In addition the pharmacokinetic characteristics of linagliptin were found to be similar in dedicated phase I

studies in Japanese, Chinese and Caucasian healthy subjects and African American type 2 diabetes patients.

Metformin:

Absorption

After an oral dose of metformin, T_{max} is reached in 2.5 hours. Absolute bioavailability of a 500 mg or 850 mg metformin hydrochloride tablet is approximately 50-60% in healthy subjects. After an oral dose, the non-absorbed fraction recovered in faeces was 20-30%.

After oral administration, metformin hydrochloride absorption is saturable and incomplete. It is assumed that the pharmacokinetics of metformin hydrochloride absorption are non-linear.

At the recommended metformin hydrochloride doses and dosing schedules, steady-state plasma concentrations are reached within 24 to 48 hours and are generally less than 1 microgram/ml. In controlled clinical trials, maximum metformin hydrochloride plasma levels (C_{max}) did not exceed 5 microgram/ml, even at maximum doses.

Food decreases the extent and slightly delays the absorption of metformin hydrochloride. Following administration of a dose of 850 mg, a 40% lower plasma peak concentration, a 25% decrease in AUC (area under the curve) and a 35 minute prolongation of the time to peak plasma concentration were observed. The clinical relevance of these decreases is unknown.

Distribution

Plasma protein binding is negligible. Metformin hydrochloride partitions into erythrocytes. The blood peak is lower than the plasma peak and appears at approximately the same time. The red blood cells most likely represent a secondary compartment of distribution. The mean volume of distribution (V_d) ranged between 63-276 l.

Biotransformation

Metformin hydrochloride is excreted unchanged in the urine. No metabolites have been identified in humans.

Elimination

Renal clearance of metformin hydrochloride is > 400 ml/min, indicating that metformin hydrochloride is eliminated by glomerular filtration and tubular secretion. Following an oral dose, the apparent terminal elimination half-life is approximately 6.5 hours.

When renal function is impaired, renal clearance is decreased in proportion to that of creatinine and thus the elimination half-life is prolonged, leading to increased levels of metformin hydrochloride in plasma.

Special populations

Paediatric population

Single dose study: after single doses of metformin hydrochloride 500 mg, paediatric patients have shown a similar pharmacokinetic profile to that observed in healthy adults.

Multiple dose study: data are restricted to one study. After repeated doses of 500 mg twice daily for 7 days in paediatric patients the peak plasma concentration (C_{max}) and systemic exposure (AUC_{0-t}) were reduced by approximately 33% and 40%, respectively compared to diabetic adults who received repeated

doses of 500 mg twice daily for 14 days. As the dose is individually titrated based on glycaemic control, this is of limited clinical relevance.

Renal impairment

The available data in subjects with moderate renal insufficiency are scarce and no reliable estimation of the systemic exposure to metformin in this subgroup as compared to subjects with normal renal function could be made. Therefore, the dose adaptation should be made upon clinical efficacy/tolerability considerations (see section 4.2).

5.3 Preclinical safety data

Linagliptin plus metformin

General toxicity studies in rats for up to 13 weeks were performed with the co-administration of linagliptin and metformin. The only observed interaction between linagliptin and metformin was a reduction of body weight gain. No other additive toxicity caused by the combination of linagliptin and metformin was observed at AUC exposure levels up to 2 and 23 times human exposure, respectively.

An embryofetal development study in pregnant rats did not indicate a teratogenic effect attributed to the co-administration of linagliptin and metformin at AUC exposure levels up to 4 and 30 times human exposure, respectively.

Linagliptin

Liver, kidneys and gastrointestinal tract are the principal target organs of toxicity in mice and rats at repeat doses of linagliptin of more than 300 times the human exposure.

In rats, effects on reproductive organs, thyroid and the lymphoid organs were seen at more than 1500 times human exposure. Strong pseudo-allergic reactions were observed in dogs at medium doses, secondarily causing cardiovascular changes, which were considered dog-specific. Liver, kidneys, stomach, reproductive organs, thymus, spleen, and lymph nodes were target organs of toxicity in Cynomolgus monkeys at more than 450 times human exposure. At more than 100 times human exposure, irritation of the stomach was the major finding in these monkeys.

Linagliptin and its main metabolite did not show a genotoxic potential.

Oral 2 year carcinogenicity studies in rats and mice revealed no evidence of carcinogenicity in rats or male mice. A significantly higher incidence of malignant lymphomas only in female mice at the highest dose (> 200 times human exposure) is not considered relevant for humans (explanation: non-treatment related but due to highly variable background incidence). Based on these studies there is no concern for carcinogenicity in humans.

The NOAEL for fertility, early embryonic development and teratogenicity in rats was set at > 900 times the human exposure. The NOAEL for maternal-, embryo-fetal-, and offspring toxicity in rats was 49 times human exposure. No teratogenic effects were observed in rabbits at > 1,000 times human exposure. A NOAEL of 78 times human exposure was derived for embryo-fetal toxicity in rabbits, and for maternal toxicity the NOAEL was 2.1 times human exposure. Therefore, it is considered unlikely that linagliptin affects reproduction at therapeutic exposures in humans.

Metformin

Preclinical data for metformin reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential or reproductive toxicity.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core

Arginine

Copovidone

Magnesium stearate

Maize starch

Silica, colloidal anhydrous

Film coating

Hypromellose

Titanium dioxide (E171)

Talc

Propylene glycol

Trajenta Duo 2.5mg/500mg - Iron oxide, yellow (E172)

Trajenta Duo 2.5mg/850mg - Iron oxide, yellow (E172), Iron oxide, red (E172)

Trajenta Duo 2.5mg/1000mg - Iron oxide, red (E172)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

Please refer to the packaging for information on shelf-life.

6.4 Special precautions for storage

Do not store above 30°C.

Store in the original package in order to protect from moisture.

6.5 Nature and contents of container

Film-coated tablets of 2.5mg/500mg, 2.5mg/850mg and 2.5mg/1000mg

Cartons containing 60 film-coated tablets in a HDPE bottle.

6.6 Special precautions for disposal and other handling

Not applicable.

7. MANUFACTURER

Manufactured by:

Boehringer Ingelheim Promeco, S.A de C.V.

Calle del Maíz No. 49,

Col. Barrio Xaltocan, C.P. 16090

Xochimilco, Mexico City, Mexico.

For:

Boehringer Ingelheim International GmbH
Ingelheim am Rhein
Germany

8. DATE OF REVISION

30 April 2025