

# TANATRIL® TABLET

Imidapril Hydrochloride (5mg, 10mg)

## What is in this leaflet

1. What Tanatril is used for
2. How Tanatril works
3. Before you use Tanatril
4. How to use Tanatril
5. While you are using it
6. Side effects
7. Storage and Disposal of Tanatril
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What Tanatril is used for

Tanatril is used to treat hypertension which is a high blood pressure.

## How Tanatril works

Tanatril contains imidapril hydrochloride which is in a class of drugs called angiotensin-converting enzyme (ACE) inhibitors. ACE is an enzyme in a body that is necessary in the formation of a chemical called angiotensin II.

Angiotensin II causes the narrowing of arteries, thereby increasing blood pressure. ACE inhibitors prevent the formation of angiotensin II, hence widening the blood vessels and helping lower the blood pressure. Tanatril can be used on its own or alongside other medicines to help lower the blood pressure.

## Before you use Tanatril

### - When you must not use it

You must not take this product when you are:

- having or ever had an allergic reaction
- having a swelling below the skin surface, usually occurs around the eyes and lips.
- undergo hemodialysis.
- pregnant or may possibly be pregnant.

### - Before you start to use it

Before you start to take this product, make sure your doctor knows if you are having any heart problem, kidney problem, stroke and ischemia (restriction of blood supply to the brain).

### - Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

## How to use Tanatril

### - How much to use

The usual adult dosage for oral use is 5 mg to 10 mg of imidapril hydrochloride once a day. The dosage may be adjusted depending on the patient's age and symptom.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

### - When to use it

Use as directed by your doctor or pharmacist.

Usually taken once a day. Try to take this drug at the same times each day to avoid missing any doses.

### - How long to use it

Continue taking Tanatril for as long as your doctor recommends.

### - If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

## While you are using it

### - Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking Tanatril.

Tell your doctor immediately if you become pregnant while taking this medication because it may cause serious harm to your baby.

This product is not recommended for nursing mothers, if necessitated, breast feeding should be avoided.

### - Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give Tanatril to anyone else, even if they have the same symptoms or condition as you.

### - Things to be careful of

#### *Driving and using machines*

This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

**Side effects**

Like all medicines, Tanatril can cause side effects, although not everybody gets them.

Common side effects include:

- Cough
- Dizziness
- Hypotension (low blood pressure)
- Headache
- Rash

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835490, or visiting the website [nptra.gov.my](http://nptra.gov.my) [Consumers ➔ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

**Storage and Disposal of Tanatril****- Storage**

Keep all medicines out of the reach and sight of children.

Store at below 30°C.  
Avoid humidity after opening.

**- Disposal**

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

**Product Description****- What it looks like**

TANATRIL 5MG TABLET:  
White plain tablet with a secant line mark TA 135 on one side and 5 on the reverse side

TANATRIL 10MG TABLET:  
White plain tablet with a secant line mark TA 136 on one side and 10 on the reverse side

**- Ingredients**

- Active ingredient  
Imidapril hydrochloride

- Inactive ingredients  
Lactose 200 mesh, Macrogol 6000, Magnesium Stearate

**- MAL number:**

TANATRIL 5MG TABLET:  
MAL20034617AZ

TANATRIL 10MG TABLET:  
MAL20034618AZ

**Manufacturer**

PT Mitsubishi Tanabe Pharma Indonesia  
Jl. Rumah Sakit No. 104, RT 001 RW 005,  
Kel. Pakemitan, Kec. Cinambo, Kota  
Bandung, Jawa Barat, Indonesia

**Product Registration Holder**

Mitsubishi Tanabe Pharma  
Malaysia Sdn. Bhd.  
Suite 8-3, Level 8, Wisma  
UOA Damansara II, No. 6,  
Changkat Semantan,  
Damansara Heights, 50490  
Kuala Lumpur, Malaysia

**Date of revision**

10/12/2020

**Serial Number**

NPRA (R1/3) 23102020/181