

HEMAVEIN FILM-COATED TABLET 500MG

Diosmin/Hesperidin (450 mg/50 mg)

What is in this leaflet

1. What Hemavein Film-Coated Tablet 500mg is used for
2. How Hemavein Film-Coated Tablet 500mg works
3. Before you use Hemavein Film-Coated Tablet 500mg
4. How to use Hemavein Film-Coated Tablet 500mg
5. While you are using it
6. Side effects
7. Storage and Disposal of Hemavein Film-Coated Tablet 500mg
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Hemavein Film-Coated Tablet 500mg is used for

It is recommended for the treatment for venous circulation disorders (heavy legs, pain, early morning restless legs) and the treatment of symptoms related to hemorrhoids (acute hemorrhoidal attack).

How Hemavein Film-Coated Tablet 500mg works

This medicine strengthens veins by increasing venous tone, and protects small blood vessels.

Before you use Hemavein Film-Coated Tablet 500mg

- When you must not use it
Do not take Hemavein Film-Coated Tablet 500mg:
If you are allergic (hypersensitive) to diosmin, hesperidin, micronized purified flavonoid fraction or any of the other ingredients of this medicine (see Ingredients section).
Do not use in children below 18 years old unless advised by your doctor.

- Before you start to use it
Ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy and breast-feeding:

If you are pregnant, think you may be pregnant, planning to have a baby, or breast-feeding, ask your doctor or pharmacist for advice before taking this medicine.

- Taking other medicines

Please tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

How to use Hemavein Film-Coated Tablet 500mg

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

- How much to use
Venous circulation disorders
The recommended dose is 2 tablets daily; i.e. one at midday and one in the evening at mealtimes.

Acute hemorrhoidal attack

The recommended dose is 6 tablets daily (in 3 divided doses) for the first 4 days, then 4 tablets daily (in 2 divided doses) for the next 3 days.

- When to use it
Use as directed by your doctor or pharmacist.
- How long to use it
Continue taking Hemavein Film-Coated Tablet 500mg for as long as your doctor or pharmacist recommends.
- If you forget to use it
Do not take a double dose to make up for a forgotten dose.
If you have any further questions on the use of this product, ask your doctor or pharmacist.
- If you use too much (overdose)
If you take more Hemavein Film-Coated Tablet 500mg than you should, contact your doctor or pharmacist immediately.
Symptoms which may be experienced with overdose of Hemavein Film-Coated Tablet 500mg include diarrhoea, nausea, abdominal pain, itching and rash.

While you are using it

- Things you must do
Take your medicine exactly as your doctor or pharmacist has told you. Tell all the doctors, dentists and

pharmacists treating you that you are taking Hemavein Film-Coated Tablet 500mg.

- Things you must not do
This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- Things to be careful of
Be careful driving or operating machinery until you know how Hemavein Film-Coated Tablet 500mg affects you.

Side effects

Like all medicines, Hemavein Film-Coated Tablet 500mg can cause side effects, although not everybody gets them.

They may include:

Diarrhoea, dyspepsia (difficult digestion), nausea, vomiting, colitis (inflammation of the large intestine), dizziness, headache, discomfort, rash (sudden redness in patches), pruritus (skin reactions such as itching) and urticaria (allergic reaction in the form of small red spots), abdominal pain, isolated oedema (swelling of the face, lips and eyelids), Quincke's oedema (swelling of the face, lips, mouth, tongue or throat which could cause breathing difficulties).

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. By reporting side effects, you can help provide more information on the safety of this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website <http://npra.gov.my> [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Hemavein Film-Coated Tablet 500mg

- Storage

The expiry date of this pack is printed on the box. Do not use this pack after this date.

Keep all medicines out of reach of children.

Do not store above 30°C.

Date of revision

19 June 2025

Serial Number

NPRA (R2) 25/083

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.

Product Description

- What it looks like

Hemavein Film-Coated Tablet 500mg is brown yellow, oval-shaped film-coated tablet, biconvex and plain on both sides.

- Ingredients

- Active ingredients

Each film-coated tablet contains:

Micronized purified flavonoid fraction corresponding to:

Diosmin 450 mg

Flavonoids expressed as hesperidin 50 mg

- Inactive ingredients

Microcrystalline cellulose, sodium starch glycolate, magnesium stearate, gelatin, hypromellose, macrogol 6000, talc, titanium dioxide, sodium lauryl sulfate, yellow iron oxide, red ironoxide, glycerin.

- MAL number:

MAL25086043X

Manufacturer

Stellapharm J.V. Co., Ltd.- Branch 1
(A Joint Venture Company of Auxilto GmbH, Germany)

No. 40 Tu Do Avenue, Vietnam-Singapore Industrial Park, An Phu Ward, Thuan An City, Binh Duong Province, Vietnam.

Product Registration Holder

Aetos Pharma Sdn Bhd
66B, Tingkat 2, Jalan Cerdas, Taman Connaught, 56000 Kuala Lumpur, Malaysia.