

Nurofen for Children-Label [YODA PACK]

Child's Weight (age)	Quantity and method of administration	Frequency in 24 hours
20 - 29kg (6 - 9 years)	1 x 200mg/5ml (using the correct end of the spoon once)	3 times
30 - 40kg (10 - 12 years)	1 x 300mg/7.5ml (using the spoon twice (5ml & 2.5ml))	3 times

For short-term use only. If the child's symptoms persist for more than 3 days, consult your doctor. The doses should be given approximately every 6-8 hours (or with a minimum of 4 hours between each dose if required). Nurofen for Children is suitable for children from 6 years old weighting more than 20kg.

For oral administration only.
Shake the bottle before use.
Please refer to enclosed leaflet for further information.

DO NOT GIVE TO CHILDREN WHO:

- have stomach ulcers or other stomach disorders
- have kidney or heart problems • are suffering dehydration
- have allergic reactions to aspirin, ibuprofen or other anti-inflammatory medicines. If an allergic reaction develops, stop taking and see your doctor immediately.

Use only if cap seal is intact at first opening.

NUROFEN[®] for Children
Orange FLAVOUR Sugar Free
Suspension 200mg/5ml

100mL

6-12 years
Ibuprofen
CONCENTRATED FORMULA
FEVER AND PAIN RELIEF
200mg/5mL oral suspension

Preserved with Domiphen Bromide 0.01% w/v
Keep all medicines out of the reach of children.
Jauhi dari kanak-kanak

Manufactured by:
Reckitt Benckiser Healthcare (UK) Ltd.
Dansom Lane, Hull HU8 7DS, United Kingdom.
Do not store above 30°C.
Discard 6 months after opening.


Imported by/Product Registration Holder:
Reckitt Benckiser
Reckitt Benckiser (M) Sdn. Bhd. (175141-H)
Level 5, Menara UAC, No 12 Jalan PJU 7/5,
Mutiar Damansara, 47800 Petaling Jaya,
Selangor Darul Ehsan, Malaysia.
Reckitt Benckiser (S) Pte Ltd (197702628K)
Guthrie House, Singapore 268802.
MALXXXXXXXXX
SINXXXXXXXXX Controlled Medicine/Ubat Terkawal

Lot, Mfg & Exp Area
Varnish Free
Prints solid Nurofen Red

® Mfg. Exp.

BARCODE

RB122009



Latest Update: 22/06/2017