

VENOFLON FILM-COATED TABLET 500MG

Diosmin/Hesperidin (450mg/50mg)

What is in this leaflet

1. What Venoflon is used for
2. How Venoflon works
3. Before you use Venoflon
4. How to use Venoflon
5. While you are using it
6. Side effects
7. Storage and disposal of Venoflon
8. Product description
9. Manufacturer and Product
Registration Holder
10. Date of revision
11. Serial Number

What Venoflon is used for

It is recommended for the treatment for venous circulation disorders (heavy legs, pain, early morning restless legs) and the treatment of symptoms related to hemorrhoids (acute hemorrhoidal attack).

How Venoflon works

This medicine strengthens veins by increasing venous tone, and protects small blood vessels.

Before you use Venoflon

When you must not use it

Do not take Venoflon:

If you are allergic (hypersensitive) to diosmin, hesperidin, micronized purified flavonoid fraction or any of the other ingredients of this medicine (see Ingredients section).

Do not use in children below 18 years old unless advised by your doctor.

Before you start to use it

Ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy and breast-feeding:

If you are pregnant, think you may be pregnant, planning to have a baby, or breast-feeding, ask your doctor or pharmacist for advice before taking this medicine.

Taking other medicines

Please tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

How to use Venoflon

Always take this medicine exactly as your doctor or pharmacist has told you.

Check with your doctor or pharmacist if you are not sure.

How much to use & When to use it

Venous circulation disorders

The recommended dose is 2 tablets daily; i.e. one at midday and one in the evening at mealtimes.

Acute hemorrhoidal attack

The recommended dose is 6 tablets daily (in 3 divided doses) for the first 4 days, then 4 tablets daily (in 2 divided doses) for the next 3 days.

How long to use it

Depending on the medical condition for which you require Venoflon, your doctor may tell you to take it for a certain duration.

If you forget to use it

Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

If you use too much (overdose)

If you take more Venoflon than you should, contact your doctor or pharmacist immediately.

Symptoms which may be experienced with overdose of Venoflon include diarrhoea, nausea, abdominal pain, itching and rash.

While you are using it

Things you must do

Take your medicine exactly as your doctor or pharmacist has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking Venoflon.

Things you must not do

This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.

Side effects

Like all medicines, Venoflon can cause side effects, although not everybody gets them.

They may include:

Diarrhoea, dyspepsia (difficult digestion), nausea, vomiting, colitis (inflammation of the large intestine), dizziness, headache, discomfort, rash (sudden redness in patches), pruritus (skin reactions such as itching) and urticaria (allergic reaction in the form of small red spots), abdominal pain, isolated oedema (swelling of the face,

lips and eyelids), Quincke's oedema (swelling of the face, lips, mouth, tongue or throat which could cause breathing difficulties).

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. By reporting side effects, you can help provide more information on the safety of this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [*Consumers* → *Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)*].

Storage and disposal of Venoflon

Storage

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the box. The expiry date refers to the last day of that month.

Store in the original outer packaging. Store below 30°C.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.

Product Description

- What it looks like

Brown-yellow, oval-shaped film-coated tablet, biconvex and plain on both sides.

- Ingredients

Active ingredient

Each film-coated tablet contains:
Micronized purified flavonoid fraction corresponding to:

Diosmin 450 mg
Flavonoids expressed as hesperidin 50 mg

Inactive ingredients

Microcrystalline cellulose, sodium starch glycolate, magnesium stearate, gelatin, hypromellose, macrogol 6000, talc, titanium dioxide, sodium lauryl sulfate, yellow iron oxide, red iron oxide, glycerin.

VENOFLON FILM-COATED TABLET 500MG

Diosmin/Hesperidin (450mg/50mg)

MAL No.: MAL25036142XC

Manufacturer

**STELLAPHARM J.V. CO., LTD. -
BRANCH 1**

(A Joint Venture Company of Auxilto
GmbH, Germany)

No. 40 Tu Do Avenue, Vietnam-
Singapore Industrial Park, An Phu Ward,
Thuan An Town, Binh Duong Province,
Vietnam.

Product Registration Holder

AUXI THERAPEUTICS SDN. BHD.

E-9-02, Oasis Square Ara Damansara

No. 2, Jalan PJU 1A/7A,

47301 Petaling Jaya,

Selangor Darul Ehsan, Malaysia.

Date of revision: 10/09/25

Serial number: NPRA (R2) 25/007