건 명	아쿠스탑(ACUSTOP) 6매 호일(말레이시아)		
규 격 (가로x세로)	466x150mm	Q A	음성
		표시문안	6pt
색 상	6도 (별색2도 포함)	후 가 공	
업체제작일	2022.12.20.	작 업 처	(주)트리스톤

주요 변경 내용
1. 바코드 추가, 관리번호 변경 2. 제품등록권자 변경

C M Y K

칼선 PANTONE 185C

※인쇄진행시, 인쇄사양과 다를경우 녹십자로 즉시 연락주시기 바랍니다.

별적

PANTONE 1797C

466

10 12 191 10 10 10 10 191 12 10

150 130

10


10

GC Biopharma

Transdermal type anti-inflammatory & analgesics

ACUSTOP CATAPLASMA

(Flurbiprofen)



- *Rheumatism*
- *Neuralgia*
- *Arthritis*

Batch No. :

Manufacturing Date :

Expiry Date :

SIN 12090P
MAL 20013452AZ

8 806436 011434

6 Sheets(10cm X 14cm)

ZIP

Fasten the laminate zip to keep the pack closed.

cut here

cut here


Fasten the laminate zip to keep the pack closed.

ZIP

Transdermal type anti-inflammatory & analgesics

ACUSTOP CATAPLASMA

(Flurbiprofen)



[Composition] 1 sheet (10cm X 14cm) contains
Flurbiprofen 40mg
[Description]
Cataplasma which is spread with white pale yellowish adhesive mass on nonwoven fabric and has a peculiar odor.
[Indication]
For the treatment of the sign and symptoms of the followings: Arthritis deformans, periarthritis humero-scapularis, tendonitis, peritendinitis, humeral epicondylitis (tennis elbow), sore muscle, swelling, pain resulted from trauma.
[Dosage & administration]
Apply 1 sheet of Acustop Cataplasma to affected area twice a day.
[Contraindication]
1. Should be avoided in patients with exudative dermatosis, eczema, sores, infected skin lesions or broken skin.
2. Do not apply to patients of following: -
 i) Patients with history of hypersensitivity to this drug.
 ii) Patients with aspirin induced asthma (nonsteroidal anti-inflammatory drug induced asthma) or with history to it.
3. Carefully apply to the following patients: -
 i) Patients with bronchial asthma.

[Warning and Precautions]
General precautions:
1. Treatment of analgesics & anti-inflammatory drug is not a causal-therapy, but a symptomatic-therapy.
2. Consider the other therapy besides drug therapy in case of using this preparation for chronic disease (arthritis deformans, ect).

Observe carefully the status of patients and be cautions of the adverse reaction.
Precaution to children:
Safety and efficacy in children is not established.
Precaution in use:
1. Do not apply to eye, damaged skin and mucous membrane.
2. Do not apply to eczema and eruption region.
3. Wipe the sweat off affected area and then apply.

[Pregnancy and Lactation]
As safety in human pregnancy has not established, this preparation should not be administered to pregnant woman or woman of child-bearing potential unless therapeutic benefit is considered to exceed the possible risk.
[Drug Interaction]
In case of using for treatment of inflammation accompanied with infection, apply cautiously with administrating appropriate antibacterial or antifungal agent together and observe carefully because the symptoms of skin inflammation may not appear by this preparation.
[Side effects]
1. Skin: Occasionally eruption, redness, itchiness, fever and photosensitivity may occur. If these symptoms occur, discontinue application.

[Storage]
Store at below 30 °C, protect from moisture and out of direct sunlight.
Keep out of Reach of Children.
Jauhi dari kanak-kanak.
[Packing Size]
1 sheet (10cm X 14 cm) X 6 sheets.

CONTROLLED MEDICINE

UBAT TERKAWAL | 控制药 | دويت تركاول

Caution; Dental Use Only

MANUFACTURED BY

GC Biopharma

GC Biopharma Corp.
26, Mugeuk-ro 65beon-gil, Geumwang-eup, Eumseong-gun, Chungcheongbuk-do, Korea

PRODUCT REGISTRATION HOLDER

PHARMA NORD SDN BHD

29-07, Tingkat 29, Q-Sentral,
Jalan Stesen Sentral 2,
Kuala Lumpur Sentral,
50470 Kuala Lumpur, Malaysia

Foil-20-4005-9