

OPTICIN 1% w/w VISCOUS EYE DROPS

Fusidic Acid (10mg/g)

What is in this leaflet

1. What OPTICIN is used for
2. How OPTICIN works
3. Before you use OPTICIN
4. How to use OPTICIN
5. While you are using OPTICIN
6. Side effects
7. Storage and Disposal of OPTICIN
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial Number

What OPTICIN is used for

OPTICIN is indicated for bacterial eye infections caused by susceptible organisms in conjunctivitis (inflammation of the white of the eye, or called as "red eye"), blepharitis (inflammation of the eyelid), sty (a lump on the edge of the upper or lower eyelid), keratitis (inflammation of the cornea, which is the clear area in the centre of the eye), dacryocystitis (inflammation of the tear sac) and in connection with removal of foreign bodies.

How OPTICIN works

OPTICIN works by killing germs (bacteria) that cause infections.

Before you use OPTICIN

-When you must not use it

Do not use OPTICIN

- If you are allergic (hypersensitive) to fusidic acid, or any of the other ingredients in your medicine. You can find a list of these ingredients in section of product description of this leaflet.

-Before you start use it

Pregnancy and breast-feeding

Please ask your doctor or pharmacist for advice before using OPTICIN:

- If you are pregnant, or think you are pregnant.

- If you are breast-feeding.

-Taking other medicines

Please tell your doctor or pharmacist if you are taking, or have recently taken any other medicines. This includes any medicines which you have bought without a prescription.

How to use OPTICIN

-How much to use

Always use OPTICIN exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

This medicine is only for use in your eyes. Do not swallow the drops. Do not put the drops inside your body or in your ears.

Always wash your hands before using OPTICIN. Do not use if the cap seal on the tube is broken. This diagram will help show how to use your medicine. It may be useful to look in a mirror.



Only take the cap off the tube when you are ready to use the medicine. It is important the tip of the tube does not touch your eye.

Tilt your head back. Pull your lower eyelid down gently. Hold the tube over your eye and look up. Squeeze one drop into your lower eyelid. The drop will come out as a thick drop. It will become more liquid in your eye, and you should still see clearly.

After using your medicine, you

may see a white powder around your eye. This can happen when the drops dry out. It is quite normal and is nothing to worry about. You can wipe the powder off with cotton wool.

If the medicine is for a child, you can put the drops in their eye while they are asleep or lying down if it is easier.

Your doctor will tell you how many drops to use, or to give to your child. To remind you to use your medicine it may help to use it when you do another regular action, such as brushing your teeth.

Adults, the elderly and children: The usual dose is one drop twice each day, in the morning and in the evening. Treatment should be continued for at least two days after the eye appears normal.

-When to use it

Use as directed by your doctor or pharmacist.

-How long to use it

It is very important to use all the medicine that your doctor has told you to use. You must use your medicine for at least 2 days after you feel better, and your eye looks normal. You must do this because otherwise your eye infection may come back.

-If you forget to use it

If you forget to use your medicine, use it as soon as you remember. Then next use your medicine at the usual time. If you have any further questions about taking this medicine, please ask your doctor or pharmacist.

-If you use too much (overdose)

If you accidentally apply too many drops in the eye(s) or a small quantity of the drops are accidentally swallowed, it is unlikely to cause any harm. If a

OPTICIN 1% w/w VISCOUS EYE DROPS

Fusidic Acid (10mg/g)

larger quantity of the drops are accidentally swallowed or you are worried contact your doctor or pharmacist for advice.

While you are using OPTICIN

-Things you must do

Use your medicine exactly as your doctor has told you.

Tell your doctor if you become pregnant while using this medicine.

-Things you must not do

Do not wear any type of contact lenses whilst using this medicine.

-Things to be careful of

OPTICIN contains:

Benzalkonium chloride. This is a preservative. It may cause eye irritation.

Please ask your doctor if you are worried about any of the ingredients in this medicine.

Driving and using machines

Following application of your medicine, a brief blurring of vision may occur. Do not drive or use machinery until your vision is clear. Check with your doctor if you feel any side effect that may stop you from driving or using machines.

Side effects

Like all medicines, OPTICIN can cause side effects, although not everybody gets them.

Transient stinging may occur. OPTICIN has a very low allergenic potential.

If any of the side effects become serious or you notice any side effects not listed in this leaflet, tell your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the

website nprr.gov.my
[Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of OPTICIN

-Storage

Keep OPTICIN out of the reach and sight of children

Store below 30°C.

Once the tube is opened, this product should be used in 15 days.

Use it in accordance with the expiry date.

Do not use OPTICIN after the expiry date which is stated on the pack.

Do not use OPTICIN if you notice any defects in the product and/or its pack.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

-What it looks like

OPTICIN white or off-white homogenous viscous solution in laminate (polyfoil) tubes with high density polyethylene tip and cap.

-Ingredients

Active ingredient: Each gram contains 10.18 mg fusidic acid hemihydrate equal to 10 mg fusidic acid as active substance.

Inactive ingredients:

Benzalkonium chloride, mannitol, disodium edetate, carbomer (974 P), sodium hydroxide, purified water.

- **MAL No.:**

MAL xxxx

Manufacturer

DEVA Holding Anonim
Sirketi
Çerkezköy Organize Sanayi
Bölgesi
Karaağaç Mah. Atatürk Cad.,
No: 32
Kapaklı -
TEKİRDAĞ/TURKEY

Product Registration Holder

InstaMed Asia Sdn Bhd.
108 Block E,
Phileo Damansara 1,
No. 9, Jalan 16/11,
46350 Petaling Jaya,
Selangor, Malaysia.

Date of Revision

18.12.2024

Serial Number

NPRA (R2) 24/102