



HurriCaine Cherry Topical Anesthetic Spray 20%

Clear tan liquid with distinctive cherry odor.

Content:

Benzocaine 20%

Indication:

Topical anesthetic used for temporary relief of occasional minor irritation and pain associated with, minor dental procedures, canker sores, sore mouth and throat, minor injury of the mouth and gums, and minor irritation of the mouth and gums.

Pharmacology:

Benzocaine is an analgesic (dulls pain), anesthetic (numbs nerves), antipyretic (calms itching).

Contraindications:

Patients with known hypersensitivity to benzocaine. Application to severely traumatized mucosa or to areas of sepsis.

Side effects/Adverse reactions:

Contact dermatitis and/or hypersensitivity to benzocaine can cause burning, stinging, pruritus, tenderness, erythema, rash, urticaria and edema. Rarely, benzocaine may induce methemoglobinemia causing respiratory distress and cyanosis.

Precautions:

Localized allergic reactions may occur in patients sensitive to benzocaine or after prolonged or repeated use. Cross-sensitivity may occur with other local anesthetic of the ester type and systemic medications such as sulphonamides, thiazides and para-amino dyes. Methemoglobinemia may rarely occur, particularly in infant and children, following application of large amounts e.g. in pharyngeal anesthesia. Topical benzocaine should be used in children below 2 years under the direction of dentist or physician. It should not be used for prolonged periods.

Use in pregnancy and lactation:

Animal reproduction studies have not been conducted with benzocaine. Topical spray should be given to a pregnant woman only if clearly indicated.

It is not known whether benzocaine is excreted in human milk. Caution should be exercised when spray is administered to a nursing woman.

Drug Interactions:

None known.

Dosage and administration:

Firmly insert extension tube into spray can valve and ensure it fits securely, into the actuator opening position the extension tube 1-2 inches from the affected area. Spray 1/2 second (press and immediately release actuator), repeat once if necessary, anesthesia is accomplished in approximately 20 seconds.

Treatment of overdose:

Accidental swallowing of large quantities causes numbness, nausea, vomiting, irritation to eyes.

In case of eye and skin contact, flush with large quantities of water. In case of accidental swallowing, to get medical attention immediately. Treatment is primarily supportive. It includes monitoring for development of methemoglobinemia.

Storage conditions: Store below 30°C. Protect from light. Keep out of reach of children.

Presentation : 60gm/bottle

Shelflife : 3 years.

Manufactured by : KIK Custom Products
1, West Hageler Lane, Danville, IL 61832 USA.

Distributed by : TK Dental Trading
e-Gate 1-03-10, 1, Lebu Tunku Kudin 2, Gelugor, 11700 Penang.