



ZINNAT
Granules
for 50ml
125mg
SUSPENSION

SANDOZ

- Unconstituted Zinnat Suspension should be stored below 30°C
- Reconstituted Zinnat Suspension must be refrigerated between 2 and 8°C
- **USE ONLY AS DIRECTED BY YOUR DOCTOR**

YOU SHOULD NOT INCREASE YOUR DOSAGE UNLESS YOUR DOCTOR TELLS YOU.

- **TO BE TAKEN WITH FOOD**
- Keep out of the reach of children
- Jauhkan daripada kanak-kanak
- Read enclosed leaflet before use

Manufactured by:
Sandoz GmbH, Biochemiestraße 10,
6250 Kundl, Austria.

Product Registration Holder:
Sandoz Products Malaysia Sdn. Bhd.
3A-2 & 3A-3, Level 3A,
Menara KEN TTDI,
No. 37, Jalan Burhanuddin Helmi,
Taman Tun Dr Ismail,
60000 Kuala Lumpur, Malaysia

MAL19921371ARZ
BRU08070090P



5 033439 008953

HOW TO PREPARE YOUR MEDICINE

1 SHAKE BOTTLE WELL TO LOOSEN GRANULES	2 FILL THE MEASURING CUP TO THE LINE WITH WATER
3 ADD WATER TO BOTTLE IN ONE GO & REPLACE CAP	4 SHAKE BOTTLE WELL TO MIX MEDICINE PROPERLY
5 TAKE MEDICINE AS PRESCRIBED*	KEEP TAKING THE MEDICINE UNTIL THE COURSE IS COMPLETED. DON'T STOP JUST BECAUSE YOU FEEL BETTER. THE MEDICINE MAY NOT HAVE KILLED ALL THE GERMS AND YOU MAY START TO FEEL UNWELL AGAIN.

*THE DOSE MAY BE ADDED TO CHILDREN'S COLD FRUIT JUICES OR MILK IMMEDIATELY BEFORE TAKING

- When prepared as directed, each 5ml contains 125mg cefuroxime as cefuroxime axetil

Controlled Medicine/Ubat Terkawal
Aspartame: Unsuitable for phenylketonurics



SANDOZ
SUSPENSION
125mg
Oral use
Granules for 50ml

Contains 125mg
cefuroxime as
cefuroxime axetil

SANDOZ
SUSPENSION
125mg
Oral use
Granules for 50ml

Contains 125mg
cefuroxime as
cefuroxime axetil



LOT:
MFD:
EXP:

