

Zofran™

Serotonin (5HT₃) antagonist

DESCRIPTION AND COMPOSITION**Pharmaceutical form****Tablets:**

Ondansetron tablets 4 mg: Yellow, oval, film-coated tablet engraved with 'GX ET3'. Each tablet contains ondansetron 4 mg as hydrochloride dihydrate.

Ondansetron tablets 8 mg: Yellow, oval, film-coated tablet engraved with 'GX ET5'. Each tablet contains ondansetron 8 mg as hydrochloride dihydrate.

Active substance

Ondansetron/Ondansetron hydrochloride dihydrate

Excipients

Lactose

Microcrystalline cellulose

Pregelatinized maize starch

Magnesium stearate

Methyl hydroxypropyl cellulose

Titanium dioxide (E171)

Iron oxide (E172)

INDICATIONS**Adults**

ZOFRAN oral formulations are indicated for the management of nausea and vomiting induced by cytotoxic chemotherapy and radiotherapy.

ZOFRAN is also indicated for the prevention of post-operative nausea and vomiting.

Paediatric Population***Oral formulation:***

ZOFRAN is indicated for the management of nausea and vomiting induced by cytotoxic chemotherapy.

No studies have been conducted on the use of orally administered ondansetron in the prevention or treatment of post-operative nausea and vomiting.

DOSAGE REGIMEN AND ADMINISTRATION

Dosage regimen

CHEMOTHERAPY AND RADIOTHERAPY INDUCED NAUSEA AND VOMITING (CINV and RINV)

The emetogenic potential of cancer treatment varies according to the doses and combinations of chemotherapy and radiotherapy regimens used. The selection of dose regimen should be determined by the severity of the emetogenic challenge.

CINV and RINV in Adults

The recommended oral dose is 8 mg taken 1 to 2 hours before chemotherapy or radiation treatment, followed by 8 mg orally every 12 hours for a maximum of 5 days.

For highly emetogenic chemotherapy a single oral dose of up to 24 mg ondansetron taken together with 12 mg oral dexamethasone sodium phosphate, 1 to 2 hours before chemotherapy, may be used. After the first 24 hours, oral treatment with ondansetron may be continued for up to 5 days after a course of treatment. The recommended oral dose is 8 mg to be taken twice daily.

CINV in Children and Adolescents (aged 3 years to 17 years)

The dose for CINV can be calculated based on body surface area (BSA) or weight. In paediatric clinical studies, ondansetron was given by IV infusion diluted in 25 to 50 mL of saline or other compatible infusion fluid and infused over not less than 15 minutes.

Dosing by BSA

Ondansetron should be administered immediately before chemotherapy as a single IV dose of 5 mg/m². The IV dose must not exceed 8 mg. Oral dosing can commence 12 hours later and may be continued for up to 5 days (Table 1). Adult doses must not be exceeded.

Table 1 BSA-based dosing for CINV (aged 3 years to 17 years)

BSA	Day 1	Days 2 - 6
≥ 0.6 m ² to ≤ 1.2 m ²	5 mg/m ² IV plus 4 mg tablet after 12 hours	4 mg tablet every 12 hours
> 1.2 m ²	5 mg/m ² IV or 8 mg IV plus 8 mg tablet after 12 hours	8 mg tablet every 12 hours

Dosing by body weight

Ondansetron should be administered immediately before chemotherapy as a single IV dose of 0.15 mg/kg. The IV dose must not exceed 8 mg. On Day 1, two further IV doses may be given in 4-hourly intervals. Oral dosing can commence 12 hours later and may be continued for up to 5 days (Table 2). Adult doses must not be exceeded.

Table 2 Weight-based dosing for CINV (aged 3 years to 17 years)

Body Weight	Day 1	Days 2 - 6
> 10 kg	Up to 3 doses of 0.15 mg/kg IV every 4 hours	4 mg tablet every 12 hours

There are no dosing recommendations for children with BSA <0.6 m², body weight ≤10 kg or who are unable to swallow tablets.

CINV and RINV in Elderly

No alteration of oral dose or frequency of administration is required.

POST-OPERATIVE NAUSEA AND VOMITING (PONV)

PONV in Adults

For prevention of post-operative nausea and vomiting, the recommended oral dose is 16 mg given 1 hour prior to anaesthesia.

For treatment of established post-operative nausea and vomiting ondansetron administration by injection is recommended.

PONV in Children and Adolescents (aged 3 years to 17 years)

No studies have been conducted on the use of orally administered ondansetron in the prevention or treatment of post-operative nausea and vomiting; slow IV injection (not less than 30 seconds) is recommended for this purpose.

PONV in Elderly

There is limited experience in the use of ondansetron in the prevention and treatment of post-operative nausea and vomiting in the elderly, however ondansetron is well tolerated in patients over 65 years receiving chemotherapy.

Special populations

Renal impairment

No alteration of daily dosage or frequency of dosing, or route of administration are required.

Hepatic impairment

Clearance of ondansetron is significantly reduced and serum half-life significantly prolonged in subjects with moderate or severe impairment of hepatic function. In such patients, a total daily dose of 8 mg IV or oral should not be exceeded.

Patients with Poor Sparteine/Debrisoquine Metabolism

The elimination half-life of ondansetron is not altered in subjects classified as poor metabolisers of sparteine and debrisoquine. Consequently in such patients repeat dosing will give drug exposure levels no different from those of the general population. No alteration of daily dosage or frequency of dosing is required.

Method of administration

The tablets should be swallowed whole with liquid.

CONTRAINDICATIONS

Based on reports of profound hypotension and loss of consciousness when ondansetron was administered with apomorphine hydrochloride, concomitant use with apomorphine is contraindicated (see section INTERACTIONS).

Hypersensitivity to any component of the preparation (see section WARNINGS AND PRECAUTIONS and section ADVERSE DRUG REACTIONS).

WARNINGS AND PRECAUTIONS

Hypersensitivity reactions have been reported in patients who have exhibited hypersensitivity to other selective 5HT₃ receptor antagonists.

Ondansetron prolongs the QT interval in a dose-dependent manner (see section CLINICAL PHARMACOLOGY). In addition, post-marketing cases of Torsade de Pointes have been reported in patients using ondansetron. Avoid ondansetron in patients with congenital long QT syndrome. Ondansetron should be administered with caution to patients who have or may develop prolongation of QTc, including patients with electrolyte abnormalities, congestive heart failure, bradyarrhythmias or patients taking other medicinal products that lead to QT prolongation or electrolyte abnormalities.

Myocardial ischemia has been reported in patients treated with ondansetron. In some cases, predominantly during intravenous administration, the symptoms appeared immediately after administration but recovered with prompt treatment. Therefore, caution should be exercised during and after administration of ondansetron.

Hypokalemia and hypomagnesemia should be corrected prior to ondansetron administration.

Serotonin syndrome has been described following the concomitant use of ondansetron and other serotonergic drugs (see section INTERACTIONS). If concomitant treatment with ondansetron and other serotonergic drugs is clinically warranted, appropriate observation of the patient is advised.

As ondansetron is known to increase large bowel transit time, patients with signs of subacute intestinal obstruction should be monitored following administration.

ADVERSE DRUG REACTIONS

Summary of the safety profile

Adverse events are listed below by system organ class and frequency. Frequencies are defined as: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1000$ to $< 1/100$); rare ($\geq$

1/10,000 to <1/1000); and very rare (<1/10,000), including isolated reports. Very common, common and uncommon events were generally determined from clinical trial data. The incidence in placebo was taken into account. Rare and very rare events were generally determined from post-marketing spontaneous data. The following frequencies are estimated at the standard recommended doses of ondansetron. The adverse event profiles in children and adolescents were comparable to that seen in adults.

Table 3 Adverse Drug Reactions

Immune system disorders	
Rare:	Immediate hypersensitivity reactions sometimes severe, including anaphylaxis.
Nervous system disorders	
Very common:	Headache
Uncommon:	Seizures, movement disorders (including extrapyramidal reactions such as dystonic reactions, oculogyric crisis and dyskinesia) have been observed without definitive evidence of persistent clinical sequelae.
Rare:	Dizziness predominantly during rapid IV administration
Eye disorders	
Rare:	Transient visual disturbances (e.g. blurred vision) predominantly during IV administration.
Very rare:	Transient blindness predominantly during IV administration.
The majority of the blindness cases reported resolved within 20 minutes. Most patients had received chemotherapeutic agents, which included cisplatin. Some cases of transient blindness were reported as cortical in origin.	
Cardiac disorders	
Uncommon:	Arrhythmias, chest pain with or without ST segment depression, bradycardia.
Rare:	QTc prolongation (including Torsade de Pointes)
Vascular disorders	
Common:	Sensation of warmth or flushing
Uncommon:	Hypotension
Respiratory, thoracic and mediastinal disorders	
Uncommon:	Hiccups
Gastrointestinal disorders	
Common:	Constipation
Hepatobiliary disorders	
Uncommon:	Asymptomatic increases in liver function tests [#]
[#] These events were observed commonly in patients receiving chemotherapy with cisplatin.	
Skin and subcutaneous tissue disorders	
Very rare:	Toxic skin eruption, including toxic epidermal necrolysis

Adverse drug reactions from spontaneous reports and literature cases (frequency not known)

The following adverse drug reactions have been derived from post-marketing experience with Zofran via spontaneous case reports and literature cases. Because these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency which is therefore categorized as not known. Adverse drug reactions are listed according to system organ classes in MedDRA.

Table 4 Adverse drug reactions from spontaneous reports and literature (frequency not known)

Cardiac disorders
Myocardial ischemia

INTERACTIONS

There is no evidence that ondansetron either induces or inhibits the metabolism of other drugs commonly co-administered with it. Specific studies have shown that there are no pharmacokinetic interactions when ondansetron is administered with alcohol, temazepam, furosemide, tramadol or propofol.

Ondansetron is metabolised by multiple hepatic cytochrome P450 enzymes: CYP3A4, CYP2D6 and CYP1A2. Due to the multiplicity of metabolic enzymes capable of metabolising ondansetron, enzyme inhibition or reduced activity of one enzyme (e.g. CYP2D6 genetic deficiency) is normally compensated by other enzymes and should result in little or no significant change in overall ondansetron clearance or dose requirement.

Caution should be exercised when ondansetron is co-administered with drugs that prolong the QT interval and/or cause electrolyte abnormalities (see section WARNINGS AND PRECAUTIONS).

Apomorphine

Based on reports of profound hypotension and loss of consciousness when ondansetron was administered with apomorphine hydrochloride, concomitant use with apomorphine is contraindicated.

Phenytoin, Carbamazepine and Rifampicin

In patients treated with potent inducers of CYP3A4 (i.e. phenytoin, carbamazepine, and rifampicin), the oral clearance of ondansetron was increased and ondansetron blood concentrations were decreased.

Serotonergic Drugs (e.g., SSRIs and SNRIs)

Serotonin syndrome (including altered mental status, autonomic instability and neuromuscular abnormalities) has been described following the concomitant use of ondansetron and other serotonergic drugs, including selective serotonin reuptake inhibitors (SSRIs) and serotonin noradrenaline reuptake inhibitors (SNRIs) (see section WARNINGS AND PRECAUTIONS).

Tramadol

Data from small studies indicate that ondansetron may reduce the analgesic effect of tramadol

PREGNANCY, LACTATION, FEMALES AND MALES OF REPRODUCTIVE POTENTIAL

PREGNANCY

Risk Summary

In human epidemiological studies, an increase in orofacial clefts was observed in infants of women administered ondansetron during the first trimester of pregnancy. Regarding cardiac malformations, the epidemiological studies showed conflicting results (see Human data). Reproductive studies in rats and rabbits did not show evidence of harm to the fetus (see Animal data)

The use of ondansetron in pregnancy is not recommended.

Human Data

Three epidemiological studies in the US assessed the risk of specific congenital anomalies, including orofacial clefts and cardiac malformations in offspring born to mothers exposed to ondansetron during the first trimester of pregnancy.

One cohort study with 88,467 pregnancies exposed to ondansetron showed an increased risk of oral clefts (3 additional cases per 10,000 women treated, adjusted relative risk (RR), 1.24 (95% CI 1.03-1.48)) without an apparent increase in risk of cardiac malformations. A separately published subgroup analysis of 23,877 pregnancies exposed to intravenous ondansetron did not find an increased risk of either oral clefts or cardiac malformations.

One case-control study using population-based birth defect registries with 23,200 cases across two datasets showed an increased risk of cleft palate in one dataset and no increased risk in the other dataset. There was no increased risk of cardiac malformations in this study.

The second cohort study with 3,733 pregnancies exposed to ondansetron found an increased risk of ventricular septal defect, adjusted RR 1.7 (95%CI 1.0-2.9), but no statistically significant increase in risk of cardiac malformations.

Animal Data

In embryo-fetal development studies in rats and rabbits, pregnant animals received oral doses of ondansetron up to 15 mg/kg/day and 30 mg/kg/day, respectively, during the period of organogenesis. With the exception of a slight decrease in maternal body weight gain in the rabbits, there were no significant effects of ondansetron on the maternal animals or the development of the offspring. At doses of 15 mg/kg/day in rats and 30 mg/kg/day in rabbits, the maternal dose was approximately 6 and 24 times the maximum recommended human oral dose of 24 mg/day, respectively, based on body surface area. In a pre- and postnatal developmental toxicity study, pregnant rats received oral doses of ondansetron up to 15 mg/kg/day from Day 17 of pregnancy to litter Day 21. With the exception of a slight reduction in maternal body weight gain, there were no effects upon the pregnant rats and the pre- and postnatal development of their offspring, including reproductive performance of the mated F1 generation. At a dose of 15 mg/kg/day in rats, the maternal dose was approximately 6 times the maximum recommended human oral dose of 24 mg/day based on BSA.

LACTATION

Risk Summary

It is not known whether Zofran is transferred into human milk. There are no data on the effects of Zofran on the breastfed child or the effects of Zofran on milk production. However, it has been demonstrated that ondansetron passes into the milk of lactating animals (rats). It is therefore recommended that mothers receiving ondansetron should not breast-feed their babies.

FEMALES AND MALES OF REPRODUCTIVE POTENTIAL

Pregnancy testing

Pregnancy status should be verified for females of reproductive potential prior to starting the treatment with Zofran.

Contraception

Females of reproductive potential should be advised that it is possible that Zofran can cause harm to the developing fetus. Sexually active females of reproductive potential are recommended to use effective contraception (methods that result in less than 1 % pregnancy rates) when using Zofran during the treatment and for two days after stopping treatment with Zofran.

Infertility

There is no effect of Zofran on fertility.

OVERDOSAGE

There is limited experience of ondansetron overdose. In the majority of cases, symptoms were similar to those already reported in patients receiving recommended doses (see section ADVERSE DRUG REACTIONS).

Ondansetron prolongs QT interval in a dose-dependent fashion. ECG monitoring is recommended in cases of overdose.

Cases consistent with serotonin syndrome have been reported in young children following oral overdose.

Treatment

There is no specific antidote for ondansetron, therefore in cases of suspected overdose, symptomatic and supportive therapy should be given as appropriate.

The use of ipecacuanha to treat overdose with ondansetron is not recommended as patients are unlikely to respond due to the anti-emetic action of ondansetron itself.

CLINICAL PHARMACOLOGY

Mechanism of action (MOA)

Ondansetron is a potent, highly selective 5HT₃ receptor antagonist. Its precise mode of action in the control of nausea and vomiting is not known.

Chemotherapeutic agents and radiotherapy may cause release of 5HT in the small intestine initiating a vomiting reflex by activating vagal afferents via 5HT₃ receptors. Ondansetron blocks the initiation of this reflex.

Activation of vagal afferents may also cause a release of 5HT in the area postrema, located on the floor of the fourth ventricle, and this may also promote emesis through a central mechanism. Thus, the effect of ondansetron in the management of the nausea and vomiting induced by cytotoxic chemotherapy and radiotherapy is probably due to antagonism of 5HT₃ receptors on neurons located both in the peripheral and central nervous system.

The mechanisms of action in post-operative nausea and vomiting are not known but there may be common pathways with cytotoxic induced nausea and vomiting.

Pharmacodynamics (PD)

Ondansetron does not alter plasma prolactin concentrations.

QT Prolongation

The effect of ondansetron on the QTc interval was evaluated in a double blind, randomized, placebo and positive (moxifloxacin) controlled, crossover study in 58 healthy adult men and women. Ondansetron doses included 8 mg and 32 mg infused intravenously over 15 minutes. At the highest tested dose of 32 mg, the maximum mean (upper limit of 90% CI) difference in QTcF from placebo after baseline-correction was 19.6 (21.5) msec. At the lower tested dose of 8 mg, the maximum mean (upper limit of 90% CI) difference in QTcF from placebo after baseline-correction was 5.8 (7.8) msec. In this study, there were no QTcF measurements greater than 480 msec and no QTcF prolongation was greater than 60 msec.

Pharmacokinetics (PK)

The pharmacokinetic properties of ondansetron are unchanged on repeat dosing.

Absorption

Following oral administration, ondansetron is passively and completely absorbed from the gastrointestinal tract and undergoes first pass metabolism. Peak plasma concentrations are attained approximately 1.5 hours after dosing. For doses above 8 mg the increase in ondansetron systemic exposure with dose is greater than proportional; this may reflect some reduction in first pass metabolism at higher oral doses.

Mean bioavailability in healthy male subjects, following the administration of a single 8 mg tablet, is approximately 55 to 60%. Bioavailability is slightly enhanced by the presence of food but unaffected by antacids.

Distribution

Ondansetron is not highly protein bound (70 to 76%).

The disposition of ondansetron following oral, IM or IV dosing in adults is similar with a steady state volume of distribution of about 140 L.

Metabolism

Ondansetron is cleared from the systemic circulation predominantly by hepatic metabolism through multiple enzymatic pathways. The absence of the enzyme CYP2D6 (the debrisoquine polymorphism) has no effect on ondansetron's pharmacokinetics.

Elimination

Ondansetron is cleared from the systemic circulation predominantly by hepatic metabolism. Less than 5% of the absorbed dose is excreted unchanged in the urine.

The disposition of ondansetron following oral, IM or IV dosing is similar with a terminal elimination half-life of about 3 hours.

Special Patient Populations

Gender

Gender differences were shown in the disposition of ondansetron, with females having a greater rate and extent of absorption following an oral dose and reduced systemic clearance and volume of distribution (adjusted for weight).

Geriatric population (65 years of age or older)

Early Phase I studies in healthy elderly volunteers showed a slight age-related decrease in clearance, and an increase in half-life of ondansetron. However, wide inter-subject variability resulted in considerable overlap in pharmacokinetic parameters between young (< 65 years of age) and elderly subjects ($\geq$ 65 years of age) and there were no overall differences in safety or efficacy observed between young and elderly cancer patients enrolled in CINV clinical trials to support a different dosing recommendation for the elderly.

Based on more recent ondansetron plasma concentrations and exposure-response modelling, a greater effect on QTcF is predicted in patients $\geq$ 75 years of age compared to young adults.

Pediatric population (aged 3 years to 17 years)

The half-life is 2.9 hours for patients 3 to 12 year age range.

In paediatric patients aged 3 to 12 years undergoing elective surgery with general anaesthesia, the absolute values for both the clearance and volume of distribution of ondansetron were reduced in comparison to values with adult patients. Both parameters increased in a linear fashion with weight and by 12 years of age, the values were approaching those of young adults. When clearance and volume of distribution values were normalised by body weight, the values for these parameters were similar between the different age group populations. Use of weight-based dosing compensates for age-related changes and is effective in normalising systemic exposure in paediatric patients.

Population pharmacokinetic analysis was performed on 428 subjects (cancer patients, surgery patients and healthy volunteers) aged 1 month to 44 years following IV administration of ondansetron. Based on this analysis, systemic exposure (AUC) of ondansetron following oral or IV dosing in children and adolescents was comparable to adults, with the exception of infants aged 1 to 4 months. Volume of distribution was related to age and was lower in adults than in infants and children. Clearance was related to weight but not to age with the exception of infants aged 1 to 4 months.

Hepatic impairment

In patients with severe hepatic impairment, ondansetron's systemic clearance is markedly reduced with prolonged elimination half-lives (15 to 32 hours) and an oral bioavailability approaching 100% due to reduced pre-systemic metabolism.

CLINICAL STUDIES

Children and adolescents (aged 1 month to 17 years)

CINV

The efficacy of ondansetron in the control of emesis and nausea induced by cancer chemotherapy was assessed in a double-blind randomized trial in 415 patients aged 1 to 18 years (S3AB3006). On the days of chemotherapy, patients received either ondansetron 5 mg/m² IV and ondansetron 4 mg orally after 8 to 12 hours or ondansetron 0.45 mg/kg IV and placebo orally after 8 to 12 hours. Post-chemotherapy both groups received 4 mg ondansetron syrup twice daily for 3 days. Complete control of emesis on worst day of chemotherapy was 49% (5 mg/m² IV and ondansetron 4 mg orally) and 41% (0.45 mg/kg IV and placebo orally). Post-chemotherapy both groups received 4 mg ondansetron syrup twice daily for 3 days. There was no difference in the overall incidence or nature of adverse events between the two treatment groups.

A double-blind randomized placebo-controlled trial (S3AB4003) in 438 patients aged 1 to 17 years demonstrated complete control of emesis on worst day of chemotherapy in:

- 73% of patients when ondansetron was administered intravenously at a dose of 5 mg/m² IV together with 2 to 4 mg dexamethasone orally
- 71% of patients when ondansetron was administered as syrup at a dose of 8 mg together with 2 to 4 mg dexamethasone orally on the days of chemotherapy.

Post-chemotherapy both groups received 4 mg ondansetron syrup twice daily for 2 days. There was no difference in the incidence or nature of adverse events between the two treatment groups.

The efficacy of ondansetron in 75 children aged 6 to 48 months was investigated in an open-label, non-comparative, single-arm study (S3A40320). All children received three 0.15 mg/kg doses of IV ondansetron, administered 30 minutes before the start of chemotherapy and then at 4 and 8 hours after the first dose. Complete control of emesis was achieved in 56% of patients.

Another open-label, non-comparative, single-arm study (S3A239) investigated the efficacy of one IV dose of 0.15 mg/kg ondansetron followed by two oral ondansetron doses of 4 mg for children aged < 12 years and 8 mg for children aged ≥ 12 years (total no. of children n= 28). Complete control of emesis was achieved in 42% of patients.

PONV

The efficacy of a single dose of ondansetron in the prevention of post-operative nausea and vomiting was investigated in a randomized, double-blind, placebo-controlled study in 670 children aged 1 to 24 months (post-conceptual age ≥ 44 weeks, weight ≥ 3 kg). Included subjects were scheduled to undergo elective surgery under general anesthesia and had an ASA status $\leq$ III. A single dose of ondansetron 0.1 mg/kg was administered within five minutes following induction of anesthesia. The proportion of subjects who experienced at least one emetic episode during the 24-hour assessment period (ITT) was greater for patients on placebo than those receiving ondansetron (28% vs. 11%, $p < 0.0001$). Four double-blind, placebo-controlled studies have been performed in 1469 male and female patients (2 to 12 years of age) undergoing general anesthesia. Patients were randomized to either single IV doses of ondansetron (0.1 mg/kg for pediatric patients weighing 40 kg or less, 4 mg for pediatric patients weighing more than 40 kg; number of patients = 735) or placebo (number of patients = 734). Study drug was administered over at least 30 seconds, immediately prior to or following anesthesia induction. Ondansetron was significantly more effective than placebo in preventing nausea and vomiting. The results of these studies are summarized in Table 5.

Table 5 Prevention and treatment of PONV in Pediatric Patients – Treatment response over 24 hours

Study	Endpoint	Ondansetron (%)	Placebo (%)	p value
S3A380	CR	68	39	≤ 0.001
S3GT09	CR	61	35	≤ 0.001
S3A381	CR	53	17	≤ 0.001
S3GT11	no nausea	64	51	0.004
S3GT11	no emesis	60	47	0.004

CR = no emetic episodes, rescue or withdrawal

NON-CLINICAL SAFETY DATA

A study in cloned human cardiac ion channels has shown ondansetron has the potential to affect cardiac repolarisation via blockade of hERG potassium channels at clinically relevant concentrations.

Dose-dependent QT prolongation has been observed in a thorough QT study in human volunteers (see section CLINICAL PHARMACOLOGY).

Reproductive toxicity

See section PREGNANCY, LACTATION, FEMALES AND MALES OF REPRODUCTIVE POTENTIAL.

Incompatibilities

None reported.

STORAGE

Special precautions for storage

Store below 30°C.

Zofran should not be used after the date marked “EXP” on the pack.

Zofran must be kept out of the reach and sight of children.

Nature and Contents of Container

Ondansetron tablets are available in blister packs of 30's.

Instructions for use and handling

None.

Product Registration Holder:

Sandoz Products Malaysia Sdn. Bhd.
Unit 1202, Level 12, Uptown 1,
No. 1, Jalan SS 21/58, Damansara Uptown,
47400 Petaling Jaya, Selangor, Malaysia.

Malaysia Package Leaflet

Information issued: Mar 2021

Revision date: Jan 2024