

# OSPAMOX FILM-COATED TABLET

Amoxicillin Trihydrate (500 mg)

## What Is In This Leaflet:

1. What is Ospamox used for
2. How Ospamox works
3. Before you use Ospamox
4. How to use Ospamox
5. While you are using Ospamox
6. Side effects
7. Storage and disposal of Ospamox
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of Revision

## What Is Ospamox Used For

Ospamox is used to treat infections caused by bacteria in different parts of the body such as:

- Infections of the ear, throat, nose and sinuses
- Chest infections such as bronchitis and pneumonia
- Infections affecting urinary and genital tract
- Prevention of heart infections during mouth or throat surgery
- Gastrointestinal infections
- Skin and soft tissues infections

Your doctor may have prescribed Ospamox for another reason.

## How Ospamox Works

Ospamox is an antibiotic. It belongs to a group of antibiotics called penicillins which is amoxicillin. It is used to treat infections caused by bacteria which are sensitive to amoxicillin.

## Before You Use Ospamox

- When you must not use it
- If you are allergic (hypersensitive) to amoxicillin, penicillins, cephalosporins or to any of the other ingredients of this medicine

### Pregnancy

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine. As a precautionary measure, amoxicillin should only be used during pregnancy if in the judgement of the physician the potential benefits outweigh the possible hazards.

### Breast-feeding

This medicine passes into breast

milk in small quantities, therefore in some cases it may be necessary to stop breastfeeding.

### - Before you start use it

Talk to your doctor or pharmacist before taking Ospamox if you have:

- any allergic conditions
- fits/ convulsions
- kidney disease
- severe and persistent diarrhea

### - Taking other medicines

Tell your doctor or pharmacist, if you are taking, have recently taken or might take any other medicines. Some medicines can cause problems if you take them with Ospamox.

Take care with the following medicines:

- Probenecid: reduces the kidney elimination of Ospamox resulting in higher blood concentration of Ospamox.
- Allopurinol (used to treat gout): you may have a higher risk of skin reactions.
- Bacteriostatic antibiotics such as tetracyclines and chloramphenicol: combination causes Ospamox to lose efficacy.
- Anticoagulants
- Methotrexate

## How To Use Ospamox

### - How much to use

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose for adults is 1,500 mg to 2,000 mg daily in two to three divided doses.

### People with kidney disease

Your doctor may have to lower the dose.

### - How to use it

For oral use.

You can take Ospamox either before or after meals.

Swallow your tablets whole with a glass of water. Do not crush or chew the tablets.

### - When to use it

Always take this medicine exactly as your doctor told you. Please ask your doctor or pharmacist if you are not sure.

### - How long to use it

Always keep taking this medicine until the course is finished, even if you feel better. If you stop taking this medicine too soon, the infection may come back. Also, the bacteria may become resistant to the medicine. If you have any further questions on the use of this product, ask your doctor or pharmacist.

### - If you forget to use it

It is important to use your medicine every day. However, if you forget to use one or more doses, take OSPAMOX as soon as you remember and then go on as prescribed.

### - If you use too much (overdose)

If you take too much of this medicine (overdose), talk to your doctor or go to a hospital straight away, and show them the pack. The most likely signs of overdose are nausea, vomiting, diarrhea, abdominal pain.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## While You Are Using It

### - Things you must do

- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking OSPAMOX.
- Tell your doctor immediately if you become pregnant while taking this medication.

### - Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor.
- Do not give OSPAMOX to anyone else, even if they have the same symptoms or condition as you.

### - Things to be careful of

Driving and using machines

This medicine has no known

# OSPAMOX FILM-COATED TABLET

Consumer Medication Information Leaflet (RiMUP)

Amoxicillin Trihydrate (500 mg)

influence on the ability to drive and use machines. In some instances, allergic reactions, dizziness and convulsions have occur. Do not drive or operate machinery until you know how this medicine affects you.

## Side Effects

Like all medicines, OSPAMOX can cause side effects, although not everyone gets them.

If you notice any of the following serious side effects, talk to your doctor immediately or go to a hospital straight away.

Nausea and diarrhea are occasional temporary side effects that may occur.

Very rarely, allergic reactions like rashes, fever, severe or itchy skin, swelling of the lower layer of skin and blood disorder problems [e.g. low levels of platelets (thrombocytopenia), severe lowered white blood cell count (agranulocytosis), decrease in white blood cell (leukopenia), increase in certain white blood cells (eosinophilia)] have occurred.

Like other antibiotics, amoxicillin may cause headache, tiredness, inflammation of tongue, inflammation of mouths and lips, fever, swelling of the lower layer of skin and inflammation of the spaces between renal tubules.

Individuals with inflammation of brain membranes (meningitis) may run an increased risk of developing fits.

Transient increases in liver enzymes are also seen.

Stop taking Ospamox and contact your doctor immediately if you experience any of the following:

- Serious allergic reactions such as Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS). DRESS appears initially as flu-like symptoms with a rash on the face and then with an extended rash, high temperature and enlarged lymph nodes.

Ospamox may cause severe allergy and serious skin reactions.

Stop using Ospamox and seek medical assistance immediately if you experience any of the following symptoms:

- Skin reddening, blisters, rash, fever, sore throat or eye irritation

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

## Storage and Disposal of Ospamox

### - Storage

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

### *Ospamox 500 mg film coated tablet*

Store below 25°C, protect from light and moisture.

### - Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## Product Description

### - What it looks like

The film-coated tablets are white to off-white, oblong, biconvex, scored on both sides.

## Ingredients

### Active substance

The active substance is amoxicillin (as trihydrate).

### Inactive ingredients

Core: magnesium stearate, povidone, sodium starch glycolate, cellulose microcrystalline

Coating: titanium dioxide, talc, hypromellose.

## MAL Number

MAL19890612AZ

## **Manufacturer and Product Registration Holder**

### Manufacturer

Sandoz GmbH  
Biochemiestrasse 10  
6250 Kundl  
Austria

### Product Registration Holder

Sandoz Products Malaysia Sdn. Bhd.  
Unit 1202, Level 12, Uptown 1,  
No. 1, Jalan SS 21/58, Damansara  
Uptown, 47400 Petaling Jaya,  
Selangor, Malaysia

## **Date of Revision**

MAR 2024

## **Serial Number**

NPRA(R2/1)200117/335