

LOT/MNF/EXP:



After first opening: The product should be used immediately. Any unused solution should be discarded.
After dilution: After dilution with infusion fluids, to reduce microbiological hazard, the diluted solution should be used as soon as practical. If storage of the diluted solution is necessary, solutions should be stored at room temperature (25°C and below) for no more than 6 hours or under refrigeration at 2°C to 8°C for no more than 24 hours.

Do not freeze. Keep the vial in the outer carton in order to protect from light.
When not protected from light, the vial should be used within 5 days.
Store at or below 30°C.
Keep out of the sight and reach of children.
Each vial of 2ml contains 200mg of sugammadex (as sugammadex sodium).
1ml contains 100mg sugammadex (as sugammadex sodium).
Intravenous use. For single use only. Read the package leaflet before use.

Manufacturer
Lek Pharmaceuticals d.d.
Verovskova Ulica 57, Ljubljana, 1526,
Slovenia.

CONTROLLED MEDICINE / UBAT TERKAWAL

Product Registration Holder
Sandoz Products Malaysia Sdn. Bhd.
Unit 1202, Level 12, Uptown 1,
No. 1, Jalan SS 21/58
Damansara Uptown,
47400 Petaling Jaya, Selangor
JAUHI UBAT DARI KANAK-KANAK

MALxxxxxxxxAZ
SIN16880P

Sugammadex Sandoz
100 mg/mL
Solution for
injection
10 vials
200mg/2ml
SANDOZ

Sugammadex Sandoz
100 mg/mL
Solution for
injection
10 vials
200mg/2ml
SANDOZ

Sugammadex Sandoz
100 mg/mL
Solution for injection
10 vials 200mg/2ml
SANDOZ