

# Lorpitan Trio Hard Capsule

Aprepitant (80 mg, 125 mg)

## What is in this leaflet

1. What Lorpitan Trio (Aprepitant Capsule) is taken for
2. How Lorpitan Trio (Aprepitant Capsule) works
3. Before you take Lorpitan Trio (Aprepitant Capsule)
4. How to take Lorpitan Trio (Aprepitant Capsule)
5. While you are taking Lorpitan Trio (Aprepitant Capsule)
6. Side effects
7. Storage and Disposal of Lorpitan Trio (Aprepitant Capsule)
8. Product description
9. Manufacturer and product registration holder
10. Date of revision
11. Serial number

## What Lorpitan Trio (Aprepitant Capsule) is taken for

Lorpitan Trio (Aprepitant Capsule) is taken **ALONG WITH OTHER MEDICINES** to prevent and control nausea (sick feeling) and vomiting caused by your cancer chemotherapy treatment.

## How Lorpitan Trio (Aprepitant Capsule) works

Lorpitan Trio (Aprepitant Capsule) is a member of a class of medicines called neurokinin 1 (NK<sub>1</sub>) receptor antagonists works by blocking signals to the area of the brain that controls nausea and vomiting.

## Before you take Lorpitan Trio (Aprepitant Capsule)

### - When you must not take it

Do not take Lorpitan Trio (Aprepitant Capsule) if you are allergic to any of its ingredients.

### - Before you start to take it

Tell your doctor about any of the following that apply to you:

- any past or present medical problems.

- any allergies.
- all medicines that you are taking or plan to take, even those you can get without a prescription or herbal products.
- Any intolerance to some sugars as this medicine contains sucrose.

### Use in children

Usage of Lorpitan Trio (Aprepitant Capsule) has not been adequately studied in children. Therefore, Lorpitan Trio (Aprepitant Capsule) should not be given to children under 18 years old.

### Use in the elderly

Lorpitan Trio (Aprepitant Capsule) works equally well in and is equally well-tolerated by older and younger adults. No dosage adjustment is necessary in the elderly.

### - Taking other medicines

Do not take Lorpitan Trio (Aprepitant Capsule) with medicines containing pimozide (used to treat psychiatric illness), terfenadine, astemizole (used for hay fever and other allergic conditions), or cisapride (used for treating digestive problems).

Taking Lorpitan Trio (Aprepitant Capsule) with these medications could result in serious or life-threatening problems

Your doctor may check that your medicines are working properly together if you are taking other medicines such as:

- anti-anxiety medicines (such as alprazolam)
- birth control medicines (which may not work as well)
- warfarin (a blood thinner)
- ketoconazole (an antifungal)
- rifampin or clarithromycin (antibiotic)
- paroxetine (a medicine used to treat a certain type of depression)
- diltiazem (a medicine used to treat high blood pressure)

- dexamethasone (a type of corticosteroids to treat inflammatory conditions)
- carbamazepine (a medicine used to treat seizures)
- St John's wort (a herbal medicine to treat mental disorder)
- tolbutamide (a medicine used to treat diabetes)
- phenytoin (a medicine used to treat seizures)

## How to take Lorpitan Trio (Aprepitant Capsule)

- How much to take and when to take  
Lorpitan Trio (Aprepitant Capsule) 3-Day Regimen

• **Day 1 (Day of chemotherapy):**  
Lorpitan Trio (Aprepitant Capsule) 125 mg will be given to you by mouth 1 hour before you start your chemotherapy treatment on Day 1.

• **Day 2 and Day 3 (Both days after chemotherapy):**  
Take one 80 mg capsule of Lorpitan Trio (Aprepitant Capsule) each morning for the 2 days following your chemotherapy treatment.

Lorpitan Trio (Aprepitant Capsule) may be taken with or without food.

### - How long to take

Lorpitan Trio (Aprepitant Capsule) is taken for 3 days.

### - If you forget to take it

Try to take Lorpitan Trio (Aprepitant Capsule) as prescribed. However, if you miss a dose, contact your physician for further instructions.

### - If you take too much (overdose)

If you take more than the prescribed dosage, contact your doctor immediately.  
Drowsiness and headache were reported in one person who taken too much of aprepitant.

# Lorpitan Trio Hard Capsule

Aprepitant (80 mg, 125 mg)

## While you are taking Lorpitan Trio (Aprepitant Capsule)

### - Things you must do

Tell your doctor if you are:

- Pregnant or plan to become pregnant.
- Breast-feeding or plan to breast-feed.

Lorpitan Trio (Aprepitant Capsule) should be taken during pregnancy and breastfeeding only if the potential benefit justifies the potential risk to the mother and the fetus.

### - Things you must not do

Remember that your doctor has prescribed this medicine only for you. Never give it to anyone else.

### - Things to be careful of

While good nutrition is important for everyone, it is especially important for people on chemotherapy. Small, frequent meals or eating a snack before your chemotherapy treatment may also help you tolerate it better. Talk to your healthcare professional for further information.

## Side Effects

Any medicine may have unintended or undesirable effects, so-called side effects.

Like all medicines, Lorpitan Trio (Aprepitant Capsule) may cause side effects. In studies, side effects usually have been mild or moderate. They generally have not caused people to stop taking Lorpitan Trio (Aprepitant Capsule)

The most common side effects in cancer patients taking the 125 mg/80 mg regimen of Lorpitan Trio (Aprepitant Capsule) included hiccups, fatigue, constipation, headache, indigestion, increased liver enzymes, and loss of appetite.

The following side effects have been reported in general use with Lorpitan Trio (Aprepitant Capsule):

- Allergic reactions, which may be serious, and may include hives, rash and itching and cause difficulty in breathing or swallowing. If you have allergic reaction, stop taking Lorpitan Trio (Aprepitant Capsule) and call your doctor right away.
- Severe skin reactions may occur rarely.

Other side effects may also occur rarely, and as with any other medicines, some side effects may be serious. Ask your doctor or pharmacist for more information. Both have a more complete list of side effects. Tell your doctor or pharmacist promptly about these or any other unusual symptoms.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) [Consumers □ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

## Storage and Disposal of Lorpitan Trio (Aprepitant Capsule)

### - Storage

Store below 30°C. Protect from moisture and light.

Keep Lorpitan Trio (Aprepitant Capsule) and all medicines safely away from children.

### - Disposal

- Do not take this medicine after the date of expiry on the packaging.
- Do not take Lorpitan Trio (Aprepitant Capsule) if you notice any changes in the appearance of the products. Contact your doctor or pharmacist.
- Do not throw away any medicines via waste water or household waste. Ask your pharmacist how to throw away medicines you no

longer use. These measures will help protect the environment.

- You may obtain further information from your doctor or pharmacist, who has more detailed information.

## Product Description

### - What it looks like

Lorpitan Trio (Aprepitant 125mg Capsule):

White to off white spherical pellets, capsule size 1, Opaque pink head and opaque white body, black imprint “125 mg”

Lorpitan Trio (Aprepitant 80mg Capsule):

White to off white spherical pellets, capsule size 2, Opaque white head and body, black imprint “80mg”

### - Ingredients

- Active ingredient:

Lorpitan Trio (Aprepitant 125mg Hard Capsule):  
125mg aprepitant per capsule

Lorpitan Trio (Aprepitant 80mg Hard Capsule):  
80mg aprepitant per capsule

- Inactive ingredients:

hydroxypropyl cellulose, sodium lauryl sulphate, sucrose, Spheres of Cellulose microcrystalline, purified water

- Composition of Capsule shell:

Cap: Titanium dioxide (E 171), Red iron oxide (E 172), Gelatin  
Body: Titanium dioxide (E 171), Gelatin

- Composition of the black ink printed on the capsule:

Shellac (NF, Ph. Eur.), Ammonia (NF, Ph. Eur.), Propylene glycol (USP, Ph. Eur.), Potassium hydroxide (NF, Ph. Eur.), Black iron oxide (E172) (NF)

# Lorpitan Trio Hard Capsule

Aprepitant (80 mg, 125 mg)

---

- MAL number

Lorpitan Trio (Aprepitant 125mg  
Hard Capsule):  
MALxxxxxxxxAZ

Lorpitan Trio (Aprepitant 80mg  
Hard Capsule):  
MALxxxxxxxxAZ

**Manufacturer**

Ethypharm  
Chemin De La Poudriere, Le Grand  
Quevilly, 76120, France

**Product Registration Holder**

GIRAFFES ENTERPRISE SDN.  
BHD.  
C18-3A, 3 TWO SQUARE, JALAN  
19/1, 46300, PETALING JAYA,  
SELANGOR

**Date of revision**

10-April-2025

**Serial Number**

NPRA (R2) 24/101