

JARDIN SALIC 20 SALICYLIC ACID 20% W/W OINTMENT

Salicylic Acid (20% w/w)

What is in this leaflet

1. What Jardin Salic 20 is used for
2. How Jardin Salic 20 works
3. Before you use Jardin Salic 20
4. How to use Jardin Salic 20
5. While you are using it
6. Side effects
7. Storage and Disposal of Jardin Salic 20
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision
11. Serial Number

What Jardin Salic 20 is used for

Jardin Salic 20 is used for removal of warts.

How Jardin Salic 20 works

Jardin Salic 20 works by softening the outer layer of your skin allowing it to loosen and shed.

Before you use Jardin Salic 20**- When you must not use it**

Do not take this medicine if:

- you are allergic to Salicylic Acid or any other ingredients of this medicine (listed in Product Description)

- Before you start to use it

Check with your doctor or pharmacist before taking Jardin Salic 20 if you:

- allergic to any other medicines.

Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Inform your doctor or pharmacist if you are taking:

- Any other medicated topical agents, e.g. benzoyl peroxide, topical retinoids including isotretinoin, alitretinon and tazarotene.
- Abrasive soap and cleanser.
- Cosmetics or soap that dry the skin or are designed to peel/exfoliate.
- Protein bound drugs.
- Alcohol containing preparations.

How to use Jardin Salic 20

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- How much to use

Apply a thin layer to entire warts.

- When to use it

Use as directed by your doctor or pharmacist.

Apply twice daily, in the morning and at night.

Subsequently, it may then be possible to maintain improvement by less frequent application.

- How long to use it

Use this medicine for the duration that your doctor recommends.

Apply the medication as instructed until the wart clears up, for up to 12 weeks or as directed by your doctor.

If the wart persists after 12 weeks, consult your doctor.

- If you forget to use it

Apply the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to apply the medicine and skip the missed dose. Do not apply a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much of this medicine may cause symptoms of Salicylism.

While you are using Jardin Salic 20**- Things you must do**

Use your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using Jardin Salic 20.

Tell your doctor immediately if you become pregnant while using this medication.

Always wash your hands before and after using the ointment.

- Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Jardin Salic 20 to anyone else, even if they have the same symptoms or condition as you.

Do not use at broken skin.

Avoid contact with mouth, eyes and mucus membrane.

Do not use for warts on the face, moles, birthmarks, hairy warts or on genital/anal warts.

Do not use over large skin areas for long period (more than 12 weeks) or more than as directed by your doctor.

JARDIN SALIC 20 SALICYLIC ACID 20% W/W OINTMENT

Salicylic Acid (20% w/w)

- Things to be careful of**Driving and using machines**

Be careful driving or operating machinery until you know how Jardin Salic 20 affects you.

Be careful to apply it only to the entire warts and not the surrounding skin.

If irritation or sensitization develops, discontinue treatment and consult your doctor.

Do not use this product after the expiry date or if the tube is leaking or the tube nozzle cover is broken.

Protect from heat, moisture and light.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Date of revision

31/12/2025

Serial Number

NPRA(R2) 25/054

Side effects

Like all medicines, Jardin Salic 20 can cause side effects, although not everybody gets them.

Possible side effects are:

- Contact dermatitis or skin irritation; flushing, unusually warm skin, burning sensation, itching and stinging of skin.
- Excessive drying and peeling of skin.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Jardin Salic 20**- Storage**

Keep out of the reach and sight of children.

Store below 30 °C.

Keep the tube tightly closed.

Product Description**- What it looks like**

White or almost white, translucent ointment.

- Ingredients

- Active ingredient
 - Salicylic Acid
- Inactive ingredients
 - White Soft Paraffin

- MAL number:

MAL25076034XZ

Manufacturer

Jardin Manufacturing Sdn Bhd,
Lot 1, 2, 3 & 4,
Kawasan Industri Batu 7, 23000
Dungun, Terengganu,
Malaysia.

Product Registration Holder

Jardin Manufacturing Sdn. Bhd,
No. 12-1, Jalan Delta U6/19,
Sunway Subang Business Park,
Jalan TUDM, 40150
Shah Alam, Selangor,
Malaysia.