

Jardin Salic 10 Salicylic Acid 10% w/w Ointment

Description

A white or almost white and translucent ointment.

Pharmacodynamics

The ointment is used as a keratolytic agent.

Pharmacokinetics

Salicylic acid may be percutaneously absorbed. However, there is no evidence of any systemic absorption from the use of Salicylic Acid Ointment..

Product Indication

For treatment of seborrhoeic dermatitis, scalp, psoriasis and hyperkeratotic skin disorders.

Recommended Dose

Apply sparingly to the affected skin areas twice daily, in the morning and at night, until improvement occurs. Subsequently, it may be possible to maintain improvement with less frequent application..

Mode Of Administration

Topical.

Contraindication

Contraindicated in patients displaying salicylate hypersensitivity or sensitivity to any other ingredient in the preparation of Jardin Salic 10 (Salicylic Acid 10% w/w Ointment).

Warnings and Precautions

1. Avoid broken skin.
2. For external use only.
3. Avoid contact with mouth, eyes and mucous membranes.
4. If irritation or sensitization develops, discontinue treatment.
5. If large areas of skin are treated, the systemic absorption of topical Salicylic Acid will be enhanced and may lead to Salicylate toxicity.
6. When using salicylic acid preparations, do not use any of the following preparations on the affected area, unless directed to do so by your doctor.
 - Any other medicated topical agents. e.g: benzoyl peroxide, topical retinoids.
 - Abrasive soaps and cleansers.
 - Cosmetics or soaps that dry the skin or are designed to peel/exfoliate.
7. Salicylic acid may cause dermatitis. Extensive prolonged use of topical Salicylic Acid may enhance the systemic absorption and may lead to Salicylate toxicity.

Pregnancy and Lactation

Topical Salicylic Acid should be used only if potential benefit outweighs the potential risk to the fetus. Topical Salicylic Acid should not be used extensively in large amount or for prolonged period in pregnant women.

Interactions With Other Medications

Other topical preparations, mainly isotretinoin, tretinoin, tazarotene.

Side Effects and Adverse Effects/Undesirable Effects

1. Contact dermatitis or skin irritation; flushing; unusually warm skin.
2. Excessive drying and peeling of skin.

Signs and Symptoms of Overdose and Treatment

Excessive prolonged use of Jardin Salic10 Ointment may result in symptoms of Salicylism. If this occurs, symptomatic treatment should be initiated.

Storage Conditions

Keep the tube tightly closed.

Store below 30 °C.

Protect from heat, moisture and light.

Keep medicine out of reach of children.

Dosage Form

Ointment.

Packaging

15 gram in printed aluminium tube.

1 (one) tubes per unicarton.

Shelf Life

24 months.

This insert contains basic prescribing information only. For further information, please consult your doctor or pharmacist.

Product Registration Holder

Jardin Manufacturing Sdn. Bhd,
No. 12-1, Jalan Delta U6/19,
Sunway Subang Business Park,
Jalan TUDM, 40150 Shah Alam,
Selangor.

Manufacturer

Jardin Manufacturing Sdn. Bhd,
Lot 1, 2, 3 & 4,
Kawasan Industri Batu 7,
23000 Dungun,
Terengganu.