

JARDIN PANADINOL SYRUP (PARACETAMOL 120 MG/5ML)

Paracetamol (120mg/5ml)

What is in this leaflet

1. What Jardin Panadinol Syrup is used for
2. How Jardin Panadinol Syrup works
3. Before you use Jardin Panadinol Syrup
4. How to use Jardin Panadinol Syrup
5. While you are using it
6. Side effects
7. Storage and Disposal of Jardin Panadinol Syrup
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision
11. Serial Number

What Jardin Panadinol Syrup is used for

Indicated for the relief of fever, headache and symptoms of cold and flu, toothache, discomfort of teething and fever after vaccination.

How Jardin Panadinol Syrup works

Paracetamol is a centrally acting analgesic and antipyretic with minimal anti-inflammatory properties.

Before you use Jardin Panadinol Syrup

- When you must not use it

Do not take this medicine if:

- you are allergic to paracetamol or any other ingredients of this medicine (listed in Product Description)

- Before you start to use it

Check with your doctor or pharmacist before taking Jardin Panadinol Syrup if you:

- Kidney problems.
- Liver problems.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Inform your doctor or pharmacist if you are taking:

- metoclopramide or domperidone (used to treat nausea and vomiting)
- cholestyramine (used to treat high cholesterol).
- anticoagulants (drugs that thin the blood, such as warfarin).
- anticonvulsants (drugs to treat epilepsy).

How to use Jardin Panadinol Syrup

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- How much to use

For an accurate dosing, weight your child and work out the dosage to be given based on a 15mg of Paracetamol for every kg body weight given every 4 or 6 hours. e.g: If your child weighs 20 kg: $20 \text{ kg} \times 15 \text{ mg} = 300 \text{ mg}$ $300 \text{ mg} / 120\text{mg} \times 5\text{ml} = 12.5 \text{ ml}$
Maximum daily dose: 60mg/kg presented in divided doses of 10-15mg/kg throughout the 24 hour period.

The table below provides an illustrative guide for dosing based on the child's weight. Please note that this is just an example, and it is recommended to consult with a healthcare professional for precise dosing information.

Child's Age	Child's Weight (kg)	How much	How often (in 24 hours)
3-6 months	4	2.5ml	4 times
6-24 months	8	5ml	4 times
2-4 months	12	7.5ml	4 times
4-8 months	14	10ml	4 times
8-10 months	24	15ml	4 times
10-12 months	32	20ml	4 times

- Below 3 months: 5 to 10mg/kg (on doctor's advise only).
- If the fever persists for more than 24 hours (4 doses), seek medical advice. This precaution ensures prompt diagnosis of any fever that might be indicative of a serious infection.
- Do not give more than 4 doses in any 24 hour period.
- Leave at least 4 hours between doses.
- Do not give this medicine to your child for more than 3 days without speaking to your doctor or Pharmacist.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Use this medicine for the duration that your doctor recommends.

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much of this medicine may cause damage to the liver and kidney.

While you are using Jardin Panadinol Syrup

- Things you must do

Use your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using Jardin Panadinol Syrup.

JARDIN PANADINOL SYRUP (PARACETAMOL 120MG/5ML)

Paracetamol (120mg/5ml)

Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Jardin Panadinol Syrup to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Please always consult your doctor or pharmacist before giving your children this product especially when there is a known to hypersensitivity.

Side effects

Like all medicines, Jardin Panadinol Syrup can cause side effects, although not everybody gets them.

Possible side effects are:

- Adverse effects are rare and usually mild, although haematological reactions have been reported.
- Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/ Toxic Epidermal Necrolysis have been reported.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blister or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug

Reaction Monitoring by visiting the website nppra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Jardin Panadinol Syrup

- Storage

Keep out of the reach and sight of children.

Store below 30 °C.

Keep the bottle tightly closed

Do not use this product after the expiry date

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

A clear red coloured syrup with strawberry flavour.

- Ingredients

- Active ingredient
 - Paracetamol
- Inactive ingredients
 - Methylparaben
 - Glycerine
 - Propylene Glycol
 - Sorbitol
 - Strawberry Flavor
 - Colour Carmosine
 - Citric Acid
 - Sucralose
 - Sodium Citrate
 - Purified Water

- MAL number:

MAL*****

Manufacturer

Jardin Manufacturing Sdn Bhd,
Lot 1,2,3 & 4,
Kawasan Industri Batu 7, 23000
Dungun, Terengganu,
Malaysia.

Product Registration Holder

Jardin Manufacturing Sdn Bhd,
No 12-1, Jalan Delta U6/19,
Sunway Subang Business Park,
Jalan TUDM, 40150
Shah Alam, Selangor,
Malaysia

Date of revision

15/06/2026

Serial Number

NPRA(R2)26/102