

Desloratadine 5 mg

What is in this leaflet:

1. What DESTAVELL is used for
2. How DESTAVELL works
3. Before you use DESTAVELL
4. How to use DESTAVELL
5. While you are using it
6. Side effects
7. Storage and Disposal of DESTAVELL
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What DESTAVELL is used for

DESTAVELL relieves symptoms associated with allergic rhinitis (inflammation of the nasal passages caused by an allergy, for example, hay fever or allergy to dust mites). These symptoms include sneezing, runny or itchy nose, itchy palate, and itchy, red or watery eyes.

DESTAVELL is also used to relieve the symptoms associated with urticaria (a skin condition caused by an allergy). These symptoms include itching and hives.

Relief of these symptoms lasts a full day and helps you to resume your normal daily activities and sleep.

How DESTAVELL works

DESTAVELL is an antiallergy medicine that does not make you drowsy. It helps control your allergic reaction and its symptoms.

Before you use DESTAVELL

- When you must not use it

Do not use DESTAVELL if you are allergic (hypersensitive) to desloratadine, to any of other ingredients of DESTAVELL or to loratadine.

- Before you start to use it

Talk to your doctor or pharmacist before taking DESTAVELL if you have medical or familial history of seizures.

Efficacy and safety of DESTAVELL in children under 12 years of age have not been established.

Pregnancy and breast-feeding

- Ask your doctor or pharmacist for advice before taking DESTAVELL during pregnancy and breast-feeding.
- If you are pregnant or nursing a baby, taking DESTAVELL is not recommended.

Important information about some of the ingredients of DESTAVELL

DESTAVELL contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

- Taking other medicines

There are no known interactions of DESTAVELL with other medicines.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

How to use DESTAVELL

- How much to use

Adults or adolescents (12 years of age and older): take one tablet once a day.

- When to use it

Swallow the tablet whole with water, with or without food.

- How long to use it

Regarding the duration of treatment, your physician will determine the type of allergic rhinitis you are suffering from and will determine for how long you should take DESTAVELL.

If your allergic rhinitis is intermittent (presence of symptoms for less than 4 days per week or for less than 4 weeks), your physician will recommend you a treatment schedule that will depend on the evaluation of the history of your disease.

If your allergic rhinitis is persistent (presence of symptoms for 4 days or more per week and for more than 4 weeks), your physician may recommend you a longer-term treatment.

For urticaria, the duration of treatment may be variable from patient to patient and therefore you should follow the instructions of your physician.

- If you forget to use it

If you forget to take your dose on time, take it as soon as possible, then go back to your regular dosing schedule. Do not take a double dose to make up for forgotten dose.

- If you use too much (overdose)

Take DESTAVELL only as it is prescribed for you. No serious problems are expected with accidental overdose. However, if you take more DESTAVELL than you were told to, contact your doctor or pharmacist.

Contact your doctor immediately or go to Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

- Take your medicine exactly as your doctor has told you.
- Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

- Do not stop taking your medicines unless advised by your doctor.
- Do not take any new medicines without consulting your doctor.
- Do not give DESTAVELL to anyone else, even if they have the same symptoms or condition.

- Things to be careful of

Driving and using machines

At the recommended dose, DESTAVELL is not expected to cause you to be drowsy or less alert. However, very rarely some people experience drowsiness, which may affect their ability to drive or use machine.

Side effects

Like all medicines, DESTAVELL can cause side effects. In adults, side effects were about the same as with dummy tablet. However, fatigue, dry mouth and headache were reported more often than with a dummy tablet.

DESTAVELL 5 mg Film-Coated Tablet

Consumer Medicine Information Leaflet (RiMUP)

Desloratadine 5 mg

During the marketing of DESTAVELL, cases of severe allergic reactions (difficulty in breathing, wheezing, itching, hives and swelling) and rash have been reported very rarely. Cases of palpitations (irregular rapid pumping of the heart), rapid heartbeat, drowsiness, liver inflammation, abnormal liver function tests and increased appetite have also been reported very rarely.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.moh.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccine (AEFI)].

Storage and Disposal of DESTAVELL

- Storage

- Keep out of the reach and sight of children.
- Do not store above 30°C. Store in the original package. Protect from excessive moisture.
- Do not use after the expiry date stated on the carton and blister.
- Tell your pharmacist if you notice any change in the appearance of the tablets.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Blue film-coated tablet, round, convex, plain on one side and breakline on the other.

- Ingredients

- Active ingredient
Desloratadine
- Inactive ingredients
Calcium hydrogen phosphate dihydrate, microcrystalline cellulose PH 102, pregelatinized starch, talc, hypromellose,

lactose monohydrate, macrogols 6000, titanium dioxide, FD&C Blue No.2 Al-lake (C.I 73015), purified water

- Packing

Box, 3 strips @ 10 film-coated tablets

- MAL Number

MAL22086029AZ

Manufacturer



PT Novell Pharmaceutical Laboratories
Jl. Wanaherang No. 35, Tlajung Udik,
Gunung Putri, Bogor 16962 – Indonesia

Product Registration Holder

MEDHUB PHARMA BERHAD
PS-G-2, Pelangi Damansara Sentral,
Persiaran Surian, 47800 Petaling Jaya,
Selangor, Malaysia

Date of revision

03/05/2024

Serial Number:

NPRA(R2/1)240423/1257