PATADAY EYE DROPS

Olopatadine hydrochloride (2mg/ml)

What is in this leaflet

1. What PATADAY is used for
2. How PATADAY works
3. Before you use PATADAY
4. How to use PATADAY
5. While you are using PATADAY
6. Side effects of PATADAY
7. Storage and Disposal of PATADAY
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial Number

What PATADAY is used for

Pataday is used for the treatment of eye itching caused by allergy to house dust, pollen, animal fur, or other allergens (allergic conjunctivitis).

How PATADAY works

Pataday works by reducing the intensity of the allergic reaction.

Before you use PATADAY

- *When you must not use it*

Do not use Pataday if you are allergic to olopatadine or any of the other ingredients of Pataday.

If this applies to you, do not use Pataday and tell your doctor.

Children and adolescents (below 18 years)

Safety and effectiveness in children below the age of 3 years have not been established.

Older people (65 years of age or above)

You can use Pataday if you are aged 65 years or over at the same dose as for other adults.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, you think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

- *Before you start to use it*

Pataday contains benzalkonium chloride which may cause eye irritation and may possibly discolor soft contact lenses. Contact lenses must be removed before administration of Pataday and reinserted at least 15 minutes later.

- *Taking other medicines*

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines, including medicines obtained without a prescription.

How to use PATADAY

Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist or nurse has told you. Check with your doctor or pharmacist or nurse if you are not sure.

- *How much to use*

The recommended dose is one drop in each affected eye once a day.

- *When to use it*

Use once a day or as directed by your doctor or pharmacist.

- *How long to use it*

Continue using Pataday for as long as your doctor recommends.

- *If you forget to use it*

If you forget to use Pataday, continue with the next dose as planned. However, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not use a double dose to make up.

- *If you use too much (overdose)*

If you use more Pataday than you should, rinse it all out with warm water. Do not put in any more drops until it is time for your next regular dose.

While you are using PATADAY

- *Things you must do*

- Only use Pataday in your eye(s).
- To avoid contamination, the dropper tip should not touch any surface. The dropper tip should also not come into contact with the eye as this may cause injury to the eye.
- If a drop misses your eye, try again.
- When closing the eyelids for 2 minutes (using nasolacrimal occlusion), the absorption of medicine in other parts of the body (systemic absorption) is reduced. This may result in a decrease in general side effects affecting other parts of the body and an increase in local activity.
- Keep the bottle tightly closed when not in use.
- If more than one eye medicines are being used, the medicinal products must be administered at least 5 minutes apart. Eye ointments should be administered last.

- *Things you must not do*

- Do not wear a contact lens in case of eye redness.
- Pataday should not be used to treat contact lens related irritation.

Side effects of PATADAY

PATADAY EYE DROPS

Olopatadine hydrochloride (2mg/ml)

Like all medicines, patients treated with Pataday may experience side effects, although not everybody gets them.

If you experience any serious side effects, stop using this medicine and tell your doctor immediately.

The following side effects have been seen with Pataday:
Common: may affect up to 1 in every 10 people

- Effects in the eye: eye discomfort

Uncommon: may affect up to 1 in every 100 people

- Effects in the eye: dry eye, blurred vision, itchy eye, redness, eyelid crusting
- General side effects: headache, disturbed sense of taste (dysgeusia), nasal dryness

Rare: may affect up to 1 in every 1,000 people

- General side effects: dry mouth

Not known: frequency cannot be estimated from the available data

- Effects in the eye: watering eye (lacrimation increased)
- General side effects: allergy, nausea

If you notice any side effects not listed in this leaflet, please inform your doctor or pharmacist or healthcare provider.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of PATADAY

- *Storage*

Do not store above 30°C.
Discard 28 days after opening.

- *Disposal*

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- *What it looks like*

Pataday is a colourless to light yellow solution.

- *Ingredients*

- Active ingredient(s)
2.22 mg of olopatadine hydrochloride in 1 mL solution (0.2%w/v)

- Inactive ingredients
Benzalkonium chloride, edetate disodium, povidone, sodium chloride, dibasic sodium phosphate (anhydrous), sodium hydroxide and/or hydrochloric acid, purified water

- *MAL number:*

MAL08021516ARZ

Manufacturer

Novartis Manufacturing NV,
Rijkweg 14,
2870 Puurs-Sint-Amands
Belgium

Product Registration Holder

Novartis Corporation (M) Sdn. Bhd.
Level 18, Imazium,
No. 8, Jalan SS21/37,
Damansara Uptown,
47400 Petaling Jaya, Selangor, Malaysia.

Information Issued

17-Jul-2020

Date of revision

14-Jan-2025

Serial Number