

Height 165mm



Exzolt 10 mg/ml solution for use in drinking water for chickens.  
Fluralaner 10mg/ml.

Treatment of poultry red mite (*Dermanyssus gallinae*) infestation in pullets, breeders and layer hens.

For Dosage and Administration, Warnings/ Precautions, and Disposal of Containers: Refer to package leaflet.

Withdrawal period(s):  
Meat and offal: 14 days.  
Eggs: zero days.

Do not store above 30°C. Protect from moisture.  
Once opened, use within 1 year.  
Once diluted, use within 24 hours.

**Keep out of the sight and reach of children / Jauhi daripada kanak-kanak.**  
**For Animal Treatment Only.**  
**Controlled Medicine / Ubat Terkawal.**

**Manufacturer:** Intervet Productions S.A.  
Rue de Lyons, 27460 Igoville, France.  
**Product Registration Holder:** Intervet (M) Sdn. Bhd.  
B-22-1& B-22-2, The Ascent Paradigm,  
No.1, Jalan SS 7/26A, Kelana Jaya,  
47301 Petaling Jaya, Selangor,  
Malaysia.


Product Registration No.: MAL20046153HAR  
Product of Intervet International B.V., Boxmeer,  
The Netherlands

**4 litre**



Base label - 110mm

PSB920F36  
Version 01 R1  
Base label (BL) 165x110mm  
FAF Unfolded 165x370mm  
FAF Folded 165x100mm  
8 pages



PSB920F36 01 R1  
BL 165x110mm, FAF Unfolded 165x370mm, FAF Folded 165x100mm, 8 pages  
CMYK

#### PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Antiparasitic products, ectoparasiticides for systemic use, isoxazolines.  
ATCvet code: QP53BE02

#### PHARMACODYNAMIC PROPERTIES

Fluralaner is an acaricide and an insecticide which has a high potency against poultry mites, mostly by exposure via feeding, i.e. it is systemically active against the target parasites.

Fluralaner is a potent inhibitor of parts of the arthropod nervous system by acting antagonistically on ligand-gated chloride channels (GABA-receptor and glutamate-receptor). In molecular on-target studies on insect gamma-aminobutyric acid (GABA) receptors of flea and fly, dieldrin resistance does not affect fluralaner.

The onset of activity against *Dermanyssus gallinae* is within four hours of the mites starting to feed on treated chickens.

The treatment kills mites feeding on treated chickens and stops egg production from female mites for 15 days after the first administration of the product. This activity breaks the mite life cycle.

*In vitro* bio-assays show that fluralaner is effective against parasites having proven field resistance, including organophosphates, pyrethroids and carbamates.

As demonstrated in a multi-site field study performed in commercial egg production farms, elimination of mites from infested chickens following treatment is associated with a statistically significant improvement in behavioral parameters indicative of animal welfare (reduction of night-time activity and head scratching, head shaking and preening of own plumage at night and during day-time) as well as a reduction of blood corticosterone concentration.

Page 6 - 90mm

#### PHARMACOKINETIC PARTICULARS

After oral administration, fluralaner is absorbed rapidly from the medicated drinking water, reaching maximum plasma concentrations 36 hours after the first dose and 12 hours after the second dose. The bioavailability is high, with approximately 91% of the dose absorbed following oral administration. Fluralaner is highly bound to protein. Fluralaner is widely distributed throughout the body, with the highest concentrations reported in the liver and skin/ fat. No significant metabolites are observed in chickens, and fluralaner is mainly eliminated via the hepatic route. The apparent elimination half-life is approximately 5 days following oral administration.

#### ENVIRONMENTAL PROPERTIES

Fluralaner has been shown to be very persistent in soil under both, aerobic and anaerobic conditions. Fluralaner degrades in aquatic sediment under anaerobic conditions while it has been shown to be very persistent under aerobic conditions.

#### LIST OF EXCIPIENTS

Alpha-tocopherol (all-rac- $\alpha$ -tocopherol) Diethylene glycol monoethyl ether  
Polysorbate 80

#### SHELF LIFE

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.  
Shelf life after first opening of the container: 1 year.  
Shelf life of the medicated drinking water: 24 hours.

#### WITHDRAWAL PERIOD(S)

Meat and offal: 14 days.  
Eggs: zero days.

#### SPECIAL STORAGE PRECAUTIONS

Do not store above 30°C.  
Protect from moisture.

Page 7 - 90mm

#### SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED VETERINARY MEDICINAL PRODUCT OR WASTE MATERIALS DERIVED FROM THE USE OF SUCH PRODUCTS

Exzolt should not enter water courses as this may be dangerous for aquatic invertebrates.  
Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

#### NATURE AND COMPOSITION OF IMMEDIATE PACKAGING

Colourless high density polyethylene (HDPE) bottle closed with an aluminium/polyester foil seal and a blue child-resistant polypropylene screw cap.  
Pack sizes: bottles of 1 litre or 4 litres.  
Not all pack sizes may be marketed.

#### For animal treatment only.

**Keep out of the sight and reach of children/ Jauhi daripada kanak-kanak.**  
**Do not use this veterinary medicinal product after the expiry date which is stated on the label after EXP.**  
**Product of Intervet International B.V., Boxmeer, The Netherlands.**

#### MANUFACTURER

Intervet Productions S.A.  
Rue de Lyons  
27460 Igoville  
France.

#### PRODUCT REGISTRATION HOLDER

Intervet (M) Sdn. Bhd.  
B-22-1 & B-22-2, The Ascent Paradigm,  
No.1, Jalan SS 7/26A, Kelana Jaya,  
47301 Petaling Jaya, Selangor,  
Malaysia.

Product registration no.: MAL20046153HAR  
DATE OF REVISION: 15<sup>th</sup> January 2024

Page 8 - 90mm



Exzolt 10 mg/ml solution for use in drinking water for chickens.  
Fluralaner 10mg/ml.

Treatment of poultry red mite (*Dermanyssus gallinae*) infestation in pullets, breeders and layer hens.

For Dosage and Administration, Warnings/ Precautions, and Disposal of Containers: Refer to package leaflet.

Withdrawal period(s):  
Meat and offal: 14 days.  
Eggs: zero days.

Do not store above 30°C. Protect from moisture.  
Once opened, use within 1 year.  
Once diluted, use within 24 hours.

**Keep out of the sight and reach of children / Jauhi daripada kanak-kanak.**  
**For Animal Treatment Only.**  
**Controlled Medicine / Ubat Terkawal.**

**Manufacturer:** Intervet Productions S.A.  
Rue de Lyons, 27460 Igoville, France.  
**Product Registration Holder:** Intervet (M) Sdn. Bhd.  
B-22-1& B-22-2, The Ascent Paradigm,  
No.1, Jalan SS 7/26A, Kelana Jaya,  
47301 Petaling Jaya, Selangor,  
Malaysia.

Product Registration No.: MAL20046153HAR  
Product of Intervet International B.V., Boxmeer,  
The Netherlands

**4 litre**



190861 R1

Height 165mm

Front cover - 100mm

**NAME OF THE VETERINARY MEDICINAL PRODUCT**  
Exzolt 10 mg/ml solution for use in drinking water for chickens

#### STATEMENT OF ACTIVE SUBSTANCE

Each ml contains:  
**Active substance:**  
10mg fluralaner

#### PHARMACEUTICAL FORM

Solution for use in drinking water.  
Light yellow to dark yellow solution.

#### TARGET SPECIES

Chickens (pullets, breeders and layer hens).

#### INDICATIONS

Treatment of poultry red mite (*Dermanyssus gallinae*) infestation in pullets, breeders and layer hens.

#### CONTRAINDICATIONS

None.

#### ADVERSE REACTIONS

None known.

#### DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For use in drinking water.  
The dose is 0.5 mg fluralaner per kg body weight (equivalent to 0.05 ml of product) administered twice, 7 days apart. The complete course of treatment must be administered for a full therapeutic effect.  
If another course of treatment is indicated, the interval between two courses of treatment should be at least 3 months.

Inside Cover - 100mm

#### Method of administration:

Determine the duration of time (between 4 and 24 hours) over which to administer the medicated water on the treatment day. This period of time must be long enough to allow all the birds to receive the required dose. Estimate how much water birds will consume during treatment based on the previous day's water consumption. The product should be added to a volume of water that the chickens will consume in one day. No other source of drinking water should be available during the medication period.  
Calculate the volume of product needed based on the total weight of all birds in the house to be treated. To ensure administration of the correct dose, the body weight should be determined as accurately as possible and an accurate measuring device should be used for measuring the calculated volume of the product to be administered.  
The required volume of product for each treatment day is calculated from the total body weight (kg) of the entire group of chickens to be treated:

$$\frac{\text{Total body weight (kg)} \times \text{Dose (mg/kg)}}{\text{Concentration (mg/ml)}} = \text{Volume of product (ml)}$$

Therefore 500 ml of product treats 10,000 kg body weight (e.g., 5,000 chickens of 2 kg body weight each) per day of treatment administration.  
The instructions below need to be followed, in the order described, to prepare the medicated water:

- Check the water system to ensure it works properly and is free of leaks; also ensure that water is available to all nipple or bell drinkers.
- For each day of treatment, medicated water must be freshly prepared.
  - Mix the required volume of the product with water into a large medication tank or create a stock solution in a small container. The stock solution must be further diluted with drinking water and administered over time, using a proportioner or dosing pump. Always add product and water simultaneously in order to avoid foaming.

Page 3 - 90mm

It is important to rinse the measuring device used to measure the required product volume during the filling phase in order to ensure that the complete dose is emptied into the medication tank or the stock solution and that no residues remain in the measuring device. Stir the stock solution or the content of the medication tank gently until the medicated water is homogeneous. Connect the medication tank or the proportioner or dosing pump to the drinking water system.

- Make sure the dosing pump is properly set to deliver the medicated water during the predetermined treatment period (hours).
- Prime the drinker lines with medicated water and check to see when medicated water has reached the end of the line. This procedure should be repeated on each day of administration.

After each treatment administration, fill the stock solution container with clean (unmedicated) water to rinse the water lines.

#### SPECIAL WARNINGS

The following practices should be avoided because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- too frequent and repeated use of acaricides from the same class, over an extended period of time
- underdosing, which may be due to underestimation of body weight, misadministration of the product, or lack of calibration of the volume measuring device.

#### Special precautions for use in animals:

Strict biosecurity measures at house and farm level should be implemented to prevent re-infestation of treated houses. To ensure long term control of the mite populations in a treated house, it is essential to treat any other infested poultry in houses in proximity to the treated one.

Page 4 - 90mm

**Special precautions to be taken by the person administering the veterinary medicinal product to animals:**

The veterinary medical product may be slightly irritating to skin and/or eyes.  
Avoid contact with skin, eyes and mucous membranes. Do not eat, drink or smoke while handling the product. Wash hands and contacted skin with soap and water after use of the product.  
In case of eye contact, immediately rinse thoroughly with water.  
If the product is spilled, remove any affected clothes.

#### Use during pregnancy, lactation or lay:

The safety of the veterinary medicinal product has been demonstrated in layers and breeders. The product can be used during lay.

#### Interaction with other medicinal products and other forms of interaction:

None known.

#### Overdose (symptoms, emergency procedures, antidotes), if necessary:

No adverse reactions were observed following the treatment of 3-week old and adult chickens dosed with up to 5 times the recommended dose for 3 times the recommended duration of treatment.  
No negative effects on egg production were observed when layer hens were treated with up to 5 times the recommended dose for 3 times the recommended duration of treatment.  
There were no adverse effects on reproductive performance when breeding chickens were treated with 3 times the recommended dose for twice the recommended duration of treatment.

#### Incompatibilities:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

Page 5 - 90mm

Height 165mm



Exzolt 10 mg/ml solution for use  
in drinking water for chickens.  
Fluralaner 10mg/ml.

Treatment of poultry red mite  
(*Dermanyssus gallinae*)  
infestation in pullets,  
breeders and layer hens.  
For the treatment and control of  
poultry mites.

**For animal treatment only.  
Keep out of the sight and reach of  
children / Jauhi daripada  
kanak-kanak.  
Controlled Medicine / Ubat  
Terkawal.**

**Read the package leaflet before  
use.**

Product of Intervet  
International B.V., Boxmeer, The  
Netherlands

Lot/Lote:

Mfg.date/

Prod.:

EXP./CAD:

Imprint area  
Varnish free area

41



**MSD**

Animal Health



191907 R4