

KLEARIT (Polyethylene Glycol 3350 and Electrolytes for Oral Solution USP)

Polyethylene Glycol 3350 (118g)/Anhydrous Sodium Sulfate (11.36g)/Sodium Bicarbonate (3.37g)/Sodium Chloride (2.93g)/Potassium Chloride (1.484g)

What is in this leaflet

1. What KLEARIT is used for
2. How KLEARIT works
3. Before you use KLEARIT
4. How to use KLEARIT
5. While you are using it
6. Side effects
7. Storage and Disposal of KLEARIT
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision

What KLEARIT is used for

Patients are prepared bycleansing for

- Endoscopic or radiological examinations
- Colonic surgery

How KLEARIT works

KLEARIT increases the amount of water in the intestine, which increases the frequency of stools, the contents of which become increasingly watery and, eventually, flushes the intestine

Before you use KLEARIT

- When you must not use it

Do not take this medicine if:

- you are allergic to Poly ethylene Glycol 3350 or any other ingredients of this medicine (listed in Product Description)

- Before you start to use it

Check with your doctor or pharmacist before taking **KLEARIT**

Pregnancy and lactation

- If you are pregnant or trying to pregnant or if you are breastfeeding, seek advices from your doctor or pharmacist before taking this medicine. KLEARIT should be used only when necessary.
- Taking other medicines
Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Inform your doctor or pharmacist if you are taking other medicines:

Due to the diarrhoea provoked by KLEARIT, the other oral treatments may not be absorbed and should be taken more than 2 hours before taking KLEARIT. Avoid intake of oral treatments before and after laxative ingestion until medical examination is completed. For medicines with small therapeutic range or with short half-life such as digoxin, antiepileptics, coumarins, and immunosuppressants, efficacy can be particularly affected

How to use KLEARIT

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- How much to use

- Adults: Instruct patients to drink a total of up to 2 liters at a rate of 200 mL every 10 minutes, until 2 liters is consumed or the rectal effluent is clear. Rapid drinking of each portion is preferred to drinking small amounts continuously. The first bowel movements should occur approximately one hour after the start of KLEARIT administration. Continue drinking until the Watery stool is clear and free of solid matter.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Use this medicine for the duration that your doctor recommends.

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much of this medicine may cause severe diarrhea and electrolyte imbalance, including hyponatremia and hypokalaemia, as well as dehydration and hypovolaemia with the signs and related symptoms

While you are using KLEARIT

- Things you must do

Use your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using KLEARIT.

Tell your doctor immediately if you become pregnant while using this medication.

Take KLEARIT only with a doctor's prescription.

- Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give KLEARIT to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of *Driving and using machines*

KLEARIT has no influence on the ability to drive and use machines.

Side effects

Like all medicines, KLEARIT can cause side effects, although not everybody gets them.

Possible side effects are:

- Very common (in more than 1 in 10 patients):
Nausea, abdominal pain and abdominal distension (bloating).
- Common (up to 1 in 10 patients):
Vomiting.
- Unknown (frequency cannot be estimated from the available data):
Severe allergic reaction involving swelling of the face, lips, tongue, or throat, or breathing difficulties (anaphylactic shock), hives, rash, itchy skin (pruritus).

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of KLEARIT

- Storage
Keep out of the reach and sight of children.

Store below 30°C.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like
- KLEARIT is a white powder

- Ingredients

- Active ingredient(s)
 1. Polyethylene Glycol 3350
 2. Anhydrous sodium sulfate
 3. Sodium bicarbonate
 4. Sodium chloride
 5. Potassium chloride

- b. Inactive ingredients
N/A

MAL number(s):

MAL*****

Manufacturer

TIL HEALTHCARE PVT LTD,
Plot No: 100 West, R 5
South Sri city SEZ,
Cherivi village
Irugulam post, Satyavedu
Mandal Tirupati district-
517588
Andhra Pradesh, INDIA

Product Registration Holder

Biocare Pharmaceutical (M) Sdn Bhd
No.28G, 28-1 & 28- 2,
Pusat Perdagangan PJCC,
Jalan PJS 5/28,
46150, Petaling Jaya,
Selangor.

Date of revision

16/10/2025

Serial Number

NPRA (R2) 25/109