

Actual Size Open: (W)160 X (H) 127mm
Close: (W) 63.5 X (H) 40mm

WeKare Vaporub

Active Ingredients:

Camphor 5.26 %w/w, Menthol 2.82 %w/w, Eucalyptus Oil 2.0 %w/w

Product Description:

White colour ointment with characteristic odour.

Indication:

Symptomatic relief of blocked nose, cough, muscular aches and pains due to colds.

Direction of Use:

Adults: Apply WeKare Vaporub on the throat, chest and back when necessary. For deeper relief, add a small amount of WeKare Vaporub to a bowl of hot (not boiling) water and inhale the vapour for 10 – 15 minutes. When using this product, do not apply heat, microwave or add to boiling water.

Children 2 years and above: Apply a small amount of WeKare Vaporub on the chest and back. Rub gently and leave clothes loose for easy inhalation.

Route of Administration:

Topical

Pharmacodynamics:

Menthol, camphor and eucalyptus oil act locally in the respiratory tract.

Pharmacokinetics:

The product exerts its action locally; little or no absorption is expected to occur.

Contraindications:

Contraindicated in children below 2 years of age.

Patient with known hypersensitivity to any of the ingredients in the product.

Warnings and Precautions:

This product is contraindicated in children below 2 years of age. Caution must be exercised when older children are treated. It is dangerous to place any camphor containing product into the nostrils of children.

A small amount applied this way may cause immediate collapse.

Initially apply a small amount to a limited skin area and observe whether there is discomfort or itch before applying to larger skin area. Discontinues use if irritation develops.

Avoid contact with the eyes and mucous membranes. Do not apply to open wounds or damaged skin. Do not apply over large areas of the body or bandage or use with heating pads.

If symptoms persist, consult your doctor.

For external use only. Not to be taken orally. Keep out of reach of children.

Product Registration Holder:

Wen Ken Marketing (M) Sdn Bhd 24,
Jalan Lambak, Taman Johor, 81200
Johor Bahru, Johor, Malaysia.

Manufacturer:

Syarikat Wen Ken Drug Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.

Date of Revision: MAY 2025

Effects on Ability to Drive and Use Machines: Not known

Interactions with Other Medicaments: Not known

Pregnancy and Lactation:

Safety for use in pregnancy has not been established. Please consult your doctor/pharmacist prior to using this product if you are pregnant, planning for pregnancy or breastfeeding. This product should not be used on the mother's chest during breastfeeding.

Adverse Effects / Undesirable Effects:

This product can cause convulsion. Headache, dizziness, nausea and vomiting, skin irritation, contact dermatitis, rash, itching, redness or swelling, burning or stinging sensation may occur.

May cause hypersensitivity/ allergic reactions in some individuals with sensitive skin.

Symptoms and Treatment of Overdose:

Symptoms

Ingestion of significant quantities is reported to cause symptoms: severe abdominal pain, nausea, vomiting, vertigo, ataxia, drowsiness and coma; instant collapse in infants after the local application of menthol or camphor to their nostrils. Seizures may be the first clinical sign of severe toxicity of camphor; however, seizures are usually self-limited. Severe toxicity of camphor can result in delirium, visual hallucinations, cerebral edema and status epilepticus. Systemic toxicity may include hypotension, tachycardia, respiratory failure and death.

Treatment

The stomach should be emptied by gastric lavage or administration of oral activated charcoal. Fluid and electrolyte management is the mainstay of treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalemia and dehydration if present. Any convulsion must be controlled first through supportive care including anticonvulsant therapy.

Storage Conditions: Store below 30 °C. Protect from direct sunlight.

Shelf Life: Please refer to the packaging. Do not use beyond the expiry date.

Dosage Form: Ointment

Packing Sizes: 25g , 45g

The information contained in this leaflet is limited. For further information, please consult your doctor or pharmacist.

40 mm

160 mm

63.5mm

127mm