SHIHAO 施豪
SHIHAO CHLORTETRACYCLINE
15% (W/W) PREMIX GRANULES
Each kg contains 150g Chlortetracycline Hcl
Manufacturing date:
Batch No:
Expiration date:
Country's Registration No:
Net Weight: 25kg
PUCHENG CHIA TAI BIOCHEMISTRY CO., LTD.
GRANULES
GRANULES

正面


SHIHAO 施豪
SHIHAO CHLORTETRACYCLINE
15% (W/W) PREMIX GRANULES
Each kg contains 150g Chlortetracycline Hcl
Indication
Pigs & Chickens
For treatment and control of respiratory and systemic infections associated with organisms sensitive to chlortetracycline.
The recommended dose rates
Pigs:10–20 mg/kg bodyweight daily
Chickens: 20-30 mg/kg bodyweight daily
These dose rates can usually be achieved by mixing 2 kg of Shihao Chlortetracycline 15% Premix Granules per tonne of complete feed to give a concentration of 300 ppm chlortetracycline hydrochloride.
However, the correct incorporation rate should always be calculated based on the feed consumption rates of the animals to be treated.
To ensure thorough dispersion of the product, it should first be mixed with a suitable quantity of feed ingredients before incorporation into the final mix. The medicated feed should be supplied to the affected pen(s) or group(s) of pigs and chickens. Treatment should be continued for a period of five to seven days for respiratory or systemic infections. During the treatment period, only feed medicated with the product should be supplied.
A manufacturer who is approved to incorporate directly at any concentration, veterinary medicinal products or premixtures containing such products must be responsible for mixing when incorporation is less than 2kg per tonne for final feed.
For incorporation into dry feed at the registered mill.
Storage
Protect from strong light. Store at 30°C.
Shelf-life after incorporation into pelleted feed: 24 hours
Warnings and Precautions
Special warnings: None
Special precautions for use
i) Special precautions for use in animals
None
ii)Special precautions to be taken by the person administering the veterinary medicinal product to animals
If you know you are hypersensitive (allergic) to chlortetracycline, do not handle the product.
When handling the product, inhalation of the dust must be avoided by wearing a disposable half-mask respirator. Avoid contact with skin and eyes.
Gloves should be worn whilst handling the product. If contact with the skin or eyes occurs, wash area immediately with clean fresh water.
If irritation persists seek medical attention.
Do not eat, drink or smoke whilst handling the product.
Hands and exposed skin should be washed thoroughly after use.
Withdrawal period
Meat and offal: Pigs: 10 days Chickens: 2 days Eggs:
Not authorized for use in laying birds producing eggs for human consumption. Do not use within 14 days of the onset of laying
Disposal of container
In accordance to local law and regulations
Manufactured by:
PUCHENG CHIA TAI BIOCHEMISTRY CO., LTD.
No.56, Dashixi, Putan Village, Pucheng, Fujian, P.R. China
Marketing authorization holder:
Prima Inter-Chem Sdn. Bhd. 4, Jalan Sungai Kayu Ara 32/38, Berjaya Industrial Park, Seksyen 32, 40460 Shah Alam, Selangor Malaysia Tel: +60357406060 Fax:+60357406061
Keep Medicine Out of Reach of Children
Jauhi Daripada Kanak-kanak
For Animal Use Only
Controlled Medicine
GRANULES
GRANULES

背面