

PACKING INSERT

Registration Number:**CONTROLLED MEDICINE****FOR ANIMAL USE ONLY****KEEP OUT OF REACH OF CHILDREN****Product name:** EsNicarb 25% W/W Premix, Granule**Strength of Active Substance(s):** 100g:25g Nicarbazine**Product Description:** Yellow to yellow-brown Granule**Pharmacodynamics:** Pharmacodynamics:

This product has effect on *Eimeria acervulina*, such as *E. Maxima*, *E. necatrix*, *E. mivata*, *E. tenella*, *E. brunetti*, etc. of chickens. It mainly inhibits the growth and reproduction of chicken coccidiosis on second generation merozoites, and its peak effect is on the 4th day after infection. Mainly used for chicken, turkey coccidiosis.

Pharmacokinetics: This product can be absorbed by the digestive tract and is widely distributed in body tissues and body fluids. The chickens were fed mixed feeding at the recommended dose for 11 days, and the residual drug could still be detected in blood and edible tissues after 2 days of drug withdrawal.

Environmental properties:

Indication: As an aid in preventing outbreaks of cecal (*Eimeria tenella*) and intestinal (*E. acervulina*, *E. maxima*, *E. necatrix*, and *E. brunetti*) coccidiosis in chickens

Target species: Poultry

Recommended Dosage: Thoroughly and evenly mixed in the feed in accordance with current good manufacturing practice for feed. 1ton (2000lb) of finished feed containing 0.01-0.02% nicarbazine (equal to 400g-800g of this finished product)

Route of Administration: Thoroughly and evenly mixed in the feed

Contraindication: Do not use in flushing mashers. Do not use as a treatment for outbreaks of coccidiosis. If losses exceed 0.5 percent in a 2-day period, obtain an accurate diagnosis and follow the instructions of your veterinarian or poultry pathologist.

Warning and Precautions:

Keep this and all drugs out of the reach of children. Not for Human use

Statement on usage during pregnancy and lactation: Do not feed to laying hens in production

Side effect: Broilers fed feed medicated with nicarbazine 25% may show reduced heat tolerance when exposed to high temperature and high humidity to which they have not been accustomed and under severe conditions, fatalities may result. An ample supply of drinking water and adequate ventilation will improve the broilers tolerance of heat.

Incompatibilities (for injections only): No applicable

Withdrawal period(s):

Withdraw 4 days before slaughter when used at levels $\leq 0.0125\%$.

Withdraw 5 days before slaughter when used at levels $>0.0125\%$

Storage conditions: Store in dry place below 25°C, protect from light and moisture.

Packaging Available: 25kg/bag

Name and address of manufacture: Zhejiang Esigma Biotechnology Co., Ltd.

Address: No 3, Chunchao Road, Chang'an Town, Haining, Jiaxing, Zhejiang, China

Product registration holder: Pahang Pharmacy SDN BHD (10826-K)

Address: Lot 5979 Jalan Teratai, 5 Mile Off Jalan Meru, 41050 Klang, Selangor Darul Ehsan, Malaysia

Batch Number:

Manufacturing Date:

Expiration Date:

Date of Revision of Package Insert: 30th Oct, 2023