

SUPAGREL 90 (Ticagrelor Film-Coated Tablets 90 mg)

1. Name of the medicinal product

SUPAGREL 90 (Ticagrelor Film-Coated Tablets 90 mg)

2. Qualitative and quantitative composition

Each film-coated tablet contains 90 mg ticagrelor.

For the full list of excipients, see section 6.1.

3. Pharmaceutical form

Film-coated tablet (tablet).

Yellow coloured, round, biconvex, film coated tablets debossed with “90” & “M” on one side and plain on other side free from physical defects.

4. Clinical particulars

4.1 Therapeutic indications

Acute Coronary Syndrome or a History of Myocardial Infarction

Supagrel 90mg, co-administered with acetylsalicylic acid (ASA), is indicated for the prevention of atherothrombotic events in adult patients with Acute Coronary Syndromes (unstable angina, non ST elevation Myocardial Infarction [NSTEMI] or ST elevation Myocardial Infarction [STEMI]); including patients managed medically, and those who are managed with percutaneous coronary intervention (PCI) or coronary artery by-pass grafting (CABG).

Acute Ischemic Stroke or Transient Ischemic Attack (TIA)

Supagrel 90mg is indicated to reduce the risk of stroke in patients with acute ischemic stroke (NIH Stroke Scale score ≤5) or high-risk transient ischemic attack (TIA).

For further information, please refer to section 5.1.

4.2 Posology and method of administration

Acute Coronary Syndrome or a History of Myocardial Infarction

Supagrel 90mg treatment should be initiated with a single 180 mg loading dose (two tablets of 90 mg) and then continued at 90 mg twice daily.

Patients taking Supagrel 90mg should also take ASA daily, unless specifically contraindicated. Following an initial dose of ASA, Supagrel 90mg should be used with a maintenance dose of ASA of 75-150 mg (see section 5.1).

Treatment is recommended for up to 12 months unless discontinuation of Supagrel 90mg is clinically indicated (see section 5.1). Experience beyond 12 months is limited.

In patients with Acute Coronary Syndromes (ACS), premature discontinuation with any antiplatelet therapy, including Supagrel 90mg, could result in an increased risk of cardiovascular (CV) death, myocardial infarction (MI) or stroke due to the patient's underlying disease. Therefore, premature discontinuation of treatment should be avoided.

In patients having an ACS event, the loading dose of 180 mg should be given as soon as possible regardless of any previous antiplatelet treatment.

Acute Ischemic Stroke or Transient Ischemic Attack (TIA)

Initiate treatment with a 180 mg loading dose of SUPAGREL 90 and then continue with 90 mg twice daily for up to 30 days. The treatment effect accrued early in the course of therapy (see section “Pharmacodynamic properties”).

Use SUPAGREL 90 with a loading dose of ASA (300 to 325 mg) and a daily maintenance dose of ASA of 75 to 100 mg (see “Special Warnings and Precautions For Use” and “Pharmacodynamic properties”).

Physicians who, desire to switch patients, with a prior ACS event, to Supagrel 90mg should administer the first dose of Supagrel 90mg 24 hours following the last dose of the other antiplatelet medication.

Discontinuation of ASA may be considered after 3 months in patients with ACS who have undergone a percutaneous coronary intervention (PCI) procedure and have an increased risk of bleeding. In that case, ticagrelor as single antiplatelet therapy should be continued for 9 months.

Missed dose

Lapses in therapy should also be avoided. A patient who misses a dose of Supagrel 90mg should take only one 90 mg tablet (their next dose) at its scheduled time.

Special populations

Elderly

No dose adjustment is required in elderly (see section “Pharmacokinetic properties”).

Renal impairment

No dose adjustment is necessary for patients with renal impairment (see section “Pharmacokinetic properties”).

Hepatic impairment

Ticagrelor has not been studied in patients with severe hepatic impairment and its use in these patients is therefore contraindicated (see section 4.3). Only limited information is available in patients with moderate hepatic impairment. Dose adjustment is not recommended, but ticagrelor should be used with caution (see sections 4.4 & 5.2). No dose adjustment is necessary for patients with mild hepatic impairment (see section 5.2).

Paediatric patients

The safety and efficacy of ticagrelor in children below the age of 18 in the approved adult indication has not been established. There is no relevant use of ticagrelor in children with sickle cell disease (see sections 5.1 and 5.2).

Method of administration

For oral use. Supagrel 90mg can be taken with or without food. For patients who are unable to swallow the tablet(s) whole, the tablets can be crushed to a fine powder and mixed in half a glass of water and drunk immediately. The glass should be rinsed with a further half glass of water and the contents drunk. The mixture can also be administered via a nasogastric tube (CH8 or greater). It is important to flush the nasogastric tube through with water after administration of the mixture.

4.3 Contraindications

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1 (see section 4.8).
- Active pathological bleeding.
- History of intracranial haemorrhage (see section 4.8).
- Severe hepatic impairment (see sections 4.2, 4.4 and 5.2).
- Co-administration of ticagrelor with strong CYP3A4 inhibitors (e.g. ketoconazole, clarithromycin, nefazodone, ritonavir and atazanavir), as co-administration may lead to a substantial increase in exposure to ticagrelor (see section 4.5).

4.4 Special warnings and precautions for use

Bleeding risk

The use of ticagrelor in patients at known increased risk for bleeding should be balanced against the benefit in terms of prevention of atherothrombotic events (see section “Undesirable effects” and “Pharmacodynamic properties”). If clinically indicated, ticagrelor should be used with caution in the following patient groups:

- Patients with a propensity to bleed (e.g. due to recent trauma, recent surgery, coagulation disorders, active or recent gastrointestinal bleeding) or who are at increased risk of trauma. The use of ticagrelor is contraindicated in patients with active pathological bleeding, in those with a history of intracranial haemorrhage, and in patients with moderate to severe hepatic impairment (see section 4.3).
 - Patients with concomitant administration of medicinal products that may increase the risk of bleeding (e.g. non-steroidal anti-inflammatory drugs (NSAIDs), oral anticoagulants and/or fibrinolytics) within 24 hours of ticagrelor dosing. Platelet transfusion did not reverse the antiplatelet effect of ticagrelor in healthy volunteers and is unlikely to be of clinical benefit in patients with bleeding. Since co-administration of ticagrelor with desmopressin did not decrease template-bleeding time, desmopressin is unlikely to be effective in managing clinical bleeding events (see section “Interaction with other medicinal products and other forms of interaction”).
- Antifibrinolytic therapy (aminocaproic acid or tranexamic acid) and/or recombinant factor VIIa may increase haemostasis. Ticagrelor may be resumed after the cause of bleeding has been identified and controlled.

Patients treated for acute ischemic stroke or TIA

Patients at NIHSS >5 and patients receiving thrombolysis were excluded from THALES and use of SUPAGREL 90 in such patients is not recommended.

Surgery

Patients should be advised to inform physicians and dentists that they are taking ticagrelor before any surgery is scheduled and before any new medicinal product is taken.

In PLATO patients undergoing coronary artery bypass grafting (CABG), ticagrelor had more bleeding than clopidogrel when stopped within 1 day prior to surgery but a similar rate of major bleeds compared to clopidogrel after stopping

therapy 2 or more days before surgery (see section “Undesirable effects”). If a patient is to undergo elective surgery and antiplatelet effect is not desired, ticagrelor should be discontinued 5 days prior to surgery (see section “Pharmacodynamic properties”).

Patients with prior ischaemic stroke

ACS patients with prior ischaemic stroke can be treated with ticagrelor for up to 12 months (PLATO study).

In PEGASUS, patients with history of MI with prior ischaemic stroke were not included. Therefore, in the absence of data, treatment beyond one year is not recommended in these patients.

Hepatic impairment

Use of ticagrelor is contraindicated in patients with severe hepatic impairment (see sections “Posology and method of administration” and “Contraindications”). There is limited experience with ticagrelor in patients with moderate hepatic impairment, therefore, caution is advised in these patients (see sections “Posology and method of administration” and “Pharmacokinetic properties”).

Patients at risk for bradycardic events

Holter ECG monitoring has shown an increased frequency of mostly asymptomatic ventricular pauses during treatment with ticagrelor compared with clopidogrel. Patients with an increased risk of bradycardic events (e.g. patients without a pacemaker who have sick sinus syndrome, 2nd or 3rd degree AV block or bradycardic-related syncope) have been excluded from the main PLATO study evaluating the safety and efficacy of ticagrelor as they may be at increased risk of developing bradyarrhythmias with ticagrelor. Therefore, due to the limited clinical experience, ticagrelor should be used with caution in these patients (see section 5.1).

Bradyarrhythmic events, including 2nd and 3rd degree AV block, have however been reported in the post-marketing setting in patients with or without history of bradyarrhythmia, in most cases, shortly after initiation of treatment with ticagrelor. Therefore, ticagrelor should be used with caution and these patients should be closely monitored during the first few weeks on treatment.

In addition, caution should be exercised when administering ticagrelor concomitantly with medicinal products known to induce bradycardia. However, no evidence of clinically significant adverse reactions was observed in the PLATO trial after concomitant administration with one or more medicinal products known to induce bradycardia (e.g., 96% beta blockers, 33% calcium channel blockers diltiazem and verapamil, and 4% digoxin) (see section “Interaction with other medicinal products and other forms of interaction”).

During the Holter substudy in PLATO, more patients had ventricular pauses >3 seconds with ticagrelor than with clopidogrel during the acute phase of their ACS. The increase in Holter- detected ventricular pauses with ticagrelor was higher in patients with chronic heart failure (CHF) than in the overall study population during the acute phase of ACS, but not at one month with ticagrelor or compared to clopidogrel. There were no adverse clinical consequences associated with this imbalance (including syncope or pacemaker insertion) in this patient population (see section “Pharmacodynamic properties”).

Dyspnoea

Dyspnoea was reported in patients treated with ticagrelor. Dyspnoea is usually mild to moderate in intensity and often resolves without need for treatment discontinuation. Patients with asthma/chronic obstructive pulmonary disease (COPD) may have an increased absolute risk of experiencing dyspnoea with ticagrelor. Ticagrelor should be used with caution in patients with history of asthma and/or COPD. The mechanism has not been elucidated. If a patient reports new, prolonged or worsened dyspnoea this should be investigated fully and if not tolerated, treatment with ticagrelor should be stopped. For further details see section “Undesirable effects”.

Central sleep apnoea

Central sleep apnoea including Cheyne-Stokes respiration has been reported in the post- marketing setting in patients taking ticagrelor. If central sleep apnoea is suspected, further clinical assessment should be considered.

Creatinine elevations

Creatinine levels may increase during treatment with ticagrelor. The mechanism has not been elucidated. Renal function should be checked according to routine medical practice. In patients with ACS, it is recommended that renal function is also checked one month after initiating the treatment with ticagrelor, paying special attention to patients ≥ 75 years, patients with moderate/severe renal impairment and those receiving concomitant treatment with an angiotensin receptor blocker (ARB).

Uric acid increase

Hyperuricaemia may occur during treatment with ticagrelor (see section “Undesirable effects”). Caution is advised in patients with history of hyperuricaemia or gouty arthritis. As a precautionary measure, the use of ticagrelor in patients with uric acid nephropathy is discouraged.

Thrombotic Thrombocytopenic Purpura (TTP)

Thrombotic Thrombocytopenic Purpura (TTP) has been reported very rarely with the use of ticagrelor. It is characterised by thrombocytopenia and microangiopathic haemolytic anaemia associated with either neurological findings, renal dysfunction or fever. TTP is a potentially fatal condition requiring prompt treatment including plasmapheresis.

Interference with platelet function tests to diagnose heparin induced thrombocytopenia (HIT)

In the heparin induced platelet activation (HIPA) test used to diagnose HIT, anti-platelet factor 4/heparin antibodies in patient serum activate platelets of healthy donors in the presence of heparin. False negative results in a platelet function test (to include, but may not be limited to the HIPA test) for HIT have been reported in patients administered ticagrelor. This is related to inhibition of the P2Y₁₂-receptor on the healthy donor platelets in the test by ticagrelor in the patient’s sera/plasma. Information on concomitant treatment with ticagrelor is required for interpretation of HIT platelet function tests.

In patients who have developed HIT, the benefit-risk of continued treatment with ticagrelor should be assessed, taking both the prothrombotic state of HIT and the increased risk of bleeding with concomitant anticoagulant and ticagrelor treatment into consideration.

Other

Based on a relationship observed in PLATO between maintenance ASA dose and relative efficacy of ticagrelor compared to clopidogrel, co-administration of ticagrelor and high maintenance dose ASA (>300 mg) is not recommended (see section “Pharmacodynamic properties”).

Sodium

Supagrel 90mg contains less than 1 mmol sodium (23 mg) per dose, i.e. is essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

Ticagrelor is primarily a CYP3A4 substrate and a mild inhibitor of CYP3A4. Ticagrelor is also a P-glycoprotein (P-gp) substrate and a weak P-gp inhibitor and may increase the exposure of P-gp substrates.

Effects of medicinal and other products on ticagrelor

CYP3A4 inhibitors

- Strong CYP3A4 inhibitors – Co-administration of ketoconazole with ticagrelor increased the ticagrelor C_{max} and AUC equal to 2.4-fold and 7.3-fold, respectively. The C_{max} and AUC of the active metabolite were reduced by 89% and 56%, respectively. Other strong inhibitors of CYP3A4 (clarithromycin, nefazodone, ritonavir and atazanavir) would be expected to have similar effects and therefore concomitant use of strong CYP3A4 inhibitors with ticagrelor is contraindicated (see section 4.3).
- Moderate CYP3A4 inhibitors – Co-administration of diltiazem with ticagrelor increased the ticagrelor C_{max} by 69% and AUC to 2.7-fold and decreased the active metabolite C_{max} by 38% and AUC was unchanged. There was no effect of ticagrelor on diltiazem plasma levels. Other moderate CYP3A4 inhibitors (e.g. amprenavir, aprepitant, erythromycin and fluconazole) would be expected to have a similar effect and can as well be co-administered with ticagrelor.
- A 2-fold increase of ticagrelor exposure was observed after daily consumption of large quantities of grapefruit juice (3 x 200 ml). This magnitude of increased exposure is not expected to be clinically relevant to most patients.

CYP3A inducers

Co-administration of rifampicin with ticagrelor decreased ticagrelor C_{max} and AUC by 73% and 86%, respectively. The C_{max} of the active metabolite was unchanged and the AUC was decreased by 46%, respectively. Other CYP3A inducers (e.g. phenytoin, carbamazepine and phenobarbital) would be expected to decrease the exposure to ticagrelor as well. Co-administration of ticagrelor with potent CYP3A inducers may decrease exposure and efficacy of ticagrelor, therefore, their concomitant use with ticagrelor is discouraged.

Cyclosporine (P-gp and CYP3A inhibitor)

Co-administration of cyclosporine (600 mg) with ticagrelor increased ticagrelor C_{max} and AUC equal to 2.3-fold and 2.8-fold, respectively. The AUC of the active metabolite was increased by 32% and C_{max} was decreased by 15% in the presence of cyclosporine.

No data are available on concomitant use of ticagrelor with other active substances that also are potent P-gp inhibitors and moderate CYP3A4 inhibitors (e.g. verapamil, quinidine) that also may increase ticagrelor exposure. If the association cannot be avoided, their concomitant use should be made with caution.

Others

Clinical pharmacology interaction studies showed that co-administration of ticagrelor with heparin, enoxaparin and ASA or desmopressin did not have any effect on the pharmacokinetics of ticagrelor or the active metabolite or on ADP- induced platelet aggregation compared with ticagrelor alone. If clinically indicated, medicinal products that alter haemostasis should be used with caution in combination with ticagrelor.

A delayed and decreased exposure to oral P2Y₁₂ inhibitors, including ticagrelor and its active metabolite, has been observed in patients with ACS treated with morphine (35% reduction in ticagrelor exposure). This interaction may be related to reduced gastrointestinal motility and apply to other opioids. The clinical relevance is unknown, but data indicate the potential for reduced ticagrelor efficacy in patients co-administered ticagrelor and morphine. In patients with ACS, in whom morphine cannot be withheld and fast P2Y₁₂ inhibition is deemed crucial, the use of a parenteral P2Y₁₂ inhibitor may be considered.

Effects of ticagrelor on other medicinal products

Medicinal products metabolised by CYP3A4

- *Simvastatin* – Co-administration of ticagrelor with simvastatin increased simvastatin C_{max} by 81% and AUC by 56% and increased simvastatin acid C_{max} by 64% and AUC by 52% with some individual increases equal to 2- to 3-fold. Co- administration of ticagrelor with doses of simvastatin exceeding 40 mg daily could cause adverse reactions of simvastatin and should be weighed against potential benefits. There was no effect of simvastatin on ticagrelor plasma levels. Ticagrelor may have similar effect on lovastatin. The concomitant use of ticagrelor with doses of simvastatin or lovastatin greater than 40 mg is not recommended.

- *Atorvastatin* - Co-administration of atorvastatin and ticagrelor increased atorvastatin acid C_{max} by 23% and AUC by 36%. Similar increases in AUC and C_{max} were observed for all atorvastatin acid metabolites. These increases are not considered clinically significant.

- A similar effect on other statins metabolised by CYP3A4 cannot be excluded. Patients in PLATO receiving ticagrelor took a variety of statins, with no concern of an association with statin safety among the 93% of the PLATO cohort taking these medicinal products.

Ticagrelor is a mild CYP3A4 inhibitor. Co-administration of ticagrelor and CYP3A4 substrates with narrow therapeutic indices (i.e. cisapride or ergot alkaloids) is not recommended, as ticagrelor may increase the exposure to these medicinal products.

P-gp substrates (including digoxin, cyclosporine)

Concomitant administration of ticagrelor increased the digoxin C_{max} by 75% and AUC by 28%. The mean trough digoxin levels were increased about 30% with ticagrelor co-administration with some individual maximum increases to 2-fold. In the presence of digoxin, the C_{max} and AUC of ticagrelor and its active metabolite were not affected.

Therefore, appropriate clinical and/or laboratory monitoring is recommended when giving narrow therapeutic index P-gp dependent medicinal products like digoxin concomitantly with ticagrelor.

There was no effect of ticagrelor on cyclosporine blood levels. Effect of ticagrelor on other P-gp substrates has not been studied.

Medicinal products metabolised by CYP2C9

Co-administration of ticagrelor with tolbutamide resulted in no change in the plasma levels of either medicinal product, which suggests that ticagrelor is not a CYP2C9 inhibitor and unlikely to alter the CYP2C9 mediated metabolism of medicinal products like warfarin and tolbutamide.

Oral contraceptives

Co-administration of ticagrelor and levonorgestrel and ethinyl estradiol increased ethinyl estradiol exposure approximately 20% but did not alter the pharmacokinetics of levonorgestrel. No clinically relevant effect on oral contraceptive efficacy is expected when levonorgestrel and ethinyl estradiol are co-administered with ticagrelor.

Medicinal products known to induce bradycardia

Due to observations of mostly asymptomatic ventricular pauses and bradycardia, caution should be exercised when administering ticagrelor concomitantly with medicinal products known to induce bradycardia (see section 4.4).

However, no evidence of clinically significant adverse reactions was observed in the PLATO trial after concomitant

administration with one or more medicinal products known to induce bradycardia (e.g. 96% beta blockers, 33% calcium channel blockers diltiazem and verapamil and 4% digoxin).

Other concomitant therapy

In clinical studies, ticagrelor was commonly administered with ASA, proton pump inhibitors, statins, beta-blockers, angiotensin converting enzyme (ACE) inhibitors and angiotensin receptor blockers as needed for concomitant conditions for long-term and also heparin, low molecular weight heparin and intravenous GpIIb/IIIa inhibitors for short durations (see section 5.1). No evidence of clinically significant adverse interactions with these medicinal products was observed.

Co-administration of ticagrelor with heparin, enoxaparin or desmopressin had no effect on activated partial thromboplastin time (aPTT), activated coagulation time (ACT) or factor Xa assays. However, due to potential pharmacodynamic interactions, caution should be exercised with the concomitant administration of ticagrelor with medicinal products known to alter haemostasis.

Due to reports of cutaneous bleeding abnormalities with SSRIs (e.g. paroxetine, sertraline and citalopram), caution is advised when administering SSRIs with ticagrelor as this may increase the risk of bleeding.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential

Women of childbearing potential should use appropriate contraceptive measures to avoid pregnancy during ticagrelor therapy.

Pregnancy

There are no or limited amount of data from the use of ticagrelor in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3). Ticagrelor is not recommended during pregnancy.

Breast-feeding

Available pharmacodynamic/toxicological data in animals have shown excretion of ticagrelor and its active metabolites in milk (see section 5.3). A risk to newborns/infants cannot be excluded. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from ticagrelor therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

Fertility

Ticagrelor had no effect on male or female fertility in animals.

4.7 Effects on ability to drive and use machines

Ticagrelor has no or negligible influence on the ability to drive and use machines. During treatment with ticagrelor, dizziness and confusion have been reported. Therefore, patients who experience these symptoms should be cautious while driving or using machines.

4.8 Undesirable effects

Summary of safety profile

The safety profile of ticagrelor has been evaluated in a large development programme where more than 58,000 patients and healthy volunteers have received ticagrelor. Data on adverse drug reactions identified in clinical studies or from post-marketing experience with ticagrelor are presented below including information from clinical studies specific to the approved indications (PLATO and THALES) (see section 5.2).

In PLATO, patients on ticagrelor had higher incidence of discontinuation due to adverse events than clopidogrel (7.4% vs 5.4%). In PEGASUS, patients on ticagrelor had a higher incidence of discontinuation due to adverse events compared to ASA therapy alone (16.1% for ticagrelor 60 mg with ASA vs. 8.5% for ASA therapy alone). In THALES, patients on ticagrelor in combination with ASA had a higher incidence of discontinuation of study drug due to adverse events compared to ASA therapy alone (9.7% for ticagrelor 90mg twice daily in combination with ASA vs 7.6% for ASA therapy alone). The most commonly reported adverse drug reactions in patients treated with ticagrelor were bleeding and dyspnoea (see section 4.4).

Tabulated list of adverse reactions

The following adverse reactions have been identified following studies or have been reported in post-marketing experience with ticagrelor (Table 1).

Adverse reactions are listed by MedDRA System Organ Class (SOC). Within each SOC the adverse reactions are ranked by frequency categories. Frequency categories are defined according to the following conventions: Very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), very

rare (<1/10,000), not known (cannot be estimated from the available data).

Table 1 – Adverse reactions by frequency and system organ class (SOC)

SOC	Very Common	Common	Uncommon	Not Known
<i>Neoplasms benign, malignant and unspecified (including cysts and polyps)</i>			Tumour bleedings ^a	
<i>Blood and lymphatic system disorders</i>	Blood disorder bleedings ^b			Thrombotic Thrombocytopenic Purpura ^c
<i>Immune system disorders</i>			Hypersensitivity including angioedema ^c	
<i>Metabolism and nutrition disorders</i>	Hyperuricaemia ^d	Gout/Gouty arthritis		
<i>Psychiatric disorders</i>			Confusion	
<i>Nervous system disorders</i>		Dizziness Syncope, Headache	Intracranial haemorrhage ^m	
<i>Eye disorders</i>			Eye haemorrhage ^c	
<i>Ear and labyrinth disorders</i>		Vertigo	Ear haemorrhage	
<i>Vascular disorders</i>		Hypotension		
<i>Cardiac disorders</i>				Bradycardia ^c AV block (2 nd and 3 rd degree) ^c
<i>Respiratory, thoracic and mediastinal disorders</i>	Dyspnoea ^a	Respiratory system bleedings ^f		
<i>Gastrointestinal disorders</i>		Gastrointestinal haemorrhage ^f Diarrhoea, Nausea, Dyspepsia, Constipation	Retroperitoneal haemorrhage	
<i>Skin and subcutaneous tissue disorders</i>		Subcutaneous or dermal bleeding ^h , Rash, Pruritus		
<i>Musculoskeletal and connective tissue disorders</i>			Muscular bleeding ⁱ	
<i>Renal and urinary disorders</i>		Urinary tract bleedings ^j		
<i>Reproductive system and breast disorders</i>			Reproductive system bleedings ^k	
<i>Investigations</i>		Blood creatinine increased ^d		

<i>Injury, poisoning and procedural complications</i>		Post procedural haemorrhage, Traumatic bleedings ^l		
---	--	---	--	--

AV = atrioventricular

- a e.g., bleeding from bladder cancer, gastric cancer, colon cancer
- b e.g., increased tendency to bruise, spontaneous haematoma, haemorrhagic diathesis
- c Identified in post-marketing experience
- d Frequencies derived from lab observations (Uric acid increases to >upper limit of normal from baseline below or within reference range. Creatinine increases of >50% from baseline.) and not crude adverse event report frequency.
- e e.g., conjunctival, retinal, intraocular bleeding
- f e.g., epistaxis, haemoptysis
- g e.g., gingival bleeding, rectal haemorrhage, gastric ulcer haemorrhage
- h e.g., ecchymosis, skin haemorrhage, petechiae
- i e.g., haemarthrosis, muscle haemorrhage
- j e.g., haematuria, cystitis haemorrhagic
- k e.g., vaginal haemorrhage, haemospermia, postmenopausal haemorrhage
- l e.g., contusion, traumatic haematoma, traumatic haemorrhage
- m i.e spontaneous, procedure related or traumatic intracranial haemorrhage

Description of selected adverse reactions

Bleeding

Bleeding findings in PLATO

Overall outcome of bleeding rates in the PLATO study are shown in Table 2.

Table 2 – Analysis of overall bleeding events, Kaplan-Meier estimates at 12 months (PLATO)

	Ticagrelor 90 mg twice daily N=9235	Clopidogrel N=9186	p-value*
PLATO Total Major	11.6	11.2	0.4336
PLATO Major Fatal/Life-Threatening	5.8	5.8	0.6988
Non-CABG PLATO Major	4.5	3.8	0.0264
Non-Procedural PLATO Major	3.1	2.3	0.0058
PLATO Total Major + Minor	16.1	14.6	0.0084
Non-Procedural PLATO Major + Minor	5.9	4.3	□0.0001
TIMI-defined Major	7.9	7.7	0.5669
TIMI-defined Major + Minor	11.4	10.9	0.3272

Bleeding category definitions:

Major Fatal/Life-threatening Bleed: Clinically apparent with >50 g/l decrease in haemoglobin or ≥4 red cell units transfused; or fatal; or intracranial; or intrapericardial with cardiac tamponade; or with hypovolaemic shock or severe hypotension requiring pressors or surgery.

Major Other: Clinically apparent with 30-50 g/L decrease in haemoglobin or 2-3 red cell units transfused; or significantly disabling.

Minor Bleed: Requires medical intervention to stop or treat bleeding.

TIMI Major Bleed: Clinically apparent with >50 g/L decrease in haemoglobin or intracranial haemorrhage.

TIMI Minor Bleed: Clinically apparent with 30-50 g/L decrease in haemoglobin.

*p-value calculated from Cox proportional hazards model with treatment group as the only explanatory variable.

Ticagrelor and clopidogrel did not differ in rates of PLATO Major Fatal/Life-threatening bleeding, PLATO total Major bleeding, TIMI Major bleeding, or TIMI Minor bleeding (Table 2). However, more PLATO combined Major + Minor bleeding occurred with ticagrelor compared with clopidogrel. Few patients in PLATO had fatal bleeds: 20 (0.2%) for ticagrelor and 23 (0.3%) for clopidogrel (see section “Special warnings and precautions for use”).

Age, sex, weight, race, geographic region, concurrent conditions, concomitant therapy, and medical history, including a previous stroke or transient ischaemic attack, all did not predict either overall or non-procedural PLATO Major bleeding. Thus, no particular group was identified at risk for any subset of bleeding.

CABG-related bleeding: In PLATO, 42% of the 1584 patients (12% of cohort) who underwent coronary artery bypass graft (CABG) surgery had a PLATO Major Fatal/Life-threatening bleeding with no difference between treatment groups. Fatal CABG bleeding occurred in 6 patients in each treatment group (see section “Special warnings and precautions for use”).

Non-CABG related bleeding and non-procedural related bleeding: Ticagrelor and clopidogrel did not differ in non-CABG PLATO-defined Major Fatal/Life-threatening bleeding, but PLATO-defined Total Major, TIMI Major, and TIMI Major + Minor bleeding were more common with ticagrelor. Similarly, when removing all procedure related bleeds, more bleeding occurred with ticagrelor than with clopidogrel (Table 2). Discontinuation of treatment due to non-procedural bleeding was more common for ticagrelor (2.9%) than for clopidogrel (1.2%; p<0.001).

Intracranial bleeding: There were more intracranial non-procedural bleeds with ticagrelor (n=27 bleeds in 26 patients, 0.3%) than with clopidogrel (n=14 bleeds, 0.2%), of which 11 bleeds with ticagrelor and 1 with clopidogrel were fatal. There was no difference in overall fatal bleeds.

Bleeding findings in PEGASUS

Overall outcome of bleeding events in the PEGASUS study are shown in Table 3.

Table 3 – Analysis of overall bleeding events, Kaplan-Meier estimates at 36 months (PEGASUS)

	Ticagrelor 60 mg twice daily + ASA N=6958		ASA alone N=6996	
Safety Endpoints	KM %	Hazard Ratio (95% CI)	KM%	p-value
TIMI-defined bleeding categories				
TIMI Major	2.3	2.32 (1.68, 3.21)	1.1	<0.0001
Fatal	0.3	1.00 (0.44, 2.27)	0.3	1.0000
ICH	0.6	1.33 (0.77, 2.31)	0.5	0.3130
Other TIMI Major	1.6	3.61 (2.31, 5.65)	0.5	<0.0001
TIMI Major or Minor	3.4	2.54 (1.93, 3.35)	1.4	<0.0001
TIMI Major or Minor or Requiring medical attention	16.6	2.64 (2.35, 2.97)	7.0	<0.0001
PLATO-defined bleeding categories				
PLATO Major	3.5	2.57 (1.95, 3.37)	1.4	<0.0001
Fatal/Life-threatening	2.4	2.38 (1.73, 3.26)	1.1	<0.0001
Other PLATO Major	1.1	3.37 (1.95, 5.83)	0.3	<0.0001
PLATO Major or Minor	15.2	2.71 (2.40, 3.08)	6.2	<0.0001

Bleeding category definitions:

TIMI Major: Fatal bleeding, OR any intracranial bleeding, OR clinically overt signs of haemorrhage associated with a drop in haemoglobin (Hgb) of ≥ 50 g/L, or when Hgb is not available, a fall in haematocrit (Hct) of 15%. **Fatal:** A bleeding event that directly led to death within 7 days.

ICH: Intracranial haemorrhage.

Other TIMI Major: Non-fatal non-ICH TIMI Major bleeding.

TIMI Minor: Clinically apparent with 30-50 g/L decrease in haemoglobin.

TIMI Requiring medical attention: Requiring intervention, OR leading to hospitalisation, OR prompting evaluation.

PLATO Major Fatal/life-threatening: Fatal bleeding, OR any intracranial bleeding, OR intrapericardial with cardiac tamponade, OR with hypovolaemic shock or severe hypotension requiring pressors/inotropes or surgery OR clinically apparent with >50 g/L decrease in haemoglobin or ≥ 4 red cell units transfused.

PLATO Major Other: Significantly disabling, OR clinically apparent with 30-50 g/L decrease in haemoglobin, OR 2-3 red cell units transfused.

PLATO Minor: Requires medical intervention to stop or treat bleeding.

In PEGASUS, TIMI Major bleeding for ticagrelor 60 mg twice daily was higher than for ASA alone. No increased bleeding risk was seen for fatal bleeding and only a minor increase was observed in intracranial haemorrhages, as compared to ASA therapy alone. There were few fatal bleeding events in the study, 11 (0.3%) for ticagrelor 60 mg and 12 (0.3%) for ASA therapy alone. The observed increased risk of TIMI Major bleeding with ticagrelor 60 mg was primarily due to a higher frequency of Other TIMI Major bleedings driven by events in the gastrointestinal SOC.

Increased bleeding patterns similar to TIMI Major were seen for TIMI Major or Minor and PLATO Major and PLATO Major or Minor bleeding categories (see Table 3). Discontinuation of treatment due to bleeding was more common with ticagrelor 60 mg compared to ASA therapy alone (6.2% and 1.5%, respectively). The majority of these bleedings were of less severity (classified as TIMI Requiring medical attention), e.g. epistaxis, bruising and haematomas.

The bleeding profile of ticagrelor 60 mg was consistent across multiple pre-defined subgroups (e.g. by age, gender, weight, race, geographic region, concurrent conditions, concomitant therapy and medical history) for TIMI Major, TIMI Major or Minor and PLATO Major bleeding events.

Intracranial bleeding: Spontaneous ICHs were reported in similar rates for ticagrelor 60 mg and ASA therapy alone (n=13, 0.2% in both treatment groups). Traumatic and procedural ICHs showed a minor increase with ticagrelor 60 mg treatment, (n=15, 0.2%) compared with ASA therapy alone (n=10, 0.1%). There were 6 fatal ICHs with ticagrelor 60 mg and 5 fatal ICHs with ASA therapy alone. The incidence of intracranial bleeding was low in both treatment groups given the significant comorbidity and CV risk factors of the population under study.

Bleeding findings in THALES

Overall outcome of bleeding events in the THALES study are shown in Table 4.

Table 4 - Analysis of bleeding events, Kaplan-Meier estimate of bleeding rates by treatment at 30 days (THALES)^a

	Ticagrelor 90 mg twice daily with ASA N=5523			ASA alone N=5493		
Safety Endpoints	Patients with events	KM%	Hazard Ratio (95% CI)	Patients with events	KM%	p-value
GUSTO-defined bleeding categories						
GUSTO Severe	28 (0.5%)	0.5%	3.99 (1.74, 9.14)	7 (0.1%)	0.1%	0.001
GUSTO Severe or Moderate	36 (0.7%)	0.6%	3.27 (1.67, 6.43)	11 (0.2%)	0.2%	<.001

Bleeding category definitions:

GUSTO Severe: Any one of the following: fatal bleeding, intracranial bleeding (excluding asymptomatic haemorrhagic transformations of ischaemic brain infarctions and excluding microhaemorrhages <10 mm evident only on gradient-echo magnetic resonance imaging), bleeding that caused haemodynamic compromise requiring intervention (e.g.

systolic blood pressure <90 mm Hg that required blood or fluid replacement, or vasopressor/inotropic support, or surgical intervention).

GUSTO Moderate: Bleeding requiring transfusion of whole blood or packed red blood cells without haemodynamic compromise (as defined above).

^a Full analysis set – intention-to-treat analysis: included all patients randomised irrespective of their protocol adherence and continued participation in the study.

In THALES, the rate of GUSTO Severe bleeding for SUPAGREL 90 90 mg twice daily in combination with ASA was higher than for ASA alone. A similar bleeding pattern was observed for the GUSTO Severe or Moderate bleeding category (see Table 4). Due to the low number of GUSTO Severe bleeding events, no conclusion can be drawn regarding bleeding risk across subgroups. Discontinuation of treatment due to bleeding was more common with SUPAGREL 90 90 mg with ASA compared to ASA therapy alone (2.9% and 0.6%, respectively).

Intracranial bleeding and fatal bleeding: In total, there were 21 intracranial haemorrhages (ICHs) (19 spontaneous, 1 traumatic, 1 procedural) for SUPAGREL 90 90 mg with ASA and 6 ICHs (3 spontaneous, 2 traumatic, 1 procedural) for ASA alone. Fatal bleedings occurred in 11 patients (10 fatal ICHs, 1 fatal gastro-intestinal bleed) for SUPAGREL 90 90 mg with ASA and in 2 patients (2 fatal ICHs) for ASA alone.

Dyspnoea

Dyspnoea, a sensation of breathlessness, is reported by patients treated with ticagrelor. In PLATO, dyspnoea adverse events (AEs) (dyspnoea, dyspnoea at rest, dyspnoea exertional, dyspnoea paroxysmal nocturnal and nocturnal dyspnoea), when combined, was reported by 13.8% of patients treated with ticagrelor and by 7.8% of patients treated with clopidogrel. In 2.2% of patients taking ticagrelor and by 0.6% taking clopidogrel investigators considered the dyspnoea causally related to treatment in the PLATO study and few were serious (0.14% ticagrelor; 0.02% clopidogrel), (see section “Special warnings and precautions for use”). Most reported symptoms of dyspnoea were mild to moderate in intensity, and most were reported as a single episode early after starting treatment.

Compared with clopidogrel, patients with asthma/COPD treated with ticagrelor may have an increased risk of experiencing non-serious dyspnoea (3.29% ticagrelor versus 0.53% clopidogrel) and serious dyspnoea (0.38% ticagrelor versus 0.00% clopidogrel). In absolute terms, this risk was higher than in the overall PLATO population. Ticagrelor should be used with caution in patients with history of asthma and/or COPD (see section “Special warnings and precautions for use”).

About 30% of episodes resolved within 7 days. PLATO included patients with baseline congestive heart failure, chronic obstructive pulmonary disease, or asthma; these patients, and the elderly, were more likely to report dyspnoea. For ticagrelor, 0.9% of patients discontinued study drug because of dyspnoea compared with 0.1% taking clopidogrel. The higher incidence of dyspnoea with ticagrelor is not associated with new or worsening heart or lung disease (see section “Special warnings and precautions for use”). Ticagrelor does not affect tests of pulmonary function.

In PEGASUS dyspnoea was reported in 14.2% of patients taking ticagrelor 60 mg twice daily and in 5.5% of patients taking ASA alone. As in PLATO, most reported dyspnoea was mild to moderate in intensity (see section “Special warnings and precautions for use”). Patients who reported dyspnoea tended to be older and more frequently had dyspnoea, COPD or asthma at baseline.

In THALES, dyspnoea led to study drug discontinuation in 1.0% and 0.2% of patients taking ticagrelor 90 mg in combination with ASA vs. ASA alone, respectively.

Investigations

Uric acid elevations: In PLATO, serum uric acid concentration increased to more than upper limit of normal in 22% of patients receiving ticagrelor compared to 13% of patients receiving clopidogrel. The corresponding numbers in PEGASUS were 9.1%, 8.8% and 5.5% for ticagrelor 90 mg, 60 mg and placebo, respectively. Mean serum uric acid concentration increased approximately 15% with ticagrelor compared to approximately 7.5% with clopidogrel and after treatment was stopped, decreased to approximately 7% on ticagrelor but with no decrease observed for clopidogrel. In PEGASUS, a reversible increase in mean serum uric acid levels of 6.3% and 5.6% was found for ticagrelor 90 mg and 60 mg, respectively, compared to a 1.5% decrease in the placebo group. In PLATO, the frequency of gouty arthritis was 0.2% for ticagrelor vs. 0.1% for clopidogrel. The corresponding numbers for gout/gouty arthritis in PEGASUS were 1.6%, 1.5% and 1.1% for ticagrelor 90 mg, 60 mg and placebo, respectively.

4.9 Overdose

Ticagrelor is well tolerated in single doses up to 900 mg. Gastrointestinal toxicity was dose-limiting in a single

ascending dose study. Other clinically meaningful adverse reactions which may occur with overdose include dyspnoea and ventricular pauses (see section “Undesirable effects”).

In the event of overdose, observe for these potential adverse reactions and consider ECG monitoring.

There is currently no known antidote to reverse the effects of ticagrelor, and ticagrelor is not dialysable (see section “Special warnings and precautions for use”). Treatment of overdose should follow local standard medical practice. The expected effect of excessive ticagrelor dosing is prolonged duration of bleeding risk associated with platelet inhibition. Platelet transfusion is unlikely to be of clinical benefit in patients with bleeding (see section “Special warnings and precautions for use”). If bleeding occurs appropriate supportive measures should be taken.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Platelet aggregation inhibitors excluding heparin, ATC code: B01AC24

Mechanism of action

SUPAGREL 90 contains ticagrelor, a member of the chemical class cyclopentyltriazolopyrimidines (CPTP), which is an oral, direct acting, selective and reversibly binding P2Y₁₂ receptor antagonist that prevents ADP-mediated P2Y₁₂ dependent platelet activation and aggregation. Ticagrelor does not prevent ADP binding, but when bound to the P2Y₁₂ receptor prevents ADP-induced signal transduction. Since platelets participate in the initiation and/or evolution of thrombotic complications of atherosclerotic disease, inhibition of platelet function has been shown to reduce the risk of CV events such as death, MI or stroke.

Ticagrelor also increases local endogenous adenosine levels by inhibiting equilibrative nucleoside transporter-1 (ENT-1).

Ticagrelor has been documented to augment the following adenosine-induced effects in healthy subjects and in patients with ACS: vasodilation (measured by coronary blood flow increases in healthy volunteers and ACS patients; headache), inhibition of platelet function (in human whole blood *in vitro*) and dyspnoea. However, a link between the observed increases in adenosine and clinical outcomes (e.g. morbidity mortality) has not been clearly elucidated.

Pharmacodynamic effects

Onset of action

In patients with stable coronary artery disease (CAD) on ASA, ticagrelor demonstrates a rapid onset of pharmacological effect as demonstrated by a mean inhibition of platelet aggregation (IPA) for ticagrelor at 0.5 hours after 180 mg loading dose of about 41%, with the maximum IPA effect of 89% by 2-4 hours post dose, and maintained between 2-8 hours. 90% of patients had final extent IPA >70% by 2 hours post dose.

Offset of action

If a CABG procedure is planned, ticagrelor bleeding risk is increased compared to clopidogrel when discontinued within less than 96 hours prior to procedure.

Switching data

Switching from clopidogrel 75 mg to ticagrelor 90 mg twice daily results in an absolute IPA increase of 26.4% and switching from ticagrelor to clopidogrel results in an absolute IPA decrease of 24.5%. Patients can be switched from clopidogrel to ticagrelor without any interruption of antiplatelet effect (see section “Posology and method of administration”).

Clinical efficacy and safety

PLATO study (Acute Coronary Syndromes)

The PLATO study included 18,624 patients who presented within 24 hours of onset of symptoms of unstable angina (UA), non-ST elevation myocardial infarction (NSTEMI) or ST elevation myocardial infarction (STEMI), and were initially managed medically, or with percutaneous coronary intervention (PCI), or with CABG.

Clinical efficacy

On a background of daily ASA, ticagrelor 90 mg twice daily showed superiority to 75 mg daily clopidogrel in preventing the composite endpoint of CV death, MI or stroke, with the difference driven by CV death and MI. Patients received a 300 mg loading dose of clopidogrel (600 mg possible if having PCI) or 180 mg of ticagrelor.


The result appeared early (absolute risk reduction [ARR] 0.6% and Relative Risk Reduction [RRR] of 12% at 30 days), with a constant treatment effect over the entire 12-month period, yielding ARR 1.9% per year with RRR of 16%. This suggests it is appropriate to treat patients with ticagrelor for up to 12 months (see section “Posology and method of administration”). Treating 54 ACS patients with ticagrelor instead of clopidogrel will prevent 1

atherothrombotic event; treating 91 will prevent 1 CV death (see Figure 1 and Table 4).

The treatment effect of ticagrelor over clopidogrel appears consistent across many subgroups, including weight; sex; medical history of diabetes mellitus, transient ischaemic attack or non- haemorrhagic stroke, or revascularisation; concomitant therapies including heparins, GpIIb/IIIa inhibitors and proton pump inhibitors (see section “Interaction with other medicinal products and other forms of interaction”); final index event diagnosis (STEMI, NSTEMI or UA); and treatment pathway intended at randomisation (invasive or medical).

A weakly significant treatment interaction was observed with region whereby the hazard ratio (HR) for the primary endpoint favours ticagrelor in the rest of world but favours clopidogrel in North America, which represented approximately 10% of the overall population studied (interaction p-value=0.045). Exploratory analyses suggest a possible association with ASA dose such that reduced efficacy was observed with ticagrelor with increasing ASA doses. Chronic daily ASA doses to accompany ticagrelor should be 75-150 mg (see section “Posology and method of administration” and “Special warnings and precautions for use”).

Figure 1 – Analysis of primary clinical composite endpoint of CV death, MI and stroke (PLATO)



Ticagrelor reduced the occurrence of the primary composite endpoint compared to clopidogrel in both the UA/NSTEMI and STEMI population (Table 5). Thus, Supagrel 90 mg twice daily together with low-dose ASA can be used in patients with ACS (unstable angina, non-ST elevation Myocardial Infarction [NSTEMI] or ST elevation Myocardial Infarction [STEMI]); including patients managed medically, and those who are managed with percutaneous coronary intervention (PCI) or coronary artery by-pass grafting (CABG).

Table 5 - Analysis of primary and secondary efficacy endpoints (PLATO)

	Ticagrelor 90 mg twice daily (% patients with	Clopidogrel 75 mg once daily (% patients with	AR R ^a (%/yr)	RRR ^a (%) (95% CI)	p-value

	event) N=9333	event) N=9291			
CV death, MI (excl. silent MI) or stroke	9.3	10.9	1.9	16 (8, 23)	0.0003
Invasive intent	8.5	10.0	1.7	16 (6, 25)	0.0025
Medical intent	11.3	13.2	2.3	15 (0.3, 27)	0.0444 ^d
CV death	3.8	4.8	1.1	21 (9, 31)	0.0013
MI (excl. silent MI) ^b	5.4	6.4	1.1	16 (5, 25)	0.0045
Stroke	1.3	1.1	-0.2	-17 (-52, 9)	0.2249
All-cause mortality, MI (excl. silent MI) or stroke	9.7	11.5	2.1	16 (8, 23)	0.0001
CV death, total MI, stroke, SRI, RI, TIA or other ATE ^c	13.8	15.7	2.1	12 (5, 19)	0.0006
All-cause mortality	4.3	5.4	1.4	22 (11, 31)	0.0003 ^d
Definite stent thrombosis	1.2	1.7	0.6	32 (8, 49)	0.0123 ^d

^a ARR = absolute risk reduction; RRR = relative risk reduction = (1-Hazard ratio) x 100%. A negative RRR indicates a relative risk increase.

^b Excluding silent MI.

^c SRI = serious recurrent ischaemia; RI = recurrent ischaemia; TIA = transient ischaemic attack; ATE = arterial thrombotic event. Total MI includes silent MI, with date of event set to date when discovered.

^d Nominal significance value; all others are formally statistically significant by pre-defined hierarchical testing.

PLATO genetic substudy

CYP2C19 and ABCB1 genotyping of 10,285 patients in PLATO provided associations of genotype groups with PLATO outcomes. The superiority of ticagrelor over clopidogrel in reducing major CV events was not significantly affected by patient CYP2C19 or ABCB1 genotype. Similar to the overall PLATO study, total PLATO Major bleeding did not differ between ticagrelor and clopidogrel, regardless of CYP2C19 or ABCB1 genotype. Non-CABG PLATO Major bleeding was increased with ticagrelor compared clopidogrel in patients with one or more CYP2C19 loss of function alleles, but similar to clopidogrel in patients with no loss of function allele.

Combined efficacy and safety composite

A combined efficacy and safety composite (CV death, MI, stroke or PLATO-defined 'Total Major' bleeding) indicates that the benefit in efficacy of ticagrelor compared to clopidogrel is not offset by the major bleeding events (ARR 1.4%, RRR 8%, HR 0.92; p=0.0257) over 12 months after ACS.

Clinical safety

Holter substudy:

To study the occurrence of ventricular pauses and other arrhythmic episodes during PLATO, investigators performed Holter monitoring in a subset of nearly 3000 patients, of whom approximately 2000 had recordings both in the acute phase of their ACS and after one month. The primary variable of interest was the occurrence of ventricular pauses ≥ 3

seconds. More patients had ventricular pauses with ticagrelor (6.0%) than with clopidogrel (3.5%) in the acute phase; and 2.2% and 1.6%, respectively, after 1 month (see section 4.4). The increase in ventricular pauses in the acute phase of ACS was more pronounced in ticagrelor patients with history of CHF (9.2% versus 5.4% in patients without CHF history; for clopidogrel patients, 4.0% in those with versus 3.6% in those without CHF history). This imbalance did not occur at one month: 2.0% versus 2.1% for ticagrelor patients with and without CHF history, respectively; and 3.8% versus 1.4% with clopidogrel. There were no adverse clinical consequences associated with this imbalance (including pacemaker insertions) in this population of patients.

THALES study (Acute Ischemic Stroke or Transient Ischemic Attack (TIA))

The THALES study (NCT03354429) was a 11016-patient, randomized, double-blind, parallel-group study of SUPAGREL 90 90 mg twice daily versus placebo in patients with acute ischemic stroke or transient ischemic attack (TIA). The primary endpoint was the first occurrence of the composite of stroke and death up to 30 days. Ischemic stroke was assessed as one of the secondary endpoints.

Patients were eligible to participate if they were ≥40 years old, with non-cardioembolic acute ischemic stroke (NIHSS score ≤5) or high-risk TIA (defined as ABCD² score ≥6 or ipsilateral atherosclerotic stenosis ≥50% in the internal carotid or an intracranial artery). Patients who received thrombolysis or thrombectomy within 24 hours prior to randomization were not eligible.

Patients were randomized within 24 hours of onset of an acute ischemic stroke or TIA to receive 30 days of either SUPAGREL 90 (90 mg twice daily, with an initial loading dose of 180 mg) or placebo, on a background of aspirin initially 300-325 mg then 75-100 mg daily. The median treatment duration was 31 days.

SUPAGREL 90 was superior to placebo in reducing the rate of the primary endpoint (composite of stroke and death), corresponding to a relative risk reduction (RRR) of 17% and an absolute risk reduction (ARR) of 1.1% (Table 6). The effect was driven primarily by a significant reduction in the stroke component of the primary endpoint (19% RRR, 1.1% ARR).

Table 6 - Incidences of the primary composite endpoint, primary composite endpoint components, and secondary endpoint (THALES)


	SUPAGREL 90 N=5523		Placebo N=5493		HR (95% CI)	p-value
	n (patients with event)	KM%	n (patients with event)	KM%		
Time to first Stroke or Death	303	5.4%	362	6.5%	0.83 (0.71, 0.96)	0.015
Time to first Stroke*	284	5.1%	347	6.3%	0.81 (0.69, 0.95)	
Time to Death*	36	0.6%	27	0.5%	1.33 (0.81, 2.19)	
Secondary Endpoint						
Time to first Ischemic Stroke	276	5.0%	345	6.2%	0.79 (0.68, 0.93)	0.004

CI = Confidence interval; HR = Hazard ratio; KM = Kaplan-Meier percentage calculated at 30 days; N = Number of patients

*The number of patients with the event of interest. In the time to first stroke, patients who died are censored at the time of death.


The Kaplan-Meier curve (Figure 2) shows the time to first occurrence of the primary composite endpoint of stroke and death.

Figure 2 – Time to First Occurrence of Stroke or Death (THALES)



KM%: Kaplan-Meier percentage evaluated at Day 30; T=Ticagrelor; P=placebo; N=Number of patients
 SUPAGREL 90's treatment effect on stroke and on death accrued over the first 10 days and was sustained at 30 days. Although not studied, this suggests that shorter treatment could result in similar benefit and reduced bleeding risk. The treatment effect of SUPAGREL 90 was generally consistent across pre-defined subgroups (Figure 3).

Figure 3 – Subgroup analyses of ticagrelor 90 mg (THALES)



Note: The figure above presents effects in various subgroups all of which are baseline characteristics and were pre-specified. The 95% confidence limits that are shown do not take into account how many comparisons were made, nor do they reflect the effect of a particular factor after adjustment for all other factors. Apparent homogeneity or heterogeneity among groups should not be over-interpreted.

At Day 30, there was an absolute reduction of 1.2% (95% CI: -2.1%, -0.3%) in the incidence of non-hemorrhagic stroke and death (excluding fatal bleed) favoring ticagrelor (294 events):

5.3%) over placebo (359 events: 6.5%) in the intention-to-treat population. In the same population, there was an absolute increase of 0.4% (95% CI: 0.2%, 0.6%) in the incidence of GUSTO severe bleeding unfavorable to ticagrelor arm (28 events: 0.5%) compared to the placebo arm (7 events: 0.1%).

Paediatric population

In a randomised, double-blind, parallel-group Phase III study (HESTIA 3), 193 paediatric patients (ages 2 to less than 18 years) with sickle cell disease were randomised to receive either placebo or ticagrelor at doses of 15 mg to 45 mg twice daily depending on body weight. Ticagrelor resulted in a median platelet inhibition of 35% at pre-dose and 56% at 2 hours post-dose at steady state.

Compared to placebo, there was no treatment benefit of ticagrelor on the rate of vaso-occlusive crises.

5.2 Pharmacokinetic properties

Ticagrelor demonstrates linear pharmacokinetics and exposure to ticagrelor and the active metabolite (AR-C124910XX) are approximately dose proportional up to 1260 mg.

Absorption

Absorption of ticagrelor is rapid with a median t_{max} of approximately 1.5 hours. The formation of the major circulating metabolite AR-C124910XX (also active) from ticagrelor is rapid with a median t_{max} of approximately 2.5 hours. Following an oral ticagrelor 90 mg single dose under fasted conditions in healthy subjects, C_{max} is 529 ng/ml and AUC is 3451 ng*h/ml. The metabolite parent ratios are 0.28 for C_{max} and 0.42 for AUC. The pharmacokinetics of ticagrelor and AR-C124910XX in patients with a history of MI were generally similar to that in the ACS population. Based on a population pharmacokinetic analysis of the PEGASUS study the median ticagrelor C_{max} was 391 ng/ml and AUC was 3801 ng*h/ml at steady state for ticagrelor 60 mg. For ticagrelor 90 mg C_{max} was 627 ng/ml and AUC was 6255 ng*h/ml at steady state.

The mean absolute bioavailability of ticagrelor was estimated to be 36%. Ingestion of a high-fat meal resulted in a 21% increase in ticagrelor AUC and 22% decrease in the active metabolite C_{max} but had no effect on ticagrelor C_{max} or the AUC of the active metabolite. These small changes are considered of minimal clinical significance; therefore, ticagrelor can be given with or without food. Ticagrelor as well as the active metabolite are P-gp substrates.

Ticagrelor as crushed tablets mixed in water, given orally or administered through a nasogastric tube into the stomach, has a comparable bioavailability to whole tablets with regards to AUC and C_{max} for ticagrelor and the active metabolite. Initial exposure (0.5 and 1 hour post-dose) from crushed ticagrelor tablets mixed in water was higher compared to whole tablets, with a generally identical concentration profile thereafter (2 to 48 hours).

Distribution

The steady state volume of distribution of ticagrelor is 87.5 l. Ticagrelor and the active metabolite is extensively bound to human plasma protein (>99.0%).

Biotransformation

CYP3A4 is the major enzyme responsible for ticagrelor metabolism and the formation of the active metabolite and their interactions with other CYP3A substrates ranges from activation through to inhibition.

The major metabolite of ticagrelor is AR-C124910XX, which is also active as assessed by *in vitro* binding to the platelet P2Y₁₂ ADP-receptor. The systemic exposure to the active metabolite is approximately 30-40% of that obtained for ticagrelor.

Elimination

The primary route of ticagrelor elimination is via hepatic metabolism. When radiolabelled ticagrelor is administered, the mean recovery of radioactivity is approximately 84% (57.8% in faeces, 26.5% in urine). Recoveries of ticagrelor and the active metabolite in urine were both less than 1% of the dose. The primary route of elimination for the active metabolite is most likely via biliary secretion. The mean $t_{1/2}$ was approximately 7 hours for ticagrelor and 8.5 hours for the active metabolite.

Special populations

Elderly

Higher exposures to ticagrelor (approximately 25% for both C_{max} and AUC) and the active metabolite were observed in elderly (≥ 75 years) ACS patients compared to younger patients by the population pharmacokinetic analysis. These differences are not considered clinically significant (see section 4.2).

Paediatric population

Limited data are available in children with sickle cell disease (see sections 4.2 and 5.1).

In the HESTIA 3 study, patients aged 2 to less than 18 years weighing ≥ 12 to ≤ 24 kg, > 24 to ≤ 48 kg and > 48 kg, were administered ticagrelor as paediatric dispersible 15 mg tablets at doses of respectively 15, 30 and 45 mg twice daily. Based on population pharmacokinetic analysis, the mean AUC ranged from 1095 ng*h/mL to 1458 ng*h/mL and the mean C_{max} ranged from 143 ng/mL to 206 ng/mL at steady state.

Gender

Higher exposures to ticagrelor and the active metabolite were observed in women compared to men. These differences are not considered clinically significant.

Renal impairment

Exposure to ticagrelor was approximately 20% lower and exposure to the active metabolite was approximately 17% higher in patients with severe renal impairment (creatinine clearance <30 ml/min) compared to subjects with normal renal function.

In patients with end stage renal disease on haemodialysis AUC and C_{max} of ticagrelor 90 mg administered on a day without dialysis were 38% and 51% higher compared to subjects with normal renal function. A similar increase in exposure was observed when ticagrelor was administered immediately prior to dialysis (49% and 61%, respectively) showing that ticagrelor is not dialysable. Exposure of the active metabolite increased to a lesser extent (AUC 13-14% and C_{max} 17-36%). The inhibition of platelet aggregation (IPA) effect of ticagrelor was independent of dialysis in patients with end stage renal disease and similar to subjects with normal renal function (see section 4.2).

Hepatic impairment

C_{max} and AUC for ticagrelor were 12% and 23% higher in patients with mild hepatic impairment compared to matched healthy subjects, respectively, however, the IPA effect of ticagrelor was similar between the two groups. No dose adjustment is needed for patients with mild hepatic impairment. Ticagrelor has not been studied in patients with severe hepatic impairment and there is no pharmacokinetic information in patients with moderate hepatic impairment. In patients that had moderate or severe elevation in one or more liver function tests at baseline, ticagrelor plasma concentrations were on average similar or slightly higher as compared to those without baseline elevations. No dose adjustment is recommended in patients with moderate hepatic impairment (see sections 4.2 and 4.4).

Ethnicity

Patients of Asian descent have a 39% higher mean bioavailability compared to Caucasian patients. Patients self-identified as black had an 18% lower bioavailability of ticagrelor compared to Caucasian patients, in clinical pharmacology studies, the exposure (C_{max} and AUC) to ticagrelor in Japanese subjects was approximately 40% (20% after adjusting for body weight) higher compared to that in Caucasians. The exposure in patients self-identified as Hispanic or Latino was similar to that in Caucasians.

PRECLINICAL SAFETY DATA

Preclinical data for ticagrelor and its major metabolite have not demonstrated unacceptable risk for adverse effects for humans based on conventional studies of safety pharmacology, single and repeated dose toxicity and genotoxic potential.

Gastrointestinal irritation was observed in several animal species at clinically relevant exposure levels (see section “Undesirable effects”).

In female rats, ticagrelor at high dose showed an increased incidence of uterine tumours (adenocarcinomas) and an increased incidence of hepatic adenomas. The mechanism for uterine tumours is likely hormonal imbalance which can lead to tumours in rats. The mechanism for the hepatic adenomas is likely due to a rodent-specific enzyme induction in the liver. Thus, the carcinogenicity findings are considered unlikely to be relevant for humans.

In rats, minor developmental anomalies were seen at a maternal toxic dose (safety margin of 5.1). In rabbits a slight delay in hepatic maturity and skeletal development was seen in foetuses from dams at high dose without showing maternal toxicity (safety margin of 4.5).

Studies in rats and rabbits have shown reproductive toxicity, with slightly reduced maternal body weight gain and reduced neonatal viability and birth weight, with delayed growth. Ticagrelor produced irregular cycles (mostly extended cycles) in female rats, but did not affect overall fertility in male and female rats. Pharmacokinetic studies performed with radio-labelled ticagrelor have shown that the parent compound and its metabolites are excreted in the milk of rats (see section “Fertility, pregnancy and lactation”).

6. Pharmaceutical particulars

6.1 List of excipients

Tablet core

Mannitol

L-Hydroxy Propyl Cellulose

Calcium hydrogen phosphate dihydrate

Magnesium stearate (E470b)

Hydroxypropylcellulose

Tablet coating

Talc

Titanium dioxide (E171)

Iron oxide yellow (E172)

Macrogol

Hypromellose (E464)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

Store below 30°C.

6.5 Nature and contents of container

- PVC/PVDC Alu Blister Pack of 10 tablets; cartons of 30 tablets (3 blisters) or 60 tablets (6 blisters).
- PVC/PVDC Alu Blister Pack of 14 tablets; cartons of 14 tablets (1 blister) or 56 tablets (4 blisters)

6.6 Special precautions for disposal and other handling

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. Manufacturer

MSN Laboratories Private Limited,

Formulation Division, Unit-II, Sy.no. 1277, 1319 to 1324,

Nandigama (Village & Mandal), Rangareddy (District),

Telangana - 509 228, India.

8. Product Registration Holder

Synercam (M) Sdn Bhd,
A-17-7, Block A, Jaya One, No. 72A,
Jalan Profesor Diraja Ungku Aziz,
Seksyen 13, 46200, Petaling Jaya
Selangor.

9. Date of revision of the text

27th March 2026

Font : Times New Roman, Size: 10