

PRESMEDIC AMBROXOL TABLET 30 MG

Ambroxol hydrochloride (30mg)

What is in this leaflet

1. What Presmedic Ambroxol Tablet 30mg is used for
2. How Presmedic Ambroxol Tablet 30mg works
3. Before you use Presmedic Ambroxol Tablet 30mg
4. How to use Presmedic Ambroxol Tablet 30mg
5. While you are using Presmedic Ambroxol Tablet 30mg
6. Side effects
7. Storage and Disposal of Presmedic Ambroxol Tablet 30mg
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial number

What Presmedic Ambroxol Tablet 30mg is used for

Presmedic Ambroxol Tablet 30mg is used in the treatment of respiratory tract disorders associated with viscid mucus.

How Presmedic Ambroxol Tablet 30mg works

Presmedic Ambroxol Tablet 30mg contains ambroxol, which helps to dissolve thick mucus, therefore improves breathing and reduces cough.

Before you use Presmedic Ambroxol Tablet 30mg

-When you must not use it

If you are allergic towards Ambroxol Hydrochloride or any other ingredients listed in this product.

If you are not sure whether you should start taking this medicine, talk to your pharmacist or doctor.

-Before you start to use it

Tell your doctor or pharmacist if you are allergic to any other medicines or you have any other health-related problems including:

- Patients with liver and kidney problems.
- Special care is recommended for patients who are in the first trimester of pregnancy.
- Allergic towards ambroxol.

- Patients with antibiotic treatment as this medicine may increase the amount of antibiotic penetration.
- If symptoms persist more than 14 days, please seek medical advice.
- Very rare cases of chronically associated severe skin impairments have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as a precaution.

-Taking other medicines

Seek for your doctor or pharmacist advice if you intend to use Presmedic Ambroxol Tablet 30mg together with other medication. You must also inform them about all the medicines you are taking, including those without a prescription.

Presmedic Ambroxol Tablet 30mg administered together with antibiotics such as amoxicillin, cefuroxime, erythromycin, doxycycline will lead to higher antibiotic concentration in the lung tissue.

How to use Presmedic Ambroxol Tablet 30mg

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

-How much to use

For adults: 1 tablet.

-When to use it

3 times daily, after meals.

-How long to use it

Presmedic Ambroxol Tablet 30mg should be taken as your doctor advised you to do so. Do not stop taking this medicine without talking to the doctor.

-If you forget to use it

If you forget to take your dose, take it as soon as you remember. Skip the missed dose if it is almost time for your next dose. Do not double up on the next dose to make up the forgotten dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using Presmedic Ambroxol Tablet 30mg

-Things you must do

Use your medicine exactly as your doctor has told you,

Tell all the doctors, dentists and pharmacists treating you that you are using Presmedic Ambroxol Tablet 30mg.

-Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Presmedic Ambroxol Tablet 30mg to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

The benefits of the medicine in pregnant women during first trimester should be weighed against a possible risk to the fetus. Consult your doctor or pharmacist if you are pregnant or breastfeeding.

Side effects

All medicines can cause unwanted side effects although not everyone experience it. See a doctor immediately if you have any of the following serious side effects. You may need urgent medical treatment.

The side effects of ambroxol include mild upper gastrointestinal effects such as heartburn, indigestion and bloated.

PRESMEDIC AMBROXOL TABLET 30 MG

Ambroxol hydrochloride (30mg)

Allergic reactions such as skin rashes, difficulty in breathing, swelling of the face, lips, mouth, tongue or throat which may cause difficulty swallowing or breathing have occurred rarely.

Reports of severe acute anaphylactic-type reactions including anaphylactic shock but their relationship to ambroxol is uncertain and are extremely rare case.

Consult your doctor/ pharmacist for further information. Other unwanted side effects not listed may occur.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction monitoring by visiting the website npra.gov.my [Consumers→Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Presmedic Ambroxol Tablet 30mg

-Storage

Keep out of reach of children.
Store below 30°C.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

-What it looks like

Round, white uncoated tablet, flat bevel-edged, with break bar line on one side and plain on the other side.

-Ingredients

-Active ingredients

Ambroxol Hydrochloride

-Inactive ingredients

Lactose

Microcrystalline Cellulose

Sodium Starch Glycolate

Magnesium Stearate

Talc

Colloidal Silicon Dioxide

-MAL Number:

MALxxxxxxxxX

Manufacturer

HOVID Bhd.

Lot 56442, 7 ½ Miles Jalan

Ipoh/Chemor,

31200, Chemor, Perak, Malaysia.

Product Registration Holder

HOVID Pharmacy Sdn. Bhd.

121, Jalan Tunku Abdul Rahman

30010 Ipoh, Perak, Malaysia.

Date of revision

25/07/2023

Serial number

NPRA (R2) 23/066