

What is in this leaflet

1. What *Carvidon-MR* is used for
2. How *Carvidon-MR* works
3. Before you use *Carvidon-MR*
4. How to use *Carvidon-MR*
5. While you are using it
6. Side effects
7. Storage and Disposal of *Carvidon-MR*
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Carvidon-MR is used for

This medicine is intended for use in combination with other medicine to treat angina pectoris (chest pain caused by coronary disease).

How Carvidon-MR works

Trimetazidine acts by inhibiting the breakdown of fatty acids which enhances glucose oxidation. Glucose oxidation requires less oxygen consumption thus optimises the heart's energy producing function, protecting vulnerable heart muscle tissues from further oxygen-deprived damage

Before you use Carvidon-MR

- When you must not use it

Do not take Carvidon-MR

- If you are allergic to Trimetazidine or to any other ingredients of this medicine.
- If you have a Parkinson disease: disease of the brain affecting movement (trembling, rigid posture, show movements and a shuffling, unbalanced walk.
- If you have severe kidney problems.

Pregnancy and lactation

Safety of use of Carvidon-MR in pregnant and nursing mothers has not been established.

As a precaution, Carvidon-MR is not to be taken during pregnancy or breastfeeding.

Tell your doctor if you are pregnant, think you are pregnant or planned to get pregnant

- Before you start use it

- Talk to your doctor or pharmacist before taking Carvidon-MR

- Taking other medicines

You must inform your doctor or pharmacist if you are taking, or have recently taken any other medicines, even if it is a medicine that you can buy without a prescription.

How to use Carvidon-MR

- How much to use

- Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.
- The recommended dose is Carvidon-MR tablet is one tablet to be taken two times a day during meals in the morning and evening.
- If you have kidney problems or if you are older than 75 years old, your doctor may adjust the recommended dose.

- When to use it

Use as directed by your doctor or pharmacist.

- One tablet to be taken two times a day during meals in the morning and evening.
- Tablet should be taken with a glass of water with meal times

- How long to use it

Continue taking *Carvidon-MR* for as long as your doctor recommends.

- If you forget to use it

Resume treatment normally. Do not double the doses the day after you miss a dose

- If you use too much (overdose)

Inform your doctor or pharmacist

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking *Carvidon-MR*.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give *Carvidon-MR* to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Driving and using machines

This medicine may make you feel dizzy and drowsy that may affect your ability to drive or use machinery.

Side effects

Like all medicines, *Carvidon-MR* can cause side effects, although not everybody gets them.

Common: Dizziness, headache, abdominal pain, diarrhoea, indigestion, feeling sick, vomiting, rash, itching, hives and feeling of weakness.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website nppra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI)"

Storage and Disposal of Carvidon-MR

- Storage

Keep out of the reach and sight of children.

Store below 30°C.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Light pink colored, circular, biconvex, film coated tablets, plain on both the faces

- Ingredients

- Active ingredient(s)
Trimetazidine Dihydrochloride
- Inactive ingredients

CARVIDON-MR

Trimetazidine Dihydrochloride Modified Release Tablets 35 mg

The other ingredients are Microcrystalline Cellulose, Polyethylene Oxide, Hypromellose, ethyl cellulose 20 cps, titanium dioxide, ferric oxide red, isopropyl alcohol and methylene chloride, Purified water.

- MAL number(s):
MAL*****

Manufacturer

MICRO LABS LIMITED
92, SIPCOT, Hosur
Tamil Nadu - – 635 126, INDIA

Product Registration Holder

Go Pharma Sdn Bhd
No 13 Jalan Rajawali 2
Bandar Puchong Jaya
47100 Puchong Selangor

Date of revision

28th June, 2017