



200 mg/ml Solution for Injection



STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each ml contains:

Active substance: Iron 200 mg (as gleptoferron complex)

Excipients: Phenol 5 mg (as preservative)

INDICATION(S)

Piglets

For the prevention and treatment of iron deficiency anaemia

PHARMACODYNAMIC

The product is a solution of macromolecular complex containing 200 mg of iron per ml: It stimulates hematopoiesis by progressive release of the dextran complex. It provides in one single injection of 1 ml 200 mg of iron, a dose necessary to prevent iron-deficiency anemia in piglet and stimulates its growth. The product allows an almost immediate reconstitution of the iron reserves necessary for the formation of hemoglobin and the recharging of iron-based enzymes which play an important role in the growth and resistance to infections.

PHARMACOKINETIC

The product is rapidly and completely absorbed.

DOSAGE AND ADMINISTRATION



200 mg of iron per piglet by intramuscular route ie 1 ml in one injection within the 3 first days of life of the piglet, preferably in muscles of the neck.



Remark: it is possible to adapt the vial with automatic injector that permits fast and sterile administration of 1 ml of the solution per piglet.

For 100 ml collapsible vial: as the vial cannot be broached more than 4 times, the use of automatic syringe equipment is recommended.

For 100 & 250 ml plastic vials: as the vial cannot be broached more than 20 times, use of automatic syringe equipment is recommended.



CONTRAINDICATION

None.



SPECIAL PRECAUTIONS FOR USE

Special warnings during use in animals

Strictly respect the aseptic conditions during the injection. Proceed with caution in case of sensitivity to iron dextran, due to a nutritional predisposition of maternal origin.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

None.

Other precautions

None.

INTERACTION WITH OTHER MEDICINAL PRODUCTS

Not known.

00000

USE DURING PREGNANCY, LACTATION OR LAY

Not applicable.

ADVERSE REACTION

Coloration at the injection site.

Very rare cases of death in piglets occurred after administration of parenteral iron preparations. These deaths have been linked to genetic factors or deficiency in Vitamin E and/or Selenium.

OVERDOSE

Not Known.

INCOMPATIBILITIES

Not known.

WITHDRAWAL PERIOD

Meat and offal: 0 days.

STORAGE

Store below 30°C, protected from light.
Once broached, it can be used within 28 days

PACK SIZES

100 ml and 250 ml.
Not all pack sizes may be marketed.

Keep out of reach of children.

Registered by:

Ceva Animal Health Malaysia Sdn Bhd
27-2, Oval Damansara.
No. 685, Jalan Damansara 60000 Kuala Lumpur.

Manufactured by:

Ceva Santé Animale
10 avenue de la Ballastière, 33500 Libourne ,France

DATE OF REVISION OF PACKAGE INSERT: 04/2021.