



Solution for injection  
for cattle, pigs and sheep



## NAME OF THE VETERINARY MEDICINAL PRODUCT

Tulaven 100 mg/ml solution for injection for cattle, pigs and sheep

## STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each ml contains:

Active substance:

Tulathromycin: 100 mg

Excipients:

Monothioglycerol: 5 mg

## PHARMACODYNAMIC

Tulathromycin is a semi-synthetic macrolide antimicrobial agent, which originates from a fermentation product. It differs from many other macrolides in that it has a long duration of action that is, in part, due to its three amine groups; therefore, it has been given the chemical subclass designation of triamilide. Macrolides are bacteriostatic acting antibiotics and inhibit essential protein biosynthesis by virtue of their selective binding to bacterial ribosomal RNA. They act by stimulating the dissociation of peptidyl-tRNA from the ribosome during the translocation process.

Tulathromycin possesses *in vitro* activity against *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni* and *Mycoplasma bovis*, and *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Mycoplasma hyopneumoniae*, *Haemophilus parasuis* and *Bordetella bronchiseptica*, the bacterial pathogens most commonly associated with bovine and swine respiratory disease, respectively. Increased minimum inhibitory concentration (MIC) values have been found in some isolates of *Histophilus somni* and *Actinobacillus pleuropneumoniae*. *In vitro* activity against *Dichelobacter nodosus* (vir), the bacterial pathogen most commonly associated with infectious pododermatitis (foot rot) in sheep, has been demonstrated. Tulathromycin also possesses *in vitro* activity against *Moraxella bovis*, the bacterial pathogen most commonly associated with infectious bovine keratoconjunctivitis (IBK).

The Clinical and Laboratory Standards Institute CLSI has set the clinical breakpoints for tulathromycin against *M. haemolytica*, *P. multocida*, and *H. somni* of bovine respiratory origin and *P. multocida* and *B. bronchiseptica* of swine respiratory origin as  $\leq 16 \mu\text{g/ml}$  susceptible and  $\geq 64 \mu\text{g/ml}$  resistant. For *A. pleuropneumoniae* of swine respiratory origin the susceptible breakpoint is set at  $\leq 64 \mu\text{g/ml}$ . CLSI has also published clinical breakpoints for tulathromycin based on a disk diffusion method (CLSI document VET08, 4th ed, 2018). No clinical breakpoints are available for *H. parasuis*. Neither EUCAST nor CLSI have developed standard methods for testing antibacterial agents against veterinary *Mycoplasma* species and thus no interpretative criteria have been set.

Resistance to macrolides can develop by mutations in genes encoding ribosomal RNA (rRNA) or some ribosomal proteins; by enzymatic modification (methylation) of the 23S rRNA target site, generally giving rise to cross-resistance with lincosamides and group B streptogramins (MLS<sup>B</sup> resistance); by enzymatic inactivation; or by macrolide efflux. MLS<sup>B</sup> resistance may be constitutive or inducible. Resistance may be chromosomal or plasmid-encoded and may be transferable if associated with transposons, plasmids integrative and conjugative elements. Additionally, the genomic plasticity of *Mycoplasma* is enhanced by the horizontal transfer of large chromosomal fragments.

In addition to its antimicrobial properties, tulathromycin demonstrates immune-modulating and anti-inflammatory actions in experimental studies. In both bovine and porcine polymorphonuclear cells (PMNs; neutrophils), tulathromycin promotes apoptosis (programmed cell death) and the clearance of apoptotic cells by macrophages. It lowers the production of the pro-inflammatory mediators leukotriene B<sub>4</sub> and CXCL-8 and induces the production of anti-inflammatory and pro-resolving lipid lipoxin A<sub>4</sub>.

## PHARMACOKINETICS

In cattle, the pharmacokinetic profile of tulathromycin when administered as a single subcutaneous dose of 2.5 mg/kg bodyweight, was characterised by rapid and extensive absorption followed by high distribution and slow elimination. The maximum concentration (C<sub>max</sub>) in plasma was approximately 0.5  $\mu\text{g/ml}$ ; this was achieved approximately 30 minutes post-dosing (T<sub>max</sub>). Tulathromycin concentrations in lung homogenate were considerably higher than those in plasma. There is strong evidence of substantial accumulation of tulathromycin in neutrophils and alveolar macrophages. However, the *in vivo* concentration of tulathromycin at the infection site of the lung is not known. Peak concentrations were followed by a slow decline in systemic exposure with an apparent elimination half-life (t<sub>1/2</sub>) of 90 hours in plasma. Plasma protein binding was low, approximately 40%. The volume of distribution at steady-state (VSS) determined after intravenous administration was 11 l/kg. The bioavailability of tulathromycin after subcutaneous administration in cattle was approximately 90%.

In pigs, the pharmacokinetic profile of tulathromycin when administered as a single intramuscular dose of 2.5 mg/kg bodyweight, was also characterised by rapid and extensive absorption followed by high distribution and slow elimination. The maximum concentration (C<sub>max</sub>) in plasma was approximately 0.6  $\mu\text{g/ml}$ ; this was achieved approximately 30 minutes post-dosing (T<sub>max</sub>). Tulathromycin concentrations in lung homogenate were considerably higher than those in plasma. There is strong evidence of substantial accumulation of tulathromycin in neutrophils and alveolar macrophages. However, the *in vivo* concentration of tulathromycin at the infection site of the lung is not known. Peak concentrations were followed by a slow decline in systemic exposure with an apparent elimination half-life (t<sub>1/2</sub>) of approximately 91 hours in plasma. Plasma protein binding was low, approximately 40%. The volume of distribution at steady-state (VSS) determined after intravenous administration was 13.2 l/kg. The bioavailability of tulathromycin after intramuscular administration in pigs was approximately 88%.

In sheep, the pharmacokinetic profile of tulathromycin, when administered as a single intramuscular dose of 2.5 mg/kg bodyweight, achieved a maximum plasma concentration (C<sub>max</sub>) of 1.19  $\mu\text{g/ml}$  in approximately 15 minutes (T<sub>max</sub>) post-dosing and had an elimination half-life (t<sub>1/2</sub>) of 69.7 hours.

Plasma protein binding was approximately 60-75%. Following intravenous dosing the volume of distribution at steady-state (VSS) was 31.7 l/kg. The bioavailability of tulathromycin after intramuscular administration in sheep was 100%.

## INDICATION(S)

**Cattle:** Treatment and metaphylaxis of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni* and *Mycoplasma bovis* sensitive to tulathromycin. The presence of the disease in the herd should be established before metaphylactic treatment.

Treatment of infectious bovine keratoconjunctivitis (IBK) associated with *Moraxella bovis* sensitive to tulathromycin.

**Pigs:** Treatment and metaphylaxis of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Mycoplasma hyopneumoniae*, *Haemophilus parasuis* and *Bordetella bronchiseptica* sensitive to tulathromycin. The presence of the disease in the herd should be established before metaphylactic treatment. The product should only be used if pigs are expected to develop the disease within 2–3 days.

**Sheep:** Treatment of the early stages of infectious pododermatitis (foot rot) associated with virulent *Dichelobacter nodosus* requiring systemic treatment.

## CONTRAINDICATIONS

Do not use in cases of hypersensitivity to macrolide antibiotics or to any of the excipients.

## ADVERSE REACTIONS

Transient pain reactions and local swellings at the injection site are very common after subcutaneous administration to cattle; they can persist for up to 30 days. No such reactions have been observed in pigs and sheep after intramuscular administration.

Pathomorphological injection site reactions (including reversible changes of congestion, oedema, fibrosis and haemorrhage) are very common for approximately 30 days after injection in cattle and pig. In sheep transient signs of discomfort (head shaking, rubbing injection site, backing away) are very common after intramuscular injection. These signs resolve within a few minutes.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

## TARGET SPECIES

Cattle, pigs and sheep.

## DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

**Cattle:** Subcutaneous use.

A single subcutaneous injection of 2.5 mg tulathromycin/kg bodyweight (equivalent to 1 ml/40 kg bodyweight). For treatment of cattle over 300 kg bodyweight, divide the dose so that no more than 7.5 ml are injected at one site.

**Pigs:** Intramuscular use.

A single intramuscular injection of 2.5 mg tulathromycin/kg bodyweight (equivalent to 1 ml/40 kg bodyweight) in the neck. For treatment of pigs over 80 kg bodyweight, divide the dose so that no more than 2 ml are injected at one site.

**Sheep:** Intramuscular use.

A single intramuscular injection of 2.5 mg tulathromycin/kg body weight (equivalent to 1 ml/40 kg bodyweight) in the neck.

## ADVICE ON CORRECT ADMINISTRATION

For any respiratory disease, it is recommended to treat animals in the early stages of the disease and to evaluate the response to treatment within 48 hours after injection. If clinical signs of respiratory disease persist or increase, or if relapse occurs, treatment should be changed, using another antibiotic, and continued until clinical signs have resolved. To ensure correct dosage bodyweight should be determined as accurately as possible to avoid underdosing. For multiple vial entry, an aspirating needle or multi-dose syringe is recommended to avoid excessive broaching of the stopper.

## WITHDRAWAL PERIOD(S)

Meat and offal for:

Cattle: 22 days.

Pigs: 13 days.

Sheep: 16 days.

Not authorised for use in animals producing milk for human consumption. Do not use in pregnant animals, which are intended to produce milk for human consumption, within 2 months of expected parturition.

## SPECIAL STORAGE CONDITIONS

Keep out of the sight and reach of children.

Store at room temperature.

Shelf-life after first broaching the vial: 28 days.

## SPECIAL WARNING(S)

### Special warnings for each target species

Cross resistance occurs with other macrolides. Do not administer simultaneously with antimicrobials with a similar mode of action such as other macrolides or lincosamides.

**Sheep:** The efficacy of antimicrobial treatment of foot rot might be reduced by others factors, such as wet environmental conditions, as well as inappropriate farm management. Treatment of foot rot should therefore be undertaken along with other flock management tools, for example providing dry environment. Antibiotic treatment of benign foot rot is not considered appropriate. Limited efficacy was shown in sheep with severe clinical signs or chronic foot rot, and should therefore only be given at an early stage of foot rot.

**Special precautions for use in animals**

Use of the veterinary medicinal product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria. Official, national and regional antimicrobial policies should be taken into account when the product is used.

Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to tulathromycin and may decrease the effectiveness of treatment with other macrolides lincosamides and group B streptogramins, due to the potential for cross resistance.

If a hypersensitivity reaction occurs, appropriate treatment should be administered without delay.

**Special precautions to be taken by the person administering the veterinary medicinal product to animals**

Tulathromycin is irritating to eyes. In case of accidental eye exposure, flush the eyes immediately with clean water. Tulathromycin may cause sensitisation by skin contact resulting in e.g. reddening of the skin (erythema) and/or dermatitis. In case of accidental spillage onto skin, wash the skin immediately with soap and water. Wash hands after use. In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

If there is suspicion of a hypersensitivity reaction following accidental exposure (recognised by e.g. itching, difficulty in breathing, hives, swelling on the face, nausea, vomiting) appropriate treatment should be administered. Seek medical advice immediately and show the package leaflet or the label to the physician.

**Pregnancy and lactation**

Laboratory studies in rats and rabbits have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. The safety of the veterinary medicinal product has not been established during pregnancy and lactation. Use only according to the benefit/risk assessment by the responsible veterinarian.

**Interaction with other medicinal products and other forms of interaction**

None known.

**Overdose (symptoms, emergency procedures, antidotes)**

In cattle at dosages of three, five or ten times the recommended dose, transient signs attributed to injection site discomfort were observed and included restlessness, head-shaking, pawing the ground, and brief decrease in feed intake. Mild myocardial degeneration has been observed in cattle receiving five to six times the recommended dose. In young pigs weighing approximately 10 kg given three or five times the therapeutic dose transient signs attributed to injection site discomfort were observed and included excessive vocalisation and restlessness. Lameness was also observed when the hind leg was used as the injection site. In lambs (approx. 6 weeks old), at dosages of three or five times the recommended dose, transient signs attributed to injection site discomfort were observed, and included walking backwards, head shaking, rubbing the injection site, lying down and getting up, bleating.

**Incompatibilities**

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

**SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIAL, IF ANY**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

**OTHER INFORMATION**

Pack sizes:

Cardboard box containing one vial.

Glass vial size: 20 ml

Plastic vial sizes: 50 ml, 100 ml, 250 ml and 500 ml.

Not all pack sizes may be marketed.

**FOR ANIMAL TREATMENT ONLY**

Manufactured by:

Ceva Santé Animale

10 Avenue De La Ballastiere

33500 Libourne

France

|                                                                                                     |                                                                             |
|-----------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|
| NOT TULAVEN 100MG/ML 20ML MY<br>680x160/FP 35.5x160<br>Recto Verso<br>CODE ARTICLE : 00000<br>BLACK | TULAVEN 100MG/ML MY<br>ID : 67627-67628-67629-67630-67631<br>CORPS : 11 pts |
|-----------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|

OLIVIER 05/10/20 - 22/02/22