

JANIPRO TRIPAXIL CAPSULE

Glucosamine Sulphate 250mg, Chondroitin Sulphate Sodium 200mg, Methylsulfonylmethane 100mg

What is in this leaflet

1. What Janipro Tripaxil Capsule is used for
2. How Janipro Tripaxil Capsule works
3. Before you use Janipro Tripaxil Capsule
4. How to use Janipro Tripaxil Capsule
5. While you are using it
6. Side effects
7. Storage and Disposal of Janipro Tripaxil Capsule
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial Number

What Janipro Tripaxil Capsule is used for

It is used as adjuvant therapy for osteoarthritis.

How Janipro Tripaxil Capsule works

It contains glucosamine which can stimulate the production of cartilage-building compounds, restore damaged cells of cartilage tissue and has some anti-inflammatory abilities. It also contains chondroitin that is needed for the making of cartilage- building compounds and helps to keep cartilage healthy by absorbing and retaining water in the joint as well as methylsulfonylmethane that can help to form connective tissues.

Before you use Janipro Tripaxil Capsule

- When you must not use it

Do not use it if you are allergic to glucosamine sulphate, chondroitin or methylsulfonylmethane.

Do not use it if you are allergic to seafood.

- Before you start to use it

1) Effects on glucose metabolism & antidiabetic agents: It has been hypothesized that glucosamine may impair insulin secretion through competitive inhibition of glucokinase in pancreatic beta cells and/or alteration of peripheral glucose uptake. Glucosamine may increase insulin resistance and consequently affect glucose tolerance. It may reduce antidiabetic agent effectiveness e.g. when used with these antidiabetic agents: Acarbose,

Acetohexamide, Chlorpropamide, Glipizide, Glyburide, Metformin, Miglitol, Pioglitazone, Repaglinide, Rosiglitazone, Glimepiride, Tolbutamide, Troglitazone, Glucosamine is likely safe in patients with well-controlled diabetes (HbA1c less than 6.5%) taking one or two oral antidiabetic medications or controlled by diet only. In patients with higher HbA1c levels or those taking insulin, monitor blood glucose levels closely/more frequently.

2) Reduced effectiveness when used with glucosamine: Doxorubicin, Etoposide, Teniposide

3) Warfarin - Elevations of International Normalized Ratio serum values and potentiation of anticoagulant effects - If concomitant therapy is necessary, the patient's INR should be more closely monitored

4) Avoid combination with chitosan as it may form complexes with chondroitin sulfate, thus decreasing its absorption

Available evidence is inconclusive or inadequate for use in pregnant or lactating mothers. Until more information is available, this product should only be used under medical supervision in pregnancy and lactating mothers if the potential benefit to the mother justifies the potential risk to the fetus.

Administration during the first 3 months of pregnancy must be avoided.

- Taking other medicines

Please inform the doctor or pharmacist if taking other medicines.

How to use Janipro Tripaxil Capsule

- How much to use

Light or moderate osteoarthritis symptoms:

2 capsules twice daily for at least 6 weeks or according to medical prescription.

Severe osteoarthritis symptoms:

Initial therapy: 2 capsules three times daily for at least 8 weeks according to medical prescription.

Follow-up therapy: Maintenance therapy of 2 capsules twice daily should be followed for 3-4 months or according to medical prescription.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Please follow the advice of the doctor or pharmacist.

- How long to use it

Please follow the advice of the doctor or pharmacist.

- If you use too much (overdose)

No cases of accidental or intentional overdose are known.

While you are using Janipro Tripaxil Capsule

- Things you must do

Inform your doctor or pharmacist of all the medications you are taking.

- Things you must not do

Do not take any new medicines without consulting your doctor. Do not give Janipro Tripaxil Capsule to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Ask your health care provider any question you may have about how to use Janipro Tripaxil Capsule.

Side effects

Cardiovascular: Peripheral oedema, tachycardia was reported in a few patients following larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.

Central Nervous System: Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%).

Gastrointestinal: Nausea, vomiting, diarrhea, dyspepsia or epigastric pain, constipation, heartburn and anorexia

JANIPRO TRIPAXIL CAPSULE

Glucosamine Sulphate 250mg, Chondroitin Sulphate Sodium 200mg, Methylsulfonylmethane 100mg

have been described rarely during oral therapy with glucosamine.

Date of revision

07.03.2024

Skin: Skin reactions such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

Serial Number

NPRA(R2)23/159

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Janipro Tripaxil Capsule**- Storage**

Store below 30°C. Protect from light and moisture.

Keep out of reach of children.

- Disposal

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description**- What it looks like**

White colour powder filled in hard capsule.

Capsule colour: scarlet-scarlet.

Capsule size 0

No marking on capsule.

- Ingredients

- Active ingredient

Glucosamine Sulphate, Chondroitin Sulphate Sodium, Methylsulfonylmethane

- MAL number:

MALXXXXX

Manufacturer

Syarikat Wen Ken Drug Sdn Bhd

24, Jalan Lambak, Taman Johor, 81200

Johor Bahru, Johor, Malaysia.

Product Registration Holder

Janipro (M) Sdn Bhd

No. 42, Jalan Seksyen 1/21,

Taman Kajang Utama,

43000 Kajang, Selangor.