

JANIPRO DOUBAXIL POWDER

Glucosamine Sulphate 1500mg, Chondroitin Sulphate Sodium 1200mg

What is in this leaflet

1. What Janipro Doubaxil Powder is used for
2. How Janipro Doubaxil Powder works
3. Before you use Janipro Doubaxil Powder
4. How to use Janipro Doubaxil Powder
5. While you are using it
6. Side effects
7. Storage and Disposal of Janipro Doubaxil Powder
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial Number

What Janipro Doubaxil Powder is used for

It is used as adjuvant therapy for osteoarthritis.

How Janipro Doubaxil Powder works

It contains glucosamine which can stimulate the production of cartilage-building compounds, restore damaged cells of cartilage tissue and has some anti-inflammatory abilities. It also contains chondroitin that is needed for the making of cartilage- building compounds and helps to keep cartilage healthy by absorbing and retaining water in the joint.

Before you use Janipro Doubaxil Powder

- When you must not use it

- If you are allergy to glucosamine sulphate or chondroitin or seafood.
- If you are taking Warfarin (or other coumarin anticoagulants).
- If you are taking oral tetracycline, do not take glucosamine at the same time.
- If you have kidney or liver problems.

- Before you start to use it

Talk to your doctor or pharmacist if you are having or have had asthma, diabetes, high cholesterol, high blood pressure, a stroke, heart problems or on a controlled sodium diet as your doctor/ pharmacist may need to monitor you more regularly.

Avoid combination with chitosan as it may form complexes with chondroitin sulfate, thus decreasing its absorption.

Pregnancy and lactation

Available evidence is inconclusive or inadequate for use in pregnant or lactating mothers. Until more information is available, this product should only be used under medical supervision in pregnancy and lactating mothers if the potential benefit to the mother justifies the potential risk to the fetus.

Administration during the first 3 months of pregnancy must be avoided.

- Taking other medicines

Please inform the doctor or pharmacist if taking other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use Janipro Doubaxil Powder

- How much to use

Adult: One sachet to be taken once a day. The entire contents of one sachet should be fully dissolved in at least 250 ml of water (one glass) and drink immediately. It is best to be taken before meal for at least 8-12 weeks or more, according to medical prescription.

- When to use it

Please follow the advice of the doctor or pharmacist.

- How long to use it

Please follow the advice of the doctor or pharmacist.

- If you forget to use it

Take the missed dose as soon as you remember and continue to take it as you would normally. If it is almost time for your next dose, skip the missed dose and take the next dose when you are meant to. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital if you or anyone else may have taken too much of this medicine.

While you are using Janipro Doubaxil Powder

- Things you must do

Tell all the doctors, dentists and pharmacists that are treating you that you are taking Janipro Doubaxil Powder.

Tell your doctor immediately if you become pregnant while taking Janipro Doubaxil Powder.

- Things you must not do

Do not take any new medicines without consulting your doctor.

- Things to be careful of

Ask your health care provider any question you may have about how to use Janipro Doubaxil Powder.

Side effects

Cardiovascular: Peripheral oedema, tachycardia was reported in a few patients following larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.

Central Nervous System: Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%).

Gastrointestinal: Nausea, vomiting, diarrhea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been described rarely during oral therapy with glucosamine.

Skin: Skin reactions such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Janipro Doubaxil Powder

- Storage

Store below 30°C. Protect from light and moisture.

Keep out of reach of children.

JANIPRO DOUBAXIL POWDER

Glucosamine Sulphate 1500mg, Chondroitin Sulphate Sodium 1200mg

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

White or almost white crystalline powder is filled in aluminium sachet. The solution is clear and colourless with salty taste after dissolved the powder in water.

- Ingredients

- Active ingredients
Glucosamine Sulphate, Chondroitin Sulphate Sodium

- MAL number:

MALXXXXX

Manufacturer

Syarikat Wen Ken Drug Sdn Bhd
24, Jalan Lambak, Taman Johor, 81200
Johor Bahru, Johor, Malaysia.

Product Registration Holder

Janipro (M) Sdn Bhd
No. 42, Jalan Seksyen 1/21,
Taman Kajang Utama,
43000 Kajang, Selangor.

Date of revision

04.02.2025

Serial Number

NPRA(R2)24/180