

REMECO PEPTICON ADVANCE ORAL SUSPENSION

Sodium Alginate, Potassium Bicarbonate (1000mg/10ml, 200mg/10ml)

What is in this leaflet

1. What Pepticon Advance Oral Suspension is used for
2. How Pepticon Advance Oral Suspension works
3. Before you take Pepticon Advance Oral Suspension
4. How to take Pepticon Advance Oral Suspension
5. While you are taking Pepticon Advance Oral Suspension
6. Side Effects
7. Storage and Disposal of Pepticon Advance Oral Suspension
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision
11. Serial Number

What Pepticon Advance Oral Suspension is used for

Treatment of symptoms of gastroesophageal reflux such as acid regurgitation, heartburn and indigestion.

How Pepticon Advance Oral Suspension works

Remeco Pepticon Advance Oral Suspension relieves the pain & discomfort of heartburn and acid indigestion by forming a protective layer on top of the stomach contents to prevent stomach acid escaping up into the food pipe where it hurts.

Before you take Pepticon Advance Oral Suspension

- When you must not take it
Do not take this medicine if you have an allergy to any of the ingredients including the excipients (parabens).

Pregnancy and lactation:

There is no significant adverse effects of the product known on the course of pregnancy or on the health of the foetus/ new-born child.

- Before you start to take it
Tell your doctor if you are allergic to any medicines or if you have any other medical condition.
- Taking other medicines
Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to take Pepticon Advance Oral Suspension

- How much to take

For oral use. Shake well before use.

Adults and children over 12 years: Take 5-10ml (One to two 5ml spoonfuls) after meal and at bedtime.

Children under 12 years: Not be given unless recommended by your doctor.

- When to take it

after meal and at bedtime

- How long to take it

Continue taking Pepticon Advance Oral Suspension for as long as your doctor recommends.

- If you forget to take it

Consult your doctor or pharmacist on what you should do if you forget to use it. Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you take too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are taking Pepticon Advance Oral Suspension

- Things you must do

Follow your doctor's or pharmacist's instruction when you are taking Pepticon Advance Oral Suspension.

- Things you must not do

Do not use this product after the expiry date which is stated on the label.

- Things to be careful of

Each 10ml dose has a sodium content of 125.5mg and 78mg of potassium. This should be taken into account when a highly restricted salt diet is recommended.

Each 10ml contains 200 mg of calcium carbonate. Care needs to be taken in treating patients with hypercalcaemia (increase of blood calcium level), nephrocalcinosis (accumulation of calcium in the kidneys) and recurrent calcium containing renal calculi.

This medicinal product contains methyl paraben and propyl paraben which may cause allergic reactions (possibly delayed).

Side Effects

Like all medicines, Pepticon Advance Oral Suspension can cause side effects, although not everybody gets them.

If you experience one of the following, stop using this product and inform your doctor immediately or go to the nearest hospital emergency department:

- Skin rash, pruritus
- Breathing difficulties
- Dizziness
- Swelling of the face, lips, tongue or throat

All of these very serious side effects are very rare.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Dutavor

- Storage

Store below 30°C. Do not refrigerate or freeze. Check that the cap seal is unbroken before first taking this product.

Use within 6 months after first opening.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

REMECO PEPTICON ADVANCE ORAL SUSPENSION

Sodium Alginate, Potassium Bicarbonate (1000mg/10ml, 200mg/10ml)

Product Description

- What it looks like

Pepticon Advance Oral Suspension is an opaque, almost white and peppermint odour viscous suspension.

- Ingredients

- Active ingredient:

Sodium alginate, potassium bicarbonate

- Inactive ingredients:

Calcium carbonate, carbomer 974P, sodium hydroxide, sodium saccharin, peppermint oil, methyl paraben, propyl paraben, purified water

- MAL number:

MAL25046041XC

Manufacturer

Vefa İlaç Sanayi ve Ticaret A.Ş.
Beylikdüzü OSB Mah. 2. Cadde No:
3/2 Beylikdüzü/ISTANBUL/TURKEY.

Product Registration Holder

Abio Marketing Sdn Bhd
No. 2, Jalan SS 13/5,
47500 Subang Jaya, Malaysia.

Date of Revision

15/05/2025

Serial Number

NPRA (R2) 25/008