

Item Number: 0756139		Version: 1		Swatches														
Profile: 0402172_1_VL_Drwg		Technical Specification: 0402172_TSEL02		Black	PANTONE 7417 li													
Artwork Dimensions/Size: 95 x 32 mm				PANTONE 185 C														
Graphics House: 106451345 / 402659595 / SGK München																		
Date: 15Dec2021				Visible / Not Printed*														
OPTIONAL: Artwork Approver outside the Shire Management System:				Dieline	Technical Info													
Role:	Name:	Signature:	Date:	 Confidential Property of Takeda														
			<table border="1"> <tr> <td>Body Text Size</td> </tr> <tr> <td>7 pt</td> </tr> </table>	Body Text Size	7 pt	<table border="1"> <tr> <td>Code ITF 2/5</td> <td>Datamatrix Code</td> <td>RSS / GS1-128 Code</td> </tr> <tr> <td>756139</td> <td>NA</td> <td>NA</td> </tr> <tr> <td>Pharmacode</td> <td>Code 3 of 9</td> <td>GTIN / EAN-13 Code</td> </tr> <tr> <td>NA</td> <td>NA</td> <td>NA</td> </tr> </table>	Code ITF 2/5	Datamatrix Code	RSS / GS1-128 Code	756139	NA	NA	Pharmacode	Code 3 of 9	GTIN / EAN-13 Code	NA	NA	NA
Body Text Size																		
7 pt																		
Code ITF 2/5	Datamatrix Code	RSS / GS1-128 Code																
756139	NA	NA																
Pharmacode	Code 3 of 9	GTIN / EAN-13 Code																
NA	NA	NA																

* Color not present in printed material.

top label / front side

IMMUNATE 1000 IU		IMMUNATE 1000 IU		IMMUNATE 1000 IU Powder and solvent for solution for injection Active Substance: Human Coagulation Factor VIII, VHSD (Vapour Heated and Solvent Detergent) treated This product contains animal part(s) (porcine/pig)	
Lot:	EXP.:	Lot:	EXP.:	Lot:	EXP.:
				MAL20061558AZ Takeda Manufacturing Austria AG, 1221 Wien/Vienna 0756139	
				 	

top label / reverse side

Powder and solvent for solution for injection

1 vial provides: 1000 IU human coagulation factor VIII and 750 IU von Willebrand Faktor activity (VWF:RCO); appr. 100 IU FVIII/ml and 75 IU VWF/ml after reconstitution. Human Albumin, Glycine, Sodium chloride, Sodium citrate, Lysine HCl, Calcium chloride. Read package leaflet before use. For single use only.

base label / front side

For intravenous use. Use the reconstituted solution immediately (within 3 hours). Slow (**max. 2 ml/min**) i. v. injection. Discard residues. Store at 2° C – 8° C (in a refrigerator). Do not freeze. Store in the original package in order to protect from light. Keep out of the reach and sight of children. Medicinal product subject to medicinal prescription.