

MEZAVANT XL gastro-resistant, prolonged release tablets

Mezalazine 1200mg

What is in this leaflet

1. What *Mezavant XL* is used for
2. How *Mezavant XL* works
3. Before you use *Mezavant XL*
4. How to use *Mezavant XL*
5. While you are using it
6. Side effects
7. Storage and Disposal of *Mezavant XL*
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What *Mezavant XL* is used for

Pharmacotherapeutic group:
Aminosalicylic acid and similar agents.

Mezavant XL gastro-resistant, prolonged release tablets contain the active substance mesalazine, which is an anti-inflammatory drug for the treatment of ulcerative colitis.

Ulcerative colitis is a disease of the colon (large bowel) and rectum (back passage), where the lining of the gut becomes red and swollen (inflamed) resulting in symptoms of frequent and bloody stools together with stomach cramps.

How *Mezavant XL* works

When given for an acute episode of ulcerative colitis, *Mezavant XL* acts through the entire colon and rectum to treat the inflammation and reduce symptoms. The tablets can also be taken to help prevent recurrence of ulcerative colitis

Before you use *Mezavant XL*

When you must not use it

Do not take *Mezavant XL*

- If you are allergic (hypersensitive) to a family of drugs known as salicylates (which include acetylsalicylic acid (Aspirin))
- If you are allergic (hypersensitive) to mesalazine or any of the other ingredients of this medicine (listed in section Ingredients)
- If you have severe kidney or severe liver problems

Before you start use it

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you

Talk to your doctor before using *Mezavant XL*

- If you have any kidney or liver problems
- If you have previously had inflammation of the heart (which may be the result of an infection in the heart)
- If you have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after using *Mezavant XL*
- If you have had a previous allergic reaction to sulphasalazine (another medicine used to treat ulcerative colitis)
- If you have narrowing or blockage of the stomach or the gut.
- If you have lung problems

Before and periodically during treatment with *Mezavant XL*, your doctor may take samples of your urine and blood to check your kidneys and liver are working well and that your blood is healthy.

Kidney stones may develop with the use of *Mezavant XL*. Symptoms may include pain in the sides of the abdomen and blood in the urine. Take care to drink a sufficient amount of liquid during treatment with *Mezavant XL*.

Serious skin reactions including Drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson syndrome (SJC), and toxic epidermal necrolysis (TEN), have been reported in association with *Mezavant XL* treatment. Stop using *Mezavant XL* and seek medical attention immediately if you notice any of the symptoms related to these serious skin reactions described in section Side Effects.

Taking other medicines

Studies have shown that *Mezavant XL* does not interfere with the following antibiotics, used to treat infections: amoxicillin, metronidazole or sulfamethoxazole.

However, *Mezavant XL* may interact with some other medicines. Tell your

doctor or pharmacist if you are taking or have recently taken or might take any other medicines,

- Mesalazine or sulphasalazine (taken for treatment of ulcerative colitis)
- Non-steroidal anti-inflammatory drugs (for example medicines containing acetylsalicylic acid (Aspirin), ibuprofen or diclofenac)
- Azathioprine or 6-mercaptopurine (known as 'immunosuppressant' medicines which reduce the activity of your body's immune system).
- Coumarin-type anticoagulants (medicines which increase the time it takes for your blood to clot e.g. warfarin)

How to use *Mezavant XL*

How much to use

The recommended dose for adults is 2.4g to 4.8g (two to four tablets) taken once a day for an acute episode of ulcerative colitis. If you are taking the highest daily dose of 4.8g/day, you should be evaluated after 8 weeks treatment. Once your symptoms have cleared and to help prevent reoccurrence of another episode, your doctor should direct you to take 2.4g (two tablets) once a day.

When to use it

Always take the medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Mezavant XL is intended for once daily, oral administration and should be taken with food. The tablets should be swallowed whole and must not be crushed or chewed. Whilst taking this medicine ensure you drink fluids to remain well hydrated especially after severe or prolonged episodes of vomiting and/or diarrhoea, high fever or heavy sweating.

Mezavant XL is not recommended to be given to children under 18 years of age due to lack of data on safety and efficacy.

MEZAVANT XL gastro-resistant, prolonged release tablets

Mezalazine 1200mg

How long to use it

Continue to take this medicine as long as your doctor prescribes it. You should not stop taking this medicine even when you don't have any symptoms of ulcerative colitis without talking to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

If you forget to use it

It is important to take your Mezavant XL tablets everyday, even when you don't have any symptoms of ulcerative colitis. Always finish the prescribed course.

If you forget to take your tablets then take them as usual the next day. Do not take a double dose to make up for a forgotten tablet.

If you use too much (overdose)

If you take too much *Mezavant XL* you may have one or more of the following symptoms: tinnitus (ringing in ears), dizziness, headache, confusion, drowsiness, shortness of breath, excess loss of water (associated with sweating, diarrhoea and vomiting), low blood sugar (which can cause light-headedness), rapid breathing, changes in the blood chemistry and increased body temperature.

If you do take too many tablets, contact your doctor, pharmacist or hospital casualty department straight away. Take your tablet pack with you.

While you are using it

Things you must not do

- Do not stop taking your medicine or change the dosage without checking with your doctor.
- Do not give *Mezavant XL* to anyone else, even if they have the same condition as you. It may not be safe for another person to take *Mezavant XL*.

- Do not take *Mezavant XL* to treat any other complaints unless your doctors tell you to. It may not be safe to use *Mezavant XL* for another complaint.

Things to be careful of

Pregnancy and breast-feeding

Since mesalazine crosses the placenta in pregnancy and is excreted in breast milk in small quantities, you should only use *Mezavant XL* during pregnancy or whilst breast-feeding if your doctor tells you to. Adverse outcomes [including low blood cell counts (white blood cells, red blood cells, and platelets)] were reported in infants born to mothers who took *Mezavant XL* during pregnancy. Diarrhoea has been reported in breastfed infants of mothers who took *Mezavant XL*.

If you are pregnant or breast feeding, think you might be pregnant or are planning to have a baby ask your doctor for advice about taking *Mezavant XL* tablet.

Interference with laboratory tests

If you are undergoing urine tests, it is important to tell the doctor or nurse you are taking, or have recently taken this medicine as it can affect some results.

Driving or operating machinery

Mezavant XL tablet is unlikely to have any effect on your ability to drive or use machines.

This medicine contains less than 1 mmol sodium (23 mg) per the maximum recommended dose (4 tablets), that is to say essentially 'sodium free'.

Side effects

Tell your doctor immediately

- If you experience symptoms such as cramping, severe stomach pain, bloody and excessive stools (diarrhoea), fever, headache or rash. These symptoms could be a sign of Acute Intolerance Syndrome which can happen during an acute episode of ulcerative colitis. This is a serious condition which occurs rarely, but means your treatment would have

to be stopped immediately

- If you develop unexplained bruising (without injury), rash, anaemia (feeling tired, weak and looking pale, especially on lips, nails and inside of eyelids), fever (high temperature), sore throat or unusual bleeding (e.g. nose bleeds)
- If you notice reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes, widespread rash, fever and enlarged lymph nodes. These serious skin rashes can be preceded by fever and flu-like symptoms.
- If you develop allergic swelling of tongue, lips and around eyes
- If you experience strong or recurrent headache, disturbed vision, or ringing or buzzing in the ears. These could be symptoms of increased pressure within your skull (idiopathic intracranial hypertension).

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Common side effects, occurring in less than 1 in 10 patients, are headache, changes in blood pressure, flatulence (passing wind), nausea (feeling sick), bloated or painful stomach, inflammation which causes abdominal pain or diarrhoea; diarrhoea, indigestion, vomiting (being sick), abnormal liver function test, itching, rash, joint pain, back pain, weakness, fatigue (feeling extremely tired); fever (high temperature).

Uncommon side effects, seen in less than 1 in 100 patients are: a reduction in blood platelets which increases the risk of bleeding and bruising; dizziness; feeling sleepy or tired; trembling or shaking; ear pain; racing heartbeat; throat pain; an inflamed pancreas (associated with pain in upper abdomen and back and feeling sick); rectal polyp (a non-cancerous growth in the back passage causing symptoms such as constipation and bleeding); acne; hair loss; muscle pain; hives; swollen face.

MEZAVANT XL gastro-resistant, prolonged release tablets

Mezalazine 1200mg

Rare side effects, seen in less than 1 in 1000 patients are: kidney failure; severe reduction in the number of white blood cells that makes infection more likely; increased sensitivity of your skin to sun and ultraviolet light (photosensitivity).

The following side effects have been reported but it is not known exactly how often they occur: Severe reduction in blood cells which can cause weakness or bruising; low blood cell counts; allergic reaction (hypersensitivity); serious allergic reaction which causes difficulty in breathing or dizziness; serious illness with blistering of the skin (possibly leading to peeling of the skin and resulting in painful, raw areas), mouth, eyes and genitals; allergic reaction which causes skin rash, fever and inflammation of internal organs; rise in the pressure of the fluid surrounding the brain; neuropathy (abnormal or damaged nerves giving a sensation of numbness and tingling); inflammation of the heart and lining around the heart; inflammation of the lung; difficulty in breathing or wheezing; interstitial lung disease (a group of lung disorders which the lung tissues become inflamed and damaged); gall stones; hepatitis (inflammation of the liver giving rise to flu-like symptoms and jaundice); hepatotoxicity (liver damage that may present as abnormal liver tests); allergic swelling of tongue, lips and around eyes; skin redness; skin rash typically on face, skin sensitivity to sunlight along with joint pain, arthritis, fatigue and overall feeling sickness; kidney problems (such as inflammation and scarring of the kidney); kidney disorder causing excessive thirst and urination; nephrolithiasis (kidney stones and associated pain); low sperm count (reversible).

If you get any side effects talk to your doctor. This includes any possible side effects not listed in

this leaflet.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website www.npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Mezavant XL

Storage

- Keep this medicine out of the sight and reach of children
- Store below 30°C
- Store in the original package in order to protect from moisture
- Do not use this medicine after the expiry date which is stated on the box after “EXP”. The expiry date refers to the last day of the month.

Disposal

Do not throw away any medicines via waste water or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

Product Description

What it looks like

The red-brown tablets are oval shaped (dimensions 20.5 × 9.5 × 7.5 mm) and stamped S476.

Mezavant XL is supplied in foil blister strips which are contained in a cardboard box. The pack contains 60 tablets.

Ingredients

Active ingredient
Mesalazine

Inactive ingredients

Tablet core: Sodium Carboxymethylcellulose 7MF; Sodium Carboxymethylcellulose 7HXF; Carnauba Wax; Stearic Acid; Silica, Colloidal Hydrated; Sodium Starch Glycolate (Type A); Talc; Magnesium Stearate; Film-coating: Talc, Methacrylic Acid Copolymer, Type A; Methacrylic Acid Copolymer, Type B; Triethylcitrate; Titanium Dioxide; Red Ferric Oxide; Polyethylene glycol 6000.

MAL number(s):

MAL17085067ACZ

Manufacturer

Cosmo SpA
Via C. Colombo 1 20045 Lainate-Milan Italy

Product Registration Holder

Takeda Malaysia Sdn. Bhd.
Unit TB-L13-1, Level 13, Tower B, Plaza 33, No.1, Jalan Kemajuan, Seksyen 13, 46200 Petaling Jaya, Selangor Malaysia

Date of revision

23/06/2025

Serial Number

NPRA (R1/3) 15032021/046