

XYCIDE C (CHLORHEXIDINE GLUCONATE 0.5% W/V AND ISOPROPYL ALCOHOL 77% W/V HAND SANITIZER)

Chlorhexidine Gluconate 0.5% w/v, Isopropyl Alcohol 77% w/v

What Is In This Leaflet

1. What **Xycide C** is used for.
2. How **Xycide C** works.
3. Before you use **Xycide C**.
4. How to use **Xycide C**.
5. While you are using it
6. Side Effects
7. Storage and disposal of **Xycide C**.
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Xycide C Is Used For

Xycide C is use as hand sanitizer for hand hygiene when soap and water not available.

How Xycide C Works

The combination of Chlorhexidine Gluconate dan Isopropyl Alcohol provides rapid acting and antimicrobial activity against a wide range of microorganisms.

Before You Use Xycide C

When you must not use it

If you are allergic to any of the ingredients.

Before you start to use it

- Inform your doctor or pharmacist if you are taking or using any other medication at the same time.
- Check the seal is not broken before first use. If it is, do not use the solution.

Taking other medicines

Please consult your doctor or pharmacist if you are taking or have taken recently any other medicines.

How to Use Xycide C

How much to use

Apply 3-5 ml (1 pump) onto hand and rub until dry. After 10 applications, it is advisable to wash hands with soap and water.

When to use it

It is used when hands are not visibly soiled or when soap and water are not readily available.

How long to use it

You need to use Xycide C as directed by your doctor or pharmacist.

If you forget to take it

Reapply as you need.

If your use too much (overdose)

If anyone accidentally swallows some of the solution, talk to a pharmacist or doctor.

While You Are Using It

Things you must do

Do not use near eyes or mucous membranes, as it may cause irritation. If it does get into the eye or the mucous membranes, it should be washed quickly with plenty of water.

Things you must not do

Do not use on open skin wounds.

Things to be careful of

If any signs or symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness, and rapid heart rate develop, immediately stop using the product.

Side Effects

Immune system disorder
Frequency not known:
Hypersensitivity including anaphylactic shock.

Xycide C contains chlorhexidine. Inform your healthcare provider if you have a known allergy to chlorhexidine.

Stop using this product and seek immediate medical assistance if you experience rash, itching, swelling, breathing difficulties, light-headedness or rapid heartbeat.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers -> Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Xycide C

Storage

- Keep container tightly closed.
- Store below 30 °C.
- Protect from heat, flame and sunlight.
- Keep out of reach of children.
- Do not keep expired solution.

Disposal

Ask your pharmacist how to dispose of solution that no longer required.

Product Description

What it looks like

A light pink solution, slight alcohol and lavender odour.

Ingredients

Active Ingredient: Chlorhexidine Gluconate, Isopropyl Alcohol.

Excipient: Glycerin, carmoisine, lavender fragrance and purified water.

MAL Number

MALxxxxxxx

Manufacturer and Product Registration Holder

XORIX SDN BHD,
Lot 1, 2, 3 & 4,
Kawasan Industri Batu 7,
23000 Dungun,
Terengganu, MALAYSIA.

Revision Date

08/11/2022

Serial number

NPRA(RA)22/020