

PACKAGE INSERT

Product Name

NW Salicylic Acid 20%w/w Cream

Name and Strength of Active Substance(s)

Salicylic Acid.....20 %w/w

Contains ethanol10%w/w

Product Description

White smooth cream

Indication

For removal of warts.

Pharmacodynamics

The pharmacologic activity of this product is generally attributed to the keratolytic activity of salicylic acid, which is incorporated into a polyacrylic, film-forming virucidal solution designed to cover the wart without the need for a bandage, which results in mechanical removal of epidermal cells infected with wart viruses.

Pharmacokinetics

Salicylic acid may be percutaneously absorbed.

Recommended dosage

Apply daily and protect surrounding skin (eg with soft paraffin or specially designed plaster), may need to continue up to 3 months.

Contraindication

If you are allergic (hypersensitive) to any ingredient in NW Salicylic Acid 20%w/w Cream.

Warning and Precautions

For EXTERNAL USED only

Can cause convulsion.

Contraindicated in children below 2 years of age. Caution must be exercised when older children are treated. Avoid direct application into the nostrils.

It is dangerous to any camphor containing product into the nostrils of children. A small amount applied this way may cause immediate collapse.

Interaction with other medicaments

Salicylic Acid is known to interact with other drugs. Consult your doctor/pharmacist.

Statement on usage during Pregnancy and Lactation

Whilst there are no known contra-indications to the use of salicylic acid during pregnancy and lactation, the safety has not been established. Salicylic acid should therefore be used with caution or following professional advice.

Adverse effect/undesirable effect

Common side effects may include minor skin irritation, rash, or peeling; or changes in the color of treated skin (usually whitening). Severe allergic reactions (rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue); skin redness or irritation.

Stop using salicylic acid topical and call your doctor at once if you have: severe headache, ringing in your ears, a light-headed feeling, like you might pass out; shortness of breath; or severe burning, dryness, or irritation of the skin.

Overdose and Treatment

Symptoms of systemic salicylate poisoning (tinnitus, dizziness and deafness) have been reported after the application of salicylic acid to large areas of skin and for prolonged periods. Salicylism may also occur in the unlikely event of large quantities being ingested. Kindly contact your doctor/pharmacist for treatment.

Incompatibilities

No known incompatibilities.

Instructions for use and handling, and disposal

Store on the original packaging.

Any unused product or waste material should be discarded in accordance with local requirements. No additional environmental precautions for disposal are necessary.

Keep out of reach of children.

Storage conditions

Store below 30° C. Keep away from heat, moisture and direct light.

Dosage Forms and Packaging Available

15g White smooth cream in HDPE white container

Name and Address of Manufacturer/ Marketing Authorization Holder

Natural Wellness Industries Sdn. Bhd.

No: 78, Jalan Kilang Midah, Taman Midah,

56000, Cheras, Kuala Lumpur.

Date of Revision of Package Insert

21/06/2017