

QUALL ECZEMA CREAM

Betamethasone (as Betamethasone Dipropionate USP) (0.05% w/w)

What is in this leaflet

1. What Quall Eczema Cream is used for
2. How Quall Eczema Cream works
3. Before you use Quall Eczema Cream
4. How to use Quall Eczema Cream
5. While you are using Quall Eczema Cream
6. Side effects
7. Storage and Disposal of Quall Eczema Cream
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision

What Quall Eczema Cream is used for

Quall Eczema Cream is used to relief skin inflammation such as redness, swelling, and itching in certain skin conditions that respond to steroid treatment.

How Quall Eczema Cream works

This medicine contain Betamethasone dipropionate, a synthetic glucocorticoid. It works by reducing redness, inflammation, swelling, and irritation of the skin.

Before you use Quall Eczema Cream

- When you must not use it

Do not use this medicine if:

- you are allergic to betamethasone or any other ingredients of this medicine (listed in Product Description)

- Before you start to use it

Tell your pharmacist or doctor:

- Your medical conditions, allergies, and all medicines you are using.
- If you have any other skin infection.
- If you have diabetes. Steroid medicines may increase the glucose (sugar) level in your blood or urine. Your doctor may also need to adjust the dose of your diabetes medications.

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before using this medicine.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use Quall Eczema Cream

- How much to use

For topical use only.

Gently massage sufficient amount into the affected and surrounding skin areas twice a day, in the morning and evening.

- When to use it

Apply twice daily or as directed by your doctor or pharmacist.

Once your skin improves, you may not need to use it as often.

- How long to use it

Continue taking Quall Eczema Cream for as long as doctor recommends.

Ask your doctor or pharmacist before prolonged using.

- If you forget to use it

Apply the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to apply the medicine and skip the missed dose. Do not apply a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Using too much of this cream can allow it to be absorbed into the body and cause side effects such as weight gain, a rounder face, high blood sugar

(feeling thirstier or urinating more often), sugar in the urine, and temporary hormone imbalance where the body produces fewer natural steroids. If this happens, your doctor may reduce how often you use it, stop the treatment, or switch to a milder cream, and in some cases give additional medication if symptoms occur after stopping. These effects are usually reversible once the treatment is adjusted or discontinued under medical advice.

While you are using Quall Eczema Cream

- Things you must do

Wash your hands before and after applying the cream.

Clean and dry the affected area before applying the cream.

Take your medicine exactly as your doctor has told you.

Tell all the doctors and pharmacists treating you that you are taking Quall Eczema Cream.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Quall Eczema Cream is for external use only. Do not put the cream into your mouth.

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give this product to anyone else, even if they have the same symptoms or condition as you.

Do not use in or around the eyes.

Do not use more than the recommended dose unless your doctor tells you to.

Do not cover the treated skin area unless your doctor tells you to. Covering the skin may increase the amount of medicine absorbed into the

skin, which may lead to unwanted side effects.

Do not use on a child without a doctor's advice. Children can absorb larger amounts of this medication through the skin (due to the large skin surface area to body weight ratio) and may be more likely to have side effects. This medicine may also affect growth in children.

- **Things to be careful of**

Watch for any skin irritation, thinning, or worsening of your condition and inform your doctor if this happens. If your notice signs of infection such as redness, swelling, or pus, seek medical advice immediately.

Side effects

Like all medicines, this product can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

Possible side effects reported following use of topical corticosteroid includes skin thinning, stretch marks and acne.

Prolonged use on large surface area, on broken skin or under occlusive dressing might cause Cushing's syndrome.

Prolonged use in children may cause growth retardation, and delayed weight gain.

If you experience any symptoms, please consult your doctor as soon as possible.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Quall Eczema Cream

- **Storage**

Keep out of the reach and sight of children.

Do not freeze.

Store below 30°C.

- **Disposal**

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- **What it looks like**

A smooth white cream in pack of aluminium tube 15 g.

- **Ingredients**

- Active ingredient
 - Betamethasone Dipropionate
- Inactive ingredients
 - Chlorocresol
 - White Soft Paraffin
 - Cetomacrogol 1000
 - Sodium Acid Phosphate
 - Liquid Paraffin
 - Purified Water
 - Cetostearyl Alcohol
 - Sodium Hydroxide
 - Phosphoric Acid

- **MAL number:**

MAL11060001AZ

Manufacturer

TORRENT PHARMACEUTICALS LTD.,
Plot No. 810, Sector III,
Industrial Area, Pithampur,
District Dhar - 454 775 (M.P.)
India.

Product Registration Holder

Zulat Pharmacy Sdn Bhd
No. 23 & 23A, Jalan Bandar 3,
Taman Melawati, 53100
Kuala Lumpur,
Malaysia.

Date of revision

19/05/2026

Serial Number

NPRA(R4/1)090426/00015