Selsun 2.5% Suspension 120ml Carton • 64 x 37 x 135mmH • v3 • 21/7/20

135

MENTHOLATUM®

selsun®

2.5% SUSPENSION

MEDICALLY PROVEN

TREATMENT

FOR PROBLEM DANDRUFF, SCALP AND SKIN

SELENIUM SULFIDE 25MG/ML

120 mL

6.4

Controlled Medicine /
Ubat Terkawal

Store below 30ºC.

Route of administration:
Topical

Contains/Mengandungi :
DMDM Hydantoin, 0.5% w/v

Made in Thailand by:
INTERTHAI PHARMACEUTICAL MANUFACTURING LTD.
Bangkok, Thailand.

Product Registration Holder
Rohto-Mentholatum (M) Sdn Bhd (354691-U), Unit 9-1, Level 9, Wisma Averis, Avenue 5, Bangsar South City, No.8, Jalan Kerinchi, 59200 Kuala Lumpur.

Diimport oleh:
Zuellig Pharma Sdn Bhd (4013-D) No.15, Persiaran Pasak Bumi, Seksyen U8, Perindustrian Bukit Jelutong, 40150 Shah Alam, Selangor Darul Ehsan.

Imported by:
Zuellig Pharma Pte Ltd
15 Changi North Way #01-01,
Singapore 498770
Tel: 6546 8188

SINXXXXXX
MAL05023182ACZ

Batch No.
Mfg. Date
Exp. Date

3.7

MENTHOLATUM®

selsun®

2.5% SUSPENSION

MEDICALLY PROVEN

TREATMENT

FOR PROBLEM DANDRUFF, SCALP AND SKIN

SELENIUM SULFIDE 25MG/ML

120 mL

8 851520 060324

222

FOR TREATMENT OF SEBORRHOEIC DERMATITIS, TINEA VERSICOLOR, DANDRUFF AND AS AN ADJUNCT IN THE TREATMENT OF TINEA CAPITIS

SHAKE BOTTLE WELL

1. SCALP TREATMENT - Apply Selsun to wet hair, lather and leave on scalp for 3 minutes. Rinse. Repeat treatment then rinse thoroughly in running water. Use twice weekly at first, then as necessary or as doctor directs.

2. SKIN TREATMENT - Apply undiluted to affected areas for 10 minutes. Rinse thoroughly. Repeat daily for 1 week or as doctor directs.

WARNING - Do not use on broken or inflamed skin. Avoid contact with eyes.

CAUTION - For external use only, do not swallow and avoid contact with eyes (Elakkan daripada terkena mata). Keep out of reach of children. (Jauhi daripada pencapaian kanak-kanak). Do not use on broken or inflamed skin. (Selenium sulphide tidak boleh digunakan pada kulit yang pecah dan radang). Skin irritation or discoloration of hair may occur and can be minimized or avoided by rinsing thoroughly. If skin irritation persists, discontinue use. Not recommended for use by patients allergic to the ingredients, on infants and for treatment of tinea versicolor in pregnant women. Avoid use 48 hours before or after permanent waving, tinting or bleaching. Avoid contact with jewellery.

SYMPTOMS OF OVERDOSE AND TREATMENT - If swallowed, vomiting should be provoked or contact a doctor or poisons information centre.

R0240027
P1/10

222

selsun®

2.5% SUSPENSION

120 mL