

ORAGEL PAIN RELIEF

Choline Salicylate / Cetylpyridinium Chloride (9% w/w / 0.05% w/w)

What is in the leaflet

1. What Oragel Pain Relief is used for
2. How Oragel Pain Relief works
3. Before you use Oragel Pain Relief
4. How to use Oragel Pain Relief
5. While you are using it
6. Side Effects
7. Storage and Disposal of Oragel Pain Relief
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Oragel Pain Relief is used for

Oragel Pain Relief is used to relieve pain and swelling caused by ulcers, wounds or infections in the mouth. It is used for the healing of mouth ulcers, infant teething, denture irritation, sore gums, cold sores and discomfort due to braces in children.

How Oragel Pain Relief works

Oragel Pain Relief contains the active ingredients Choline Salicylate and Cetylpyridinium Chloride.

Choline Salicylate is an anti-inflammatory pain reliever agent. It is used to decrease swelling and to relieve pain and discomfort.

Cetylpyridinium Chloride is an antiseptic that works to kill bacteria and other microorganisms.

Before you use Oragel Pain Relief

– When you must not use it

Do not use Oragel Pain Relief if:

- you are allergic to salicylates or any other ingredients in this product (listed in Product Description).
- you are suffering from a stomach ulcer.
- you are taking medicine for blood clots or gout.
- patients are infants below 4 months.

– Before you start to use it

Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine. Salicylate appears in breast milk and crosses the placenta. Therefore, breast-feeding mothers should avoid using this product.

– Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines as below:

- Warfarin (blood thinner)
- Antacid (heartburn medicine)
- Corticosteroids

How to use Oragel Pain Relief

– How much to use

With clean finger or cotton bud, gently rub a small amount of gel (approximately one centimeter) onto the affected areas.

– When to use it

Adults:

Repeat the application every 3 hours.

Children:

Repeat the application every 3 hours. Do not exceed 6 applications in 24 hours.

– How long to use it

Use for the length of time advised by your doctor or pharmacist.

– If you forget to use it

If you forget to use Oragel Pain Relief, use it as soon as you remember. Never double up on the next dose to make up for the one missed.

– If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Using too much of this medication may cause dizziness, ringing in the ears, deafness, sweating, nausea and vomiting, headache, or confusion.

While you are using Oragel Pain Relief

– Things you must do

Tell all the doctors, dentists and pharmacists treating you that you are using Oragel Pain Relief.

Tell your doctor immediately if you become pregnant while using this medication.

– Things you must not do

Do not take or give aspirin to young children when using this product to

avoid any risk of excessive salicylate levels.

Do not use Oragel Pain Relief to treat any other conditions unless your doctor or pharmacist tells you so.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Oragel Pain Relief to anyone else, even if they have the same symptoms or condition as you.

– Things to be careful of

Do not apply more than once every 3 hours for infant teething pain.

Excessive and prolonged usage should be avoided, especially in infants.

Side Effects

Like all medicines, Oragel Pain Relief can cause side effects, although not everybody gets them. If they occur, most are likely to be minor and temporary. However, some may be serious and need medical attention.

If you think the gel is causing the baby any problems, please consult your doctor or pharmacist.

Stop taking the medicine if you experience: any symptoms of allergy (hypersensitivity).

If you get any side effects, even those not mentioned in this leaflet, tell your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Oragel Pain Relief

– Storage

- Store below 30°C.
- Protect from light.
- Keep cap tightly closed.
- Keep out of reach of children.

– Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

ORAGEL PAIN RELIEF

Choline Salicylate / Cetylpyridinium Chloride (9% w/w / 0.05% w/w)

Product Description

– *What it looks like*

Clear and colourless cooling gel with a pleasant taste. Alcohol free and sugar free.

– *Ingredients*

- Active ingredient
Choline Salicylate,
Cetylpyridinium Chloride.
- Inactive ingredients
Poloxamer 407, Sucralose,
Glycerin, Aniseed Oil, Menthol,
Purified Water.

– *MAL Number:*

MAL09082895X

Manufacturer and Product Registration Holder

Winwa Medical Sdn Bhd
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Date of Revision

03/02/2025

Serial Number

NPRA(R4/1)280125/00007