

ORAGEL PAIN RELIEF

(Analgesic - Anti-inflammatory - Antibacterial)

Active Ingredients:

Choline Salicylate 9%w/w
Cetylpyridinium Chloride 0.05%w/w

Presentations:

Tubes of 8g and 15g.

Product Descriptions:

A clear and colourless cooling gel with a pleasant taste. Alcohol free and sugar free.

Pharmacology:

Choline Salicylate has analgesic, anti-inflammatory and antipyretic properties. It alleviates pain by inhibition of prostaglandin synthesis. It acts as inhibitors of the enzyme cyclo-oxygenase, which results in the direct inhibition of the biosynthesis of prostaglandins and thromboxanes from arachidonic acid.

Cetylpyridinium Chloride is a quaternary pyridinium antiseptic with actions and uses typical of cationic surfactant which dissociates in aqueous solution into a smaller inactive anion and into a large and complex cation responsible for the surface activity. It also has bactericidal activity against gram positive bacteria and at a higher concentration, it is effective against some gram negative bacteria.

Choline Salicylate is a salicylic acid derivative. It is absorbed rapidly from the gastrointestinal tract when taken orally but absorption after rectal doses is less reliable. It can also be absorbed through the skin. Salicylate is excreted unchanged in the urine.

There is no significant absorption of Cetylpyridinium Chloride through the mucous membranes or skin.

Indications:

For the relief of pain and inflammation caused by ulceration, abrasion and oral infections. It is used for the healing of mouth ulcers, infant teething, denture irritation, sore gums, cold sores and discomfort due to braces in children.

Directions of Use:

Adults: With clean finger or cotton bud, rub gently a small amount of gel (approximately one centimeter) onto the affected areas. Repeat every 3 hours.

Children: With clean finger or cotton bud, rub gently a small amount of gel (approximately one centimeter) onto the affected areas. Repeat every 3 hours. Do not exceed 6 applications in 24 hour.

Symptoms and Treatments of Overdose:

Do not exceed the recommended dose. In case of overdose, seek medical advice immediately.

Choline Salicylate

Symptoms of overdosage include dizziness, tinnitus, deafness, sweating,

nausea and vomiting, headache and confusion. In acute oral Salicylate overdosage, repeated doses of activated charcoal may be given by mouth if the patient is suspected of ingesting more than 120mg/kg of Salicylate within 1 hour of presentation.

Cetylpyridinium Chloride

In the case of ingestion, this may cause nausea and vomiting. Treatment is symptomatic; demulcents and diluents may be given.

Contraindications:

Not recommended for infants below 4 months.

Do not use if have known hypersensitivity towards Choline Salicylate or Cetylpyridinium Chloride.

Do not take or give Aspirin to young children when using this product to avoid any risk of excessive Salicylate levels.

Do not use this product when taking medicine for blood clots or gout.

Do not use this product if patient has a stomach ulcer.

Precaution:

Do not apply more than once every 3 hour for infants teething pain.

Excessive and prolonged usage should be avoided, especially in infants.

If symptoms persist, please consult a doctor.

Use in Pregnancy and Lactation:

There are no data to confirm the safety of Choline Salicylate and Cetylpyridinium Chloride use in pregnancy and lactation. Consult your doctor before using this product if you are pregnant. Salicylate appears in breast milk and crosses the placenta. Therefore, breast-feeding mothers should avoid using this product.

Interactions with Other Medicaments:

Choline Salicylate may increase the toxicity of Warfarin.

When taking with antacid, it may reduce the Salicylate effects.

Choline Salicylate may increase the risk of gastrointestinal bleeding when used with corticosteroids.

There are no clinically significant interactions for Cetylpyridinium Chloride.

Undesirable Effects:

It is unlikely to occur if the product is used as directed. Sometimes, asthmatic or allergic reaction may occur if sensitive to Salicylate.

Storage:

Store below 30°C. Protect from light. Keep cap tightly closed after opening.

Keep out of reach of children. Jauhkan daripada capaian kanak-kanak.

Shelf Life:

3 years from the date of manufacture.

Manufacturer and Product Registration Holder

WINWA MEDICAL SDN. BHD.

1952, Lorong I.K.S. Bukit Minyak 3,

Taman I.K.S. Bukit Minyak,

14000 Bukit Mertajam,


Pulau Pinang, Malaysia.

MAL09082895X BRU15121981NP

Date of Revision: 07 August 2024

X.IOG-1.08

Size: 75mm x 110mm

 Dark Blue (Pantone 2767U)