

ANTISMINE SYRUP

Chlorpheniramine Maleate (2mg/5ml)

What is in the leaflet

1. What Antismine Syrup is used for
2. How Antismine Syrup works
3. Before you use Antismine Syrup
4. How to take Antismine Syrup
5. While you are using it
6. Side Effects
7. Storage and Disposal of Antismine Syrup
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Antismine Syrup is used for

Antismine Syrup is used for the treatment in allergic reactions including hay fever, runny or stuffy nose, sneezing, skin rash, itchy, watery eyes, itching of the nose or throat.

How Antismine Syrup works

Antismine Syrup contains chlorpheniramine maleate that relieves the allergic conditions by blocking the effects of histamine. Histamine is produced by the body during allergic reactions.

Before you use Antismine Syrup

– When you must not take it

Do not take if you have hypersensitivity to chlorpheniramine maleate, antihistamine or any other ingredients in this product.

Do not take if you are receiving MAOI (Monoamine Oxidase Inhibitor) therapy within the previous 14 days.

– Before you start to use it

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breastfeeding before using any medicine. Small amounts of antihistamines are secreted in breast milk and may affect breastfeeding.

Consult your doctor if you have:

- fits
- serious heart and blood vessels disease
- liver disease
- increased pressure in the eye
- urination retention
- prostate gland enlargement
- gastric outflow obstruction
- asthma
- respiratory inflammation
- overactive thyroid

- very high blood pressure

Special care should be taken when using chlorpheniramine maleate in children and the elderly as they are more prone to developing central nervous system effects.

– Taking other medicines

Tell your doctor if you are taking any other medicines as below:

- sedative or hypnotic
- opioid pain killer
- phenytoin, used to treat fits

How to take Antismine Syrup

– How much to take

Adults and children above 12 years:
Give 10ml 3 to 4 times a day. Do not exceed 60ml in a day.

Children 6 to 12 years:
Give 5ml 3 to 4 times a day. Do not exceed 30ml in a day.

Children 1 to 5 years:
Give 2.5ml to 5ml 3 times a day. Do not exceed 15ml in a day.

Children up to 1 year:
Give 2.5ml 2 times a day. Do not exceed 5ml in a day.

– When to take it

Follow the recommended dose and should not take more than the prescribed dose.

– How long to take it

Take as directed by your doctor or pharmacist.

– If you forget to take it

If you forget to take Antismine Syrup, take it as soon as you remember. Never double up on the next dose to make up for the one missed.

– If you take too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much of this medication may cause sedation, confusion or hallucination, fits, temporary inability to breathe, anticholinergic effects such as blurred vision, dry mouth and skin, irregular heartbeat.

While you are using Antismine Syrup

– Things you must do

If symptoms persist for more than 5 days, please consult a doctor or pharmacist.

– Things you must not do

Do not take Antismine Syrup to treat any other conditions unless your doctor or pharmacist tells you so.

– Things to be careful of

This medicine may cause drowsiness. If affected, do not drive or operate machinery. Avoid alcoholic drinks.

Side Effects

The most common side effects include drowsiness, gastrointestinal upset such as nausea, vomiting, diarrhoea or constipation, and epigastric pain, tiredness, dizziness, irregular heartbeat and allergic reactions.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Antismine Syrup

– Storage

- Store below 30°C.
- Protect from light.
- Keep cap tightly closed.
- Keep out of reach of children.

– Disposal

Do not take this after the expiry date. An expiry date is printed on the label and the box of the medicine.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

– What it looks like

Clear, greenish yellow coloured syrupy liquid with fruity flavor. Alcohol free.

– Ingredients

- Active ingredient
Chlorpheniramine Maleate

ANTISMINE SYRUP

Chlorpheniramine Maleate (2mg/5ml)

- Inactive ingredients
Methyl Hydroxybenzoate, Propyl Hydroxybenzoate, Sucrose, Xanthan Gum, Liquid Glucose, Sorbitol Solution 70%, Sodium Citrate Dihydrate, Strawberry Flavour, Tutti Frutti Flavour, Quinoline Yellow, Purified Water.

– MAL Number:

MAL19900149AZ

**Manufacturer and Product
Registration Holder**

Winwa Medical Sdn Bhd
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Date of revision

13/01/2025

Serial Number:

NPRA(R4/1)200624/00075